

Product – Chemical Profile for Treatments Containing Perfluoroalkyl or Polyfluoroalkyl Substances for Use on Converted Textiles or Leathers

NOVEMBER 2019 • DISCUSSION DRAFT



TABLE OF CONTENTS

1. About this Profile	4
2. Rationale for Product-Chemical Selection.....	6
Potential for exposure to PFASs in treatments for converted textiles or leathers	7
Potential for significant or widespread adverse impacts.....	8
Adverse impacts relevant to this Priority Product.....	9
3. Product Definition and Scope	10
4. Candidate Chemical Definition and Properties	11
Relevant physicochemical properties.....	15
Environmental fate.....	21
Degradation, reaction, or metabolic products of concern.....	30
Hazard Traits and Environmental or Toxicological Endpoints.....	32
Structural or mechanistic similarity to chemicals with known adverse impacts.....	38
5. Potential for Exposures to the Candidate Chemical in the Priority Product.....	39
Market presence and trends	39
Potential exposures to the Candidate Chemical during the product’s life cycle.....	40
Aggregate effects	45
Indicators of potential exposures to the Candidate Chemical in the product	46
6. Potential for Significant or Widespread Adverse Impacts	65
Adverse impacts linked to the Candidate Chemical’s hazard traits	66
Cumulative effects.....	68
Adverse waste and end-of-life effects	70
Populations that may be adversely impacted	72
7. Other Regulatory Programs	78
8. Potential Alternatives	80
9. Additional Considerations	82

10. Conclusions	87
Acronyms and Abbreviations	88
PFAS (sub)classes	88
Individual PFASs	89
Other abbreviations used in this document	90
Units.....	91
References	92
Appendix 1 – Additional PFAS Candidate Chemicals Database Listings.....	148
Appendix 2 – PFAS Hazard Traits	157
Appendix 3 – Report Preparation.....	172
Appendix 4 – Appendix References.....	173

1. ABOUT THIS PROFILE

The Department of Toxic Substances Control (DTSC) identifies product-chemical combinations for consideration as Priority Products in accordance with the process identified in Article 3 of the Safer Consumer Products (SCP) regulations.¹ DTSC has determined that treatments containing any member of the class of perfluoroalkyl and polyfluoroalkyl substances (PFASs) for use on converted² textiles or leathers meet the key prioritization criteria³ for listing a Priority Product:

(1) There must be potential public and/or aquatic, avian, or terrestrial animal or plant organism exposure to the Candidate Chemical(s) in the product; and

(2) There must be the potential for one or more exposures to contribute to or cause significant or widespread adverse impacts.

This Product-Chemical Profile (Profile) demonstrates that the regulatory criteria have been met and serving as the basis for Priority Product rulemaking. The Profile does not provide a comprehensive assessment of all available adverse impact and exposure literature on perfluoroalkyl and polyfluoroalkyl substances (PFASs) or on treatments for converted textiles or leathers. If this Priority Product regulation is adopted, the responsible entities must follow the reporting requirements pursuant to the SCP regulations.⁴

Candidate Chemical: A chemical that exhibits a hazard trait and is listed on one or more authoritative lists in the SCP regulations.

Product-Chemical Profile: A report generated by DTSC to explain its determination that a proposed Priority Product meets the SCP regulatory criteria for potential significant or widespread adverse impacts to humans or the environment.

Priority Product: A product-chemical combination as identified in regulation by DTSC that has the potential to contribute to significant or widespread adverse impacts to humans or the environment.

¹ CAL. CODE REGS. tit. 22, Division 4.5, Chapter 55, Article 3: Process for Identifying and Prioritizing Product-Chemical Combinations.

² The term “converted” indicates textiles and leather that manufacturers and craftspeople have turned into consumer products such as clothing, shoes, upholstery, etc.

³ CAL. CODE REGS. tit. 22, § 69503.2(a).

⁴ CAL. CODE REGS. tit. 22, § 69503.7 and Article 5 (Alternatives Analysis)

Readers should consider the following:

- This Profile is not a regulatory document and does not impose any regulatory requirements.
- The Profile summarizes information compiled by DTSC as of December 2019.
- DTSC requests that stakeholders provide data on the chemical and product described in this document to assist us in the evaluation process that may lead to our regulatory proposal. Written comments can be submitted using our information management system, CalSAFER,⁵ prior to 11:59 p.m. on December 31, 2019.
- By proposing to list this product-chemical combination as a Priority Product containing a Chemical of Concern, DTSC is not asserting that the product cannot be used safely. The proposal indicates only that there is a potential for people or the environment to be exposed to the Chemical of Concern in the Priority Product; that such exposure has the potential to cause or contribute to significant or widespread adverse impacts; and that safer alternatives should be explored.

⁵ <https://calsafes.dtsc.ca.gov/workflows/comment/>

2. RATIONALE FOR PRODUCT-CHEMICAL SELECTION

The Department of Toxic Substances Control (DTSC) proposes to list treatments containing any member of the class of perfluoroalkyl and polyfluoroalkyl substances (PFASs) for use on converted textiles or leathers as a Priority Product. This product-chemical combination meets the identification and prioritization factors outlined in the Safer Consumer Products regulations: (1) there is potential for human and other organism exposure to PFASs from treatments for converted textiles or leathers; and (2) the exposure has the potential to contribute to or cause significant and widespread adverse impacts.

The class of PFASs contains more than 3,000 man-made chemicals with at least one fully fluorinated carbon atom. All PFASs are [Candidate Chemicals](#) because the California Environmental Contaminant Biomonitoring Program lists the entire class as Priority Chemicals for measuring in the blood or urine of Californians.

DTSC is also taking a class approach to PFASs because:

- PFASs are a wide and varied group of chemicals used in many applications. All PFASs share one common trait – highly stable carbon-fluorine (C-F) bonds that make them or their final degradation products highly persistent in the environment.
- Persistence is a hazard trait identified in the Office of Environmental Health Hazard Assessment’s Green Chemistry Hazard Traits regulation 69405.3: “Persistence of a chemical in the environment promotes sustained exposure and contributes to accumulation in the environment.”
- Within the overall PFAS class, perfluoroalkyl acids (PFAAs) are the most problematic subclass. They are the most widely and thoroughly characterized subclass and are associated with a number of different health hazards, including endocrine disruption, developmental and reproductive toxicity, and immune dysregulation.
- In the case of PFAAs, their ubiquity in combination with their persistence results in continuous exposure from multiple sources, including contaminated drinking water sources and the food supply. They can accumulate in meat, plants, and drinking water, and ultimately in humans and wildlife.
- The vast majority of PFASs degrade into PFAAs, which typically increases their mobility in the environment, making containment or removal a challenge.
- While persistence alone warrants enough concern to include any member of the PFAS class in product prioritization, the health hazards associated with exposure to PFAAs are additionally concerning and underlie our listing.

Potential for exposure to PFASs in treatments for converted textiles or leathers

PFASs have a wide variety of applications, including in numerous consumer products. Once released to the environment during product manufacture, use, or disposal, PFASs become part of a virtually closed cycle leading to chronic human and ecological exposures. They are found in the aquatic, atmospheric, and terrestrial environments, including remote locations far from any point source. They are present in indoor dust, drinking water sources, food, wildlife, and humans – including the milk and serum of breastfeeding women. Because persistent PFASs lack a natural degradation route, their levels in the environment, humans, or biota will continue to rise for as long as PFASs are produced and used in consumer products, and even after production of these compounds has ceased.

Once released to the environment during product manufacture, use, or disposal, PFASs become part of a virtually closed cycle leading to chronic human and ecological exposures. Treatments for converted textiles or leathers contribute to the widespread environmental contamination and exposures.

DTSC has identified treatments for converted textiles or leathers as significant sources of human and ecological PFAS exposures, especially via inhalation during product use. Treatments for converted textiles or leathers are frequently used in homes and in several occupations. They contribute to the ubiquitous environmental contamination and exposures, as do other consumer products such as food packaging, cosmetics, and waterproof clothing.

Human exposure to PFASs begins early in a person's life, since mothers transfer these chemicals to their babies via the placenta and breastfeeding. In the general population, PFAS exposure occurs mainly via ingestion of contaminated food and drinking water, but other sources of exposure may contribute. This contamination is partly due to releases of PFASs from the manufacturing, use, and disposal of treatments for converted textiles or leathers, as well as from the use and disposal of the treated consumer products. When used, treatments for converted textiles or leathers release PFASs into indoor air and dust, which people inhale or ingest. Most waste or end-of-life converted textiles or leathers in California are disposed of in landfills, where they become sources of PFASs to the environment via leachates and gaseous emissions. Wastewater treatment plants that collect landfill leachates, surface runoff, and residential and commercial wastewater do not effectively remove PFASs. As a result, when wastewater effluent is discharged into surface waters, PFASs are released into the environment, contaminating aquatic ecosystems and drinking water sources. Sewage sludge also contains PFASs, thus the application of biosolids on soil can contaminate terrestrial ecosystems, drinking water, and human food supplies.

Carpets, rugs, upholstery, clothing, shoes, and other consumer products to which treatments containing PFASs have been applied become major sources of exposure for infants and children via direct contact and incidental indoor dust ingestion. Young children have been shown to ingest more soil and dust than adults, due to greater hand-to-mouth transfer; this can result in higher exposure to PFASs found in these contaminated environmental media. Carpet and upholstery cleaners, workers in stores selling upholstered furniture, furnishings, carpets, clothing, or shoes, and auto dealership workers and auto detailing technicians may also experience above-average PFAS exposure levels.

Potential for significant or widespread adverse impacts

Due to the strength of the carbon-fluorine bond, PFASs are characterized by high environmental persistence, which leads to continuous and poorly reversible accumulation in the environment, and hence to likely increasing exposures. Many PFASs display significant mobility in environmental media, which makes them widespread in the environment and in living organisms. Some members of the PFAS class bioaccumulate significantly in animals or plants, including in foods consumed by humans, and undergo lactational or transplacental transfer from mothers to offspring. Certain PFASs also contribute to global warming.

There is evidence that exposure to PFASs can lead to adverse health outcomes in humans. If humans are exposed to PFASs through diet, drinking water or inhalation, these chemicals remain in the body for a long time. As people continue to be exposed to PFASs the levels in their bodies may increase to the point where they suffer from adverse health effects. Studies indicate that some PFASs can cause reproductive and developmental, liver and kidney, and immunological effects, as well as tumors in laboratory animals. The most consistent findings from human epidemiology studies are a small increase in serum cholesterol levels among exposed populations, with more limited findings related to infant birth weights, effects on the immune system, cancer, and thyroid hormone disruption. Some PFASs have also been linked to phytotoxicity, aquatic toxicity, and terrestrial ecotoxicity.

Under a voluntary agreement with the U.S. Environmental Protection Agency (U.S. EPA), in 2015, major PFAS manufacturers phased out the production and emission of longer-chain PFASs with known adverse health impacts and their precursors. As a result, biomonitoring studies have shown decreasing levels of certain longer-chain PFASs in human tissues. However, most replacements are also PFASs, including fluorinated ethers and shorter-chain PFASs. Fluorinated ethers were thought to degrade easily, but recent studies found they also persist indefinitely in the environment. Shorter-chain PFASs are marketed as less toxic compared to the longer chains, mainly because they appear to bioaccumulate less and to be more readily eliminated from some organisms. Nevertheless, they are equally persistent and more mobile in the environment than the chemicals they are replacing, and also show potential for toxicity. Toxicological and epidemiological data clearly indicating the safety of aggregate, chronic, and low-dose exposures to PFASs found in stain and soil repellents are lacking.

All PFASs are Candidate Chemicals under the Safer Consumer Products (SCP) Program due to their designation on December 22, 2015, as Priority Chemicals under the California Environmental Contaminant Bio-monitoring Program.

Based on the criteria in the Safer Consumer Products regulations, DTSC has determined that treatments containing PFASs for use on converted textiles or leathers have the potential to cause significant and widespread adverse impacts to sensitive subpopulations, including fetuses, infants, young children, pregnant women, carpet and upholstery cleaners, workers in stores selling upholstered furniture, furnishings, carpets, clothing, or shoes, and auto dealership workers and auto detailing technicians; to environmentally sensitive habitats; and to threatened and endangered species. Given the known hazard traits, replacing currently used PFASs in the treatments for converted textiles or leathers with other members of the PFAS class could constitute a regrettable substitution. Hence, this proposal covers treatments containing any member of the class of PFASs intended for use on converted textiles or leathers.

Adverse impacts relevant to this Priority Product

Relevant Factors are used in SCP's Alternatives Analysis (AA) to make a focused and meaningful comparison of adverse impacts during the product's life cycle between the Priority Product and the alternative. This Profile has identified adverse impacts in the following categories:

- Adverse environmental impacts
- Adverse public health impacts
- Adverse waste and end-of-life effects
- Environmental fate
- Physicochemical properties
- Associated exposure pathways and life cycle segments, including:
 - Manufacture
 - Use
 - Operation and maintenance
 - Waste generation and management
 - Reuse and recycling
 - End-of-life disposal

3. PRODUCT DEFINITION AND SCOPE

This section describes the product that forms the basis for the proposed product-chemical combination.

“Treatments containing PFASs for use on converted⁶ textiles or leathers” means any product containing PFASs placed into commerce in California that may be marketed or sold for the purpose of:

- eliminating dirt or stains from carpets, rugs, clothing, shoes, upholstery, or other converted fabrics; or
- repelling stains, dirt, oil, or water from carpets, rugs, clothing, shoes, upholstery, or other converted fabrics.

Note that these treatments are sometimes also referred to as impregnating agents (Jensen et al. 2008).

Subproducts include:

- cleaner: a product marketed or sold for the purpose of eliminating dirt or stains;
- protectant: a product marketed or sold to protect the surface from soiling when in contact with dirt or other impurities, or to reduce liquid absorption;
- spot remover: a product marketed or sold to clean localized areas, or to remove localized spots or stains; and
- water proofer or water repellent: a product marketed or sold to repel water.

“Treatments containing PFASs for use on converted textiles or leathers” does not include products marketed or sold exclusively for use at industrial facilities during the process of carpet, rug, clothing, shoe, or furniture manufacturing.

⁶ The term “converted” indicates textiles and leather that manufacturers and craftspeople have turned into consumer products such as carpets, upholstery, furnishings, clothing, shoes, etc.

4. CANDIDATE CHEMICAL DEFINITION AND PROPERTIES

This section introduces the Candidate Chemical(s) in the proposed product-chemical combination.

This proposal covers all perfluoroalkyl and polyfluoroalkyl substances (PFASs)⁷ in current or future production. All PFASs are Candidate Chemicals (DTSC 2016) under the Safer Consumer Products (SCP) Program due to their designation on December 22, 2015, as Priority Chemicals under the California Environmental Contaminant Biomonitoring Program (California Biomonitoring Program or CECBP). The CECBP listing is based on the chemicals' potential for widespread exposures, persistence, bioaccumulation, and emerging evidence for toxicity. In addition, DTSC's Candidate Chemicals database includes 73 other listings for individual PFASs (see Appendix 1).

PFASs are a class of fluorinated organic chemicals containing at least one fully fluorinated carbon atom (CECBP 2015a). This class includes more than 3,000 chemicals (KEMI 2015), although fewer may be in current production. The Organisation for Economic Cooperation and Development (OECD) has identified 4,730 Chemical Abstracts Service (CAS) numbers related to individual PFASs or commercial PFAS mixtures available on the global market (OECD 2018). PFASs are manufactured and not found naturally in the environment. All PFASs contain carbon and fluorine atoms, in addition to other elements such as oxygen, hydrogen, nitrogen, sulfur, or chlorine. PFASs vary in chain length, i.e., the number of carbon atoms forming the backbone of their molecule, from a chain of two carbons to large molecular-weight polymers.

Based on common characteristics, and for the purposes of this designation, PFASs can be roughly subdivided into four main categories (Buck et al. 2011; Wang et al. 2017b) listed below and depicted in Figure 1. However, any other PFAS that exists or will be developed in the future and is used in carpets and rugs also falls under the scope of this proposal:

- 1. Perfluoroalkyl acids (PFAAs).** These are perfluorinated substances in which fluorine atoms have replaced all hydrogen atoms attached to carbon atoms (except for those associated with functional groups). As a result, these compounds are recalcitrant to degradation and extremely persistent in the environment. This subgroup includes:
 - Perfluoroalkyl carboxylic acids (PFCAs) such as perfluorooctanoate (PFOA);⁸
 - Perfluoroalkyl sulfonic acids (PFSAs) such as perfluorooctane sulfonate (PFOS);⁸
 - Perfluoroalkyl sulfinic acids (PFSiAs);
 - Perfluoroalkyl phosphonic acids (PFPAs);

⁷ Note that U.S. EPA and others use the acronym "PFAS" for this class of chemicals. For the purpose of this document, we use the acronym "PFASs" to emphasize that it refers to a class of chemicals, and to make it clear when we refer to multiple members of this class rather than an individual chemical.

⁸ PFOA and PFOS are referred to as "C8" because they contain eight carbon atoms in their molecules.

- Perfluoroalkyl phosphinic acids (PFPIAs);
- Perfluoroether carboxylic acids (PFECAs); and
- Perfluoroether sulfonic acids (PFESAs).


2. PFAA precursors. These are mostly polyfluoroalkyl substances, meaning fluorine atoms have replaced all hydrogen atoms attached to at least one (but not all) carbon atoms. Polyfluorinated substances have the potential to degrade into perfluoroalkyl substances, i.e., they are precursors to perfluoroalkyl substances. Examples include:

- PFASs used for surface treatments:
 - Side-chain fluorinated polymers (e.g., fluorinated acrylate, methacrylate, urethane, oxetane polymers);
 - n:2⁹ polyfluoroalkyl phosphoric acid esters, also known as polyfluoroalkyl phosphates, or fluorotelomer phosphates (PAPs);
 - Polyfluorinated organosiloxanes; and
 - Fluorotelomer ethoxylates (FTEOs).
- PFASs commonly used as raw materials for surface treatment products:
 - *N*-alkyl perfluoroalkane sulfonamides (FASAs);
 - Perfluoroalkane sulfonamide ethanols (FASEs) and *N*-alkyl perfluoroalkane sulfonamidoethanols;
 - *N*-alkyl perfluoroalkane sulfonamidoethyl acrylates and methacrylates (FAS(M)ACs);
 - Fluorotelomer iodides (n:2 FTIs);
 - Fluorotelomer olefins (n:2 FTOs);
 - Fluorotelomer alcohols (n:2 FTOHs); and
 - Fluorotelomer acrylates (n:2 FTACs) and methacrylates (n:2 FTMACs).
- PFASs formed as intermediate environmental transformation products:
 - Fluorotelomer aldehydes (n:2 FTALs) and unsaturated aldehydes (n:2 FTUALs);
 - Fluorotelomer carboxylic acids (n:2 FTCAs) and unsaturated carboxylic acids (n:2 FTUCAs);
 - Fluorotelomer sulfonic acids (n:2 FTSs);
 - Fluorotelomer ethoxycarboxylates (FTEOCs); and
 - n:3 saturated and unsaturated acids.

⁹ “N” refers to the number of perfluorinated carbon atoms; the number after the colon refers to the number of nonfluorinated carbon atoms in the molecule.

- 3. Perfluoropolyethers (PFPEs).** These are perfluoroalkyl substances that are highly persistent in the environment, with large molecular size (oligomers, polymers, and copolymers), and ether linkages. They are unlikely to degrade to PFAAs under typical environmental conditions, but may be manufactured using PFAAs and contain them as impurities, and may release PFAAs during combustion.
- 4. Fluoropolymers.** These polymers are materials (as opposed to surface treatments), and are highly persistent in the environment. They cannot degrade to PFAAs under typical environmental conditions, but certain PFAAs have been used in their manufacturing and can occur as impurities. Moreover, fluoropolymers may release PFCAs, including PFOA, during combustion at temperatures between 180 and 800°C (Feng et al. 2015; Schlummer et al. 2015). Examples of fluoropolymers include polytetrafluoroethylene (PTFE), polyvinylidene fluoride (PVDF), and polyvinyl fluoride (PVF).

Thus, while PFAAs constitute a small subset (approximately one percent) of PFASs, they are terminal degradation products, manufacturing aids/feedstocks, or impurities of other PFAS class members, which makes their hazard traits relevant to the entire class. PFAAs and some of their precursors are frequently subdivided into longer- and shorter-chain PFASs. The longer-chain PFASs have six or more perfluorinated carbons; longer-chain PFCAs, PFPAs, and PFPiAs have seven or more perfluorinated carbons (Bowman 2017).



* PFASs in RED are those that have been restricted under national/regional/global regulatory or voluntary frameworks, with or without specific exemptions (for details, see OECD (2015), Risk reduction approaches for PFASs. <http://oe.cd/hAN>).

** The numbers of articles (related to all aspects of research) were retrieved from SciFinder® on Nov. 1, 2016.

Figure 1: General classification of PFASs. Reprinted (adapted) with permission from Environ. Sci. Technol. 51, 5, 2508-2518 (Wang et al. 2017b). Copyright 2017 American Chemical Society.

Relevant physicochemical properties

Reference: CAL. CODE REGS. tit. 22, § 69503.3(a)(1)(D).

Physicochemical properties can be helpful in predicting a chemical's behavior. A chemical's behavior in humans, wildlife, ecosystems, and the environment may indicate potential adverse public health and environmental impacts.

PFASs possess unique physicochemical characteristics that confer increased stability in extreme temperatures, antistatic properties, and resistance to wettability, staining, and corrosion (Buck et al. 2011, 2012). PFASs are used in a wide variety of applications, including (KEMI 2015; FluoroCouncil 2019).

- Surface treatments for leather and textiles
- Surface treatments for food packaging, processing, and preparation
- Metal (chromium) plating fume suppressants
- Detergents, pharmaceuticals, pesticides
- Aqueous firefighting foams (AFFFs)
- Aerospace, automotive, construction, and electronics manufacturing industries

These unique properties of PFASs also contribute to their environmental persistence, global distribution, and accumulation in water, soils, plants, and animals. Compared to other organic chemicals, PFASs are very persistent in the environment and biota, partly because fluorine (F) forms the strongest single bond with carbon (Kiplinger et al. 1994). Due to its high electronegativity, the F atom pulls the shared electrons of the covalent C-F bond towards itself, which renders it partially negative and the C atom partially positive, creating a polar covalent bond. This bond requires high energy, up to 544 kJ/mol, to dissociate (Lemal 2004), compared to the 337 kJ/mol needed to separate carbon and hydrogen (Dean 1999). Consequently, PFASs are thermally, chemically, and biologically stable, and resistant to abiotic degradation such as atmospheric photo-oxidation, direct photolysis, and hydrolysis (Wang et al. 2016). Perfluoroethers are more thermally stable than other PFASs due to their strong C-O bonds (Ceretta 2013). Polymeric PFASs also have high thermal, chemical, aging, and weather resistance, and are inert to solvents, hydrocarbons, acids and bases (Banks et al. 1994). Even though bacteria and fungi can degrade organic chemicals by utilizing the carbon present in the compounds as an energy source, the F atoms in PFASs surround and shield the carbon backbone, so organisms are largely unable to fully biodegrade these chemicals (Frömel and Knepper 2010). PFASs that can degrade in the environment typically transform into smaller PFASs that do not degrade further under environmental conditions, such as PFAAs (Buck et al. 2011; D'eon and Mabury 2011; Frömel and Knepper 2010; Krafft and Riess 2015a; Wang et al. 2014a; Washington et al. 2015).

For most PFASs, physicochemical properties data are absent from the public domain. The data available in the literature can range widely depending on the measurement methods or model used for

the estimates. Table 1 lists data available for PFAAs. Less is known about PFECAs and PFESAs, but despite structural differences, their physicochemical properties are believed to be similar to those of their PFCA or PFSA counterparts (Gomis et al. 2015). In particular, they appear to be similarly persistent and mobile in the environment as the longer-chain PFASs (Gomis et al. 2015).

Most PFASs are solid at room temperature. Notable exceptions include 6:2 FTOH and 4:2 FTOH, which exist as a liquid at room temperature, and 8:2 FTOH, which exists as a solid but can sublime and volatilize at room temperature (Concawe 2016). The boiling points of PFASs are similar to those of the homologous hydrocarbons (i.e., molecules with the same structure but only carbon-hydrogen bonds instead of carbon-fluorine bonds), except for perfluoroethers and perfluoroketones, which boil at significantly lower temperatures compared to their hydrocarbon homologues due to lower intermolecular forces (Ceretta 2013).

Most PFASs exhibit low vapor pressures, but compared to homologous hydrocarbons they tend to be similarly or somewhat more volatile, despite higher molecular weights (Krafft and Riess 2015a). Most PFAAs are semi-volatile and can adsorb onto indoor dust (Bohlin-Nizzetto et al. 2015; Dreyer et al. 2015; Ericson et al. 2012; Fraser et al. 2013; Haug et al. 2011; Knobeloch et al. 2012; Liu et al. 2014, 2015a; Strynar and Lindstrom 2008). Some of their precursors however, including FTUCAs and FTOHs, tend to be more volatile and can undergo atmospheric long-range transport (Ding and Peijnenburg 2013). In general, shorter-chain PFASs are more volatile than longer chains, and thus more likely to be released from products (U.S. EPA 2017a).

PFAAs are proteinophilic (protein-binding), accumulating particularly in blood, liver, stomach, kidney, lung, gall bladder, brain, muscle, and yolk sac tissues.

PFASs display a range of solubility in water, from sparingly soluble to miscible, though models may significantly underestimate PFAS solubility (Xiao 2017). Solubility tends to increase with decreasing carbon chain length and molecular weight (Table 1). PFCAs and PFSAAs are relatively more soluble than other PFASs of similar chain length due to the hydrophilicity of the carboxylate and sulfonate groups on these molecules. Solubility of PFCAs and PFSAAs tends to decrease with increasing molecular weight due to a concomitant increase in the length of the hydrophobic perfluorinated alkyl chains (Concawe 2016). Their water solubility makes these PFASs mobile in the environment and capable of undergoing long-range transport via ocean currents (Ahrens et al. 2009a; Armitage et al. 2009; Krafft and Riess 2015b; Prevedouros et al. 2006). Model results suggest, however, that PFPiAs are highly hydrophobic; therefore, sediments may be an important sink for PFPiAs in the aquatic environment (Xiao 2017). PFAAs ionize in water at environmentally relevant pH, according to their negative pK_a (Gomis et al. 2015; Wang et al. 2017b). For PFPAs, full deprotonation (i.e., two negative charges) occurs at a pH of 5.13-7.11 (Xiao 2017).

The bonds between the carbon-fluorine (CF_3 and CF_2) groups present in PFASs create very low critical surface energy, resulting in lipophobic and hydrophobic characteristics. This means PFASs repel both oil and water (Audenaert et al. 1999), making them surface active agents (surfactants) useful for resisting stains from soils, liquids and grease on the surface of carpets, rugs, and other textiles. Due to their low critical surface energy, PFASs tend to accumulate and form micelles at liquid (e.g., water)/air phase boundaries (Concawe 2016; Eftaiha et al. 2012; Mason Chemical Company 2017).

Their hydrophilic and hydrophobic properties pose challenges to determining PFAS partitioning coefficients experimentally, leading to reliance on modeled data and estimates (Shoeib et al. 2004). For example, K_{ow} , which represents a chemical's partitioning behavior between octanol and water, cannot be determined experimentally for PFASs because they form a third phase in addition to octanol and water. All PFAAs listed in Table 1 except for PFBA, PFPeA, PFBS, PFBPA, and PFHxPA have modeled $\log K_{ow} > 4$, which means they are considered bioaccumulative according to the California Code of Regulations, Title 22, Division 4.5, Chapter 54, Article 5. However, K_{ow} , a common screening criterion of a neutral compound's ability to partition from water into lipid-rich tissues within an organism (Mackay and Fraser 2000), has limited applicability to PFAAs, due to their unique properties. This is because, in general, PFASs have a low potential to accumulate in adipose (lipid) tissues, unlike most other persistent organic pollutants. PFAAs are proteinophilic (protein-binding), accumulating particularly in blood, liver, stomach, kidney, lung, gall bladder, brain, muscle, and yolk sac tissues (Chen and Guo 2009; Hebert and MacManus-Spencer 2010; Greaves et al. 2012; Jones et al. 2003; Mackay and Fraser 2000; Pérez et al. 2013). Additionally, the concept of K_{ow} for PFAAs has little to no practical meaning or usefulness because PFAAs exist in anion rather than neutral form at environmentally relevant conditions (Valsecchi et al. 2017).

Data on the physicochemical properties of PFASs are limited to mostly model results, which vary widely because the underlying data set is based on hydrocarbons. Models and experimental data that can adequately characterize PFASs, especially beyond longer-chain PFCAs and PFSAs, are still lacking (Xiao 2017).

Table 1. Physicochemical properties of some PFAAs¹⁰

Chemical Name	Water Solubility (20 – 25 °C) [g/L]	Melting Point [°C]	Boiling Point [°C]	Vapor Pressure [Pa]	log K _{ow} [-]	log K _{oc} [L/kg]	K _d (pH 7)	Dissociation Constant (pK _a)	Soil Adsorption Coefficient (L/kg)
Perfluorobutanoic acid (PFBA)	Miscible	-18	121	1307	2.8	1.9	--	-0.2 to 0.7	48*
Perfluoropentanoic acid (PFPeA)	113	14*	124	1057	3.4	1.4	--	-0.06	96*
Perfluorohexanoic acid (PFHxA)	22	14	143	457	4.1	1.9	--	-0.13	1,070*
Perfluoroheptanoic acid (PFHpA)	4.2	30	175	158	4.7	2.2	0.4 – 1.1	-0.15	2,110*
Perfluorooctanoic acid (PFOA)	3.4 – 9.5	37 – 60	188 – 192	4 – 1300	5.3	1.3 – 2.4	0 – 3.4	-0.16 to 3.8	1,160*
Perfluorononanoic acid (PFNA)	9.5	59 – 66	218	1.3	5.9	2.4	2.6 – 5.9	-0.17	2,830*
Perfluorodecanoic acid (PFDA)	9.5	77 – 88	218	0.2	6.5	2.8	2.0 – 31	-0.17	397*
Perfluoroundecanoic acid (PFUnA)	0.004	83 – 101	160 – 230	0.1	7.2	3.3	12 – 103	-0.17	16,700*

¹⁰ Data excluding those marked with an asterisk are from peer-reviewed literature as summarized by Concawe (2016) Environmental fate and effects of poly and perfluoroalkyl substances (PFAS). Concawe Soil and Groundwater Taskforce, Brussels, Belgium. Prepared by ARCADIS: T. Pancras, G. Schrauwen, T. Held, K. Baker, I. Ross, H. Slenders, Report # 8/16. Accessed March 15, 2017. Concawe (www.concawe.eu) is a division of the European Petroleum Refiners Association based in Brussels, Belgium. Entries marked with an asterisk (*) were predicted using QSAR models by U.S. EPA (2017a) U.S. Environmental Protection Agency (U.S. EPA): Chemistry Dashboard. Available at: <https://comptox.epa.gov/dashboard>. Data are calculated and updated regularly by OPERA [OPEN (quantitative) Structure-activity Relationship Application], a standalone free and open source command line application in Matlab (Version 8.2) providing QSAR models predictions. Model validation data set may be found here: <http://esc.syrres.com/interkow/EpiSuiteData.htm>. Accessed November 6, 2017.

Perfluorododecanoic acid (PFDoA)	0.0007	107 – 109	245	0.01	7.8	--	24 – 269	-0.17 to 0.8	85,500*
Perfluorotridecanoic acid (PFTTrDA)	0.0002	99*	249*	0.3	8.3	--	--	--	184,000*
Perfluorotetradecanoic acid (PFTeDA)	0.00003	98*	276	0.1	8.9	--	--	--	233,000*
Perfluorobutane sulfonic acid (PFBS)	46 – 57	76 – 84	211	631	3.9	1	--	-6.0 to -5.0	288*
Perfluorohexane sulfonic acid (PFHxS)	2.3	58*	231*	59	5.2	1.8	0.6 – 3.2	-6.0 to -5.0	2,300*
Perfluorooctane sulfonic acid (PFOS)	0.52 – 0.57	54	237* to > 400	6.7	6.4	2.5 – 3.1	0.1 – 97	-6.0 to -2.6	1,460*
Perfluorodecane sulfonic acid (PFDS)	0.002	--	--	0.7	7.7	3.5	--	--	19,200*
Perfluorobutyl phosphonic acid (PFBPA)	14260	--	--	0.18	2.2	--	--	--	--
Perfluorohexyl phosphonic acid (PFHxPA)	515	--	--	0.04	3.5	--	--	--	--
Perfluorooctyl phosphonic acid (PFOPA)	25	96*	210*	0.01 5.7e-03*	4.7 4.8*	--	--	--	658*
Perfluorodecyl phosphonic acid (PFDPA)	0.5	87*	210*	0.0002 1.5e-03*	6.0 3.9*	--	--	--	18,100*
Perfluorobutyl phosphinic acid (PFBPIA)	2.3e-06 to 8.2e-03*	-9.9 to 70.1*	208 to 245*	6.8e-07 to 1.2e-02*	2.4 to 11.9*	--	--	--	1,390*

Perfluorohexyl phosphinic acid (PFHxPiA)	3.6e-07 to 3.1e-04*	-25.1 to 106*	213 to 301*	8.5e-11 to 9.6e-03*	4.4 to 15.1*	--	--	--	231,000*
Perfluorooctyl phosphinic acid (PFOPiA)	1.7e-09 to 2.8e-04*	-9.9 to 138*	206 to 353*	9.0e-13 to 1.7e-02*	7.2 to 18.6*	--	--	--	134,000*
Perfluorodecyl phosphinic acid (PFDPiA)	2.9e-08*	10.4*	265*	1.9e-09*	7.6*	--	--	--	132,000*
Perfluorododecyl phosphinic acid (PFDODPiA)	5.4e-07*	50.7*	290*	2.2e-09*	7.5*	--	--	--	133,000*

Environmental fate

Reference: CAL. CODE REGS. tit. 22, § 69503.3(a)(1)(E).

Environmental fate describes a chemical's mobility in environmental media, transformation (physical, chemical, or biological), or accumulation in the environment or biota. A chemical's environmental fate in air, water, soil, and living organisms relates to its exposure potential hazard traits, as defined in the California Code of Regulations, Title 22, Chapter 54.

PFASs are widespread in indoor and outdoor environments, humans, and biota. They are found even in raindrops and snowflakes, in high-altitude atmospheric wind currents (Taniyasu et al. 2013), and the deep sea (Sanchez-Vidal et al. 2015; Zhao et al. 2012). The accumulation of PFASs in different environmental compartments and living organisms appears to depend on their partitioning behavior, which depends in part on the perfluoroalkyl chain length and the functional group of the parent compound (EFSA 2011; Wang et al. 2017b).

Numerous factors, including intrinsic chemical properties combined with physical and biogeochemical processes, collectively determine the occurrence and fate of PFASs (ITRC 2018). Longer-chain PFASs bioaccumulate up the food chain (Concawe 2016). Shorter-chain PFASs are particularly mobile in water and air, can undergo long-range transport, and are difficult to remove from contaminated water (Brendel et al. 2018; Li et al. 2020). PFAAs are the terminal environmental degradation products of most PFASs, and all are extremely persistent in the environment (Wang et al. 2017b). Shorter-chain PFAAs are distributed globally by oceanic currents and sea spray, or by the degradation of volatile precursors that undergo long-range atmospheric transport (Ateia et al. 2019b). Long-range atmospheric transport followed by wet or dry deposition has led to the ubiquitous presence of PFBA and other PFAAs in snow, water, soil, and sediment samples from a remote alpine nature reserve in China (Wang et al. 2019).

Some studies distinguish between PFAAs entering the environment from direct and indirect sources (Buck et al. 2011; Prevedouros et al. 2006). "Direct" sources refer to PFAA emissions from the life cycle of products containing PFAAs or their derivatives as ingredients, unreacted raw materials, or unintended by-products; "indirect" sources refer to PFAA formation from the degradation of precursors (OECD 2013). Other references refer to these as either "point" or "diffuse" sources (ITRC 2018). Monitoring data suggest that a large number of PFASs are PFAA precursors, meaning they can transform into PFAAs in the environment (Ahrens 2011; ITRC 2018).

Regardless of their sources, due to their unique physicochemical properties, PFASs released to the environment end up virtually everywhere, including air, dust, wastewater treatment plant (WWTP) effluent, biosolids, soil, inland and ocean waters, drinking water, and food (Lindstrom et al. 2011b; Wang et al. 2016). PFAAs and their precursors are found in the deep ocean and in underground

aquifers, in rainwater, snow, ice, glaciers, and pristine Arctic lakes, far from any point source (Kim and Kannan 2007; Kirchgeorg et al. 2016; Muir et al. 2019; Prevedouros et al. 2006; Zhao et al. 2012). Wildlife and humans are often sinks of environmental PFASs (Cariou et al. 2015; Chang et al. 2014; Fraser et al. 2013; Giesy and Kannan 2001; Kang et al. 2016; Liu et al. 2011a; Monroy et al. 2008; OECD 2013; Olsen et al. 2007b; Tao et al. 2008a, b).

Volatile PFAA precursors, including FTOHs, are distributed globally as gases or adsorbed to airborne particles (Dreyer et al. 2015; Liu et al. 2015a; Wang et al. 2014b), and can re-volatilize back into the atmosphere from water or snow (Xie et al. 2015). PFAAs can become airborne and deposit onto water surfaces (Kim and Kannan 2007). Global distribution also occurs due to movement of oceanic currents, with PFHxA and PFBS being among the most frequently detected compounds in seawater (Zhao et al. 2012).

Although a considerable number of studies document levels of some PFASs in the environment and various biological media, knowledge regarding the sorption/desorption patterns and mechanisms of transport, transformation, and fate of most PFASs is limited. Relatively few studies are available on PFASs in sediments, as environmental studies worldwide have focused more attention on the quantities of PFASs in biota and water, and less on sediments and their partitioning behavior and fate (Nakata et al. 2006; Zareitalabad et al. 2013). Data on persistence and half-life estimates in biological and environmental media are also limited, and are sometimes conflicting or inaccurate. Half-lives are estimated based on acute exposures rather than on the low-dose chronic exposures experienced by most people and biota. Major data and knowledge gaps remain regarding the sources, transformation, distribution, accumulation, and temporal and spatial trends of various PFASs in wildlife and humans.

The following sections summarize publicly available data on the behavior of PFASs in different environmental compartments and sinks, without following a particular pathway, since the transport and cycling mechanisms of all PFASs in the environment are not fully understood. The available environmental fate data are limited to PFAAs and a few of their precursors. However, since PFAAs are used in the manufacturing of fluoropolymers and PFPEs, the following discussion can be relevant to the life cycle of all PFASs.

This potential for long-range atmospheric transport followed by deposition onto land and water has resulted in PFAS-contaminated food chains and accumulation in plants and animals, even in remote Arctic and Antarctic regions.

PFASs in air

PFASs display a wide range of airborne behaviors. Some PFASs, such as FTOHs, are relatively volatile and detected in indoor air and dust (Ericson et al. 2012; Fraser et al. 2013; Karásková et al. 2016). Release of these PFASs into air can occur during chemical manufacturing, consumer product manufacturing, product use, around WWTPs, and under landfill conditions after disposal (Ahrens 2011; Ahrens et al. 2011b). The most volatile PFASs are PFAA precursors, which can oxidize to PFAAs in the atmosphere (Ellis et al. 2004; Wallington et al. 2006). PFAAs are semi-volatile and dissociate in water, as the perfluoroalkyl chain forms micelles on the water surface and partitions into the air (Eftaiha et al. 2012; ITRC 2018; Mason Chemical Company 2017).

Once PFASs enter the air, they can be carried long distances by air currents or adsorb to particulate matter (Barber et al. 2007; Dreyer et al. 2015; Liu et al. 2015a; Shoeib et al. 2010; Wang et al. 2014b), which may be directly inhaled or deposited in rain and snow (Casal et al. 2017; Sammut et al. 2017; Taniyasu et al. 2013). Neutral volatile PFAS precursors, such as FTOHs, are the dominant PFASs found in the gas phase (Ahrens et al. 2012), whereas ionic PFASs, such as PFOA and PFOS, characterized by low vapor pressure and high water solubility, tend to be the dominant species in airborne particulate matter (ITRC 2018). According to the limited publicly available data, PFASs have long atmospheric half-lives, for instance approximately 20 days for 8:2 FTOH (Ellis et al. 2003).

Due to their potential for long-range atmospheric transport followed by deposition onto land and water, PFASs have contaminated food chains and accumulated in plants and animals, even in remote Arctic and Antarctic regions (Houde et al. 2006; Butt et al. 2010; Casal et al. 2017; Muir et al. 2019; Roscales et al. 2019). Despite the U.S.-wide phase-out of longer-chain PFASs, atmospheric emissions of these compounds from offshore manufacturing may still impact the United States. Trans-Pacific atmospheric transport from China leading to adverse impacts on air quality in California has previously been documented for other pollutants (Lin et al. 2014b). Considering the large number of PFASs in commerce today, their continuous air emissions, and their relatively long atmospheric half-lives, atmospheric transport is of significant concern due to the potential for widespread PFAS environmental distribution and contamination impacting humans and other organisms.

PFASs in water

PFASs enter surface water and groundwater through a variety of routes, including industrial discharges from manufacturing sites, WWTP effluent, runoff and leaching from contaminated soil or landfills, deposition of contaminated particulate matter onto water bodies, and glacial meltwater. Their individual solubility and partitioning behavior, combined with the physical and chemical characteristics of receiving waters, determine whether specific PFASs are found dissolved or suspended in the water column, in micelles at the air/water interface (Eftaiha et al. 2012; Mason Chemical Company 2017), or

adsorbed to solid particles (Campo et al. 2016). Longer-chain PFAAs tend to partition into suspended particles and sediments, while shorter-chain PFAAs tend to remain in the dissolved phase (Chen et al. 2019).

Due to their low pKa values, most PFAAs are present in their anionic form in surface waters and tend to remain in solution (ITRC 2018); from there, PFAAs can contaminate groundwater through groundwater recharge (Liu et al. 2016) or be transported to the oceans where they become globally distributed by ocean currents (Benskin et al. 2012b). Stormwater runoff may contribute significantly to the PFAS load in surface waters (Wilkinson et al. 2017; Zushi and Masunaga 2009). Municipal and industrial WWTPs can release PFASs to the environment via point source effluent discharges, leakage or unintended releases, air emissions, and disposal of biosolids and other treatment byproducts (ITRC 2018). During the wastewater treatment process, volatile PFASs can be released into the air, while others selectively partition into biosolids and are partially removed from wastewater. WWTP effluent tends to have higher concentrations of PFAAs than the influent because of the transformation of precursors during the treatment process (Appleman et al. 2014; Arvaniti and Stasinakis 2015; Sinclair and Kannan 2006). PFASs also enter surface water and groundwater via landfill leachate (Ahrens et al. 2016a; Banzhaf et al. 2017).

PFAAs appear to not degrade under environmental conditions, even in activated sewage sludge, and can be more persistent than other legacy persistent organic pollutants such as dichlorodiphenyltrichloroethane (DDT), polychlorinated biphenyls (PCBs), and dieldrin (EWG 2003; European Commission 2017). Like other persistent organic pollutants, oceans are considered the ultimate long-term reservoir for PFASs (Prevedouros et al. 2006), including PFAAs and their precursors (González-Gaya et al. 2014). Marine organisms are chronically exposed via contaminated food chains (Ahrens et al. 2009a; Casal et al. 2017; Houde et al. 2011; Xiao 2017). Many physical and biogeochemical processes and factors collectively determine the oceanic occurrence and fate of PFASs, including distance to coastal urban industrial regions, oceanic subtropical gyres, currents, and biogeochemical processes (González-Gaya et al. 2014). PFASs are also transported downwards from the ocean surface mainly by sinking zooplankton fecal pellets and decaying phytoplankton, but these removal mechanisms are slow (González-Gaya et al. 2019). Oceanic currents can transport PFASs over long distances, resulting in PFAS contamination of water in distant pristine regions of the world, far from industrial activities (Prevedouros et al. 2006; Ahrens et al. 2011a).

Snow and ice in the northern latitudes also accumulate PFASs, including the shorter chains (Del Vento et al. 2012). Climate change resulting in snow, ice, and permafrost melt can remobilize PFASs and increase their concentrations in coastal areas (Zhao et al. 2012).

The ubiquitous presence of PFASs in surface water, groundwater, oceans, sediments, and air has also resulted in drinking water contamination, especially with the more soluble and mobile shorter-chain

PFAAs (Boone et al. 2019; Gellrich et al. 2013). Shorter-chain PFASs are generally more mobile in groundwater and will leach faster from soil to groundwater (NGWA 2017). Contamination of drinking water with PFASs and irrigation of food crops with contaminated water is generally considered a major source of human exposure, as discussed in subsequent sections of this document.

PFASs in sediments

Sediments are an important sink and reservoir of PFASs in the aquatic environment (Chen et al. 2016; Pan et al. 2015). Freshwater and marine sediments can become contaminated when PFASs dissolved in water adsorb onto particulate matter and are deposited. PFASs exhibit a range of sediment-water partitioning coefficients that influence their environmental fate. Long-term production and use patterns for PFASs are documented in sediment cores (Codling et al. 2014). Longer-chain PFASs such as PFOS and PFOA are typically the predominant PFASs found in surface sediments (Rankin et al. 2016). Spatial distribution of PFASs in sediments and the occurrence of “hot spots” in sampling areas can provide evidence of specific manufacturing activities and point source discharges (Munoz et al. 2015; White et al. 2015). Monitoring studies (see Chapter 5) suggest that urbanization and industrial activities contribute to PFAS accumulation in sediments, particularly in coastal regions and estuaries near urban centers. PFAS-contaminated coastal waters can be carried to deep sea regions through deep shelf water cascading events (Sanchez-Vidal et al. 2015). Persistent organic pollutants similar to PFASs have been found in ocean sediments deeper than 10,000 meters, where they can bioaccumulate in marine organisms and biomagnify through food chains (Jamieson et al. 2017).

The ubiquitous presence of PFASs in surface water, groundwater, oceans, sediments, and air has also resulted in drinking water contamination, especially with the more soluble and mobile shorter-chain PFAAs.

PFASs in soil

PFASs can contaminate the soil via atmospheric transport followed by deposition, or directly through application of biosolids (Sepulvado et al. 2011), soil conditioners, or other treatments. From soil, they can migrate into surface water or groundwater, including drinking water sources (Lilienthal et al. 2017). The sorption of PFASs to soil is influenced by several factors, including soil organic matter content, clay content, pH, and the presence of cations (Higgins and Luthy 2006; Li et al. 2018). The fate and transport of a specific PFAS in soil depends on its physicochemical properties. Longer-chain PFASs tend to partition preferentially into soil (Dalahmeh et al. 2018), while shorter-chain PFASs partition into soil water (Ahrens et al. 2009b; Guo et al. 2015). Longer-chain PFAAs added to an agricultural soil were found to sorb strongly to the soil, whereas the shorter-chain PFAAs were readily transported in water from the soil surface to the root zone and into the groundwater (McLachlan et al. 2019; Muir et al.

2019). Shorter-chain PFCAs have also been observed to migrate out of soil following biosolid application, with disappearance half-lives ranging from 88 to 866 days (Venkatesan and Halden 2014). According to the study authors, the fate of these PFCAs after loss from soils could include leaching, plant uptake, and volatilization (Zhang et al. 2019a).

Publicly available data on the half-lives of PFASs in soils are scarce and vary widely across models and experiments. For example, one industry study reports a half-life of 1,200 to 1,700 years for the biodegradation of a side-chain fluoroacrylate polymer to PFOA in aerobic soil (Russell et al. 2008). A team of U.S. EPA scientists (Washington et al. 2009) pointed out several issues with that study and arrived at an estimated half-life for fluoroacrylate polymers of 10-17 years based on experimental results and modeling. Subsequently, they evaluated the degradability of two commercial side-chain fluoroacrylate polymers in saturated soil and found half-lives of 33 to 112 years (Washington et al. 2015). Russell et al. (2010) estimated the half-life in soil of a side-chain fluorourethane polymer to be 28 to 241 years. More recently, U.S. EPA scientists calculated an environmental half-life for these polymers between nine and 60 years and concluded that the side-chain fluorinated polymers “now in production might constitute considerable sources to the environment of the new generation of PFASs” (Washington et al. 2019). Dasu and Lee (2016) studied the inherent biotransformation potential of two urethane monomers and found negligible degradation in agricultural soil, but significant degradation in forest soil, with an estimated half-life ranging between three and 22.2 months. Royer et al. (2015) studied the biotransformation of 8:2 FTAC and 8:2 FTMAC to 8:2 FTOH in soils, finding half-lives of five days or less and 15 days, respectively.

PFASs in biota

Plants and animals can bioaccumulate PFASs, with large interspecies differences, particularly between aquatic and terrestrial organisms (Concawe 2016). PFAS uptake has been documented in fruits and vegetables grown in backyard and garden soils irrigated with contaminated water (Scher et al. 2018), agricultural soils spiked with PFASs (McLachlan et al. 2019), potted soils spiked with PFASs (Stahl et al. 2009; Zhang et al. 2019a), and biosolids-amended soil (Blaine et al. 2013, 2014a; Yoo et al. 2011). Crops grown in agricultural soils without significant PFAS contamination were found to contain small levels of mainly shorter-chain PFASs from rainwater and atmospheric deposition (Eun et al. 2020). Longer-chain PFAAs, especially PFSAs, tend to partition in the roots, whereas the more soluble, shorter-chain PFAAs, especially PFCAs, partition in other parts of the plant (Blaine et al. 2013, 2014a; Gobelius et al. 2017; Lechner and Knapp 2011; Scher et al. 2018; Stahl et al. 2009; Yoo et al. 2011). Studies have also reported PFAA uptake in tree leaves, possibly from a combination of soil uptake and atmospheric deposition, in an urban environment (Zhang et al. 2015) and near industrial emission sources (Shan et al. 2014).

The extent of bioaccumulation is traditionally measured using several different indices. The bioconcentration factor (BCF) is a measure of the amount of a contaminant found in an aquatic organism compared to the amount found in water. The biomagnification factor (BMF) compares the amount of the contaminant in the organism to that in its diet. The bioaccumulation factor (BAF) combines uptake from all sources, including water, diet, and environmental exposures (Arnot and Gobas 2006). Chemicals with BAF or BCF values greater than 1,000 or BMF greater than 1 meet the bioaccumulation hazard trait criteria outlined in the CAL. CODE REGS. tit. 22, Division 4.5, Chapter 54, Article 5. However, these traditional measures of bioaccumulation have limited applicability when gauging the potential for PFAS exposures to cause adverse health effects (Cousins et al. 2016). Although the elimination kinetics and toxicity of some single PFASs have been experimentally determined, the ubiquity and extreme persistence of PFASs in the environment lead to continuous internal exposures to mixtures of PFASs, which have not been adequately assessed (Wang et al. 2017b).

Table 2(a-c) summarizes BCFs, BMFs, and BAFs for some PFASs with publicly available data. The BCFs of some longer-chain PFAAs approach 1,000, and a few – PFOS and perfluorohexadecanoic acid (PFHxDA) – exceed it. Animals tend to have greater concentrations of PFASs in their bodies compared to their diets. BAFs in aquatic organisms depend on the type and concentration of specific PFASs in water and are generally proportional to the length of the carbon chain or molecular weight, with PFASs showing higher bioaccumulation than PFCAs of the same chain length – although there is interspecies variation (Hong et al. 2015).

Once ingested via food or drinking water, some PFASs remain in the human body for years. Limited studies suggest that serum elimination half-life can range from a few days for shorter-chain PFASs (Chang et al. 2008; Nilsson et al. 2010; Olsen et al. 2007a) to several years for longer-chain PFASs (Bartell et al. 2010; Olsen et al. 2007a; Seals et al. 2011). It can take up to 56 years for total elimination of a chlorinated PFESA (Shi et al. 2016). Reported half-lives for PFAAs and FTOHs in other organisms such as rats, mice, and monkeys are shorter than in humans, typically hours, days, or months (Butenhoff et al. 2004; Chang et al. 2008, 2012; Gannon et al. 2011; Ohmori et al. 2003; Sundström et al. 2012; Tatum-Gibbs et al. 2011).

Table 2a. Examples of bioconcentration factors (BCFs)

Chemical	Organism	BCF	Reference
PFOS	Bluegill	1,866 – 4,312	Drottar (2001)
PFOS	Rainbow trout	1,100 – 5,400	Drottar (2001)
PFOS	Rainbow trout	2,900 (liver), 3,100 (blood)	Martin et al. (2003)
PFOA	Water breathing animals	1.8 – 8.0	ECHA (2014)
PFOA	Rainbow trout	12 (liver), 25 (blood)	Martin et al. (2003)
PFHxDA	Carp	4,700 – 4,800	U.S. EPA (2009)
Perfluorooctadecanoic acid (PFODA)	Carp	320 – 430	U.S. EPA (2009)

Table 2b. Examples of bioaccumulation factors (BAFs)

Chemical	Organism	BAF	Reference
PFOS	Zooplankton/water	240	Houde et al. (2008)
PFOS	Mysis/water	1,200	Houde et al. (2008)
PFOS	Sculpin/water	95,000	Houde et al. (2008)
PFOS	Lake trout/water	16,000	Houde et al. (2008)
PFOA	Water breathing animals	0.9 – 266	ECHA (2014)
PFOA	Copepods/water	200 – 500	Munoz et al. (2019)
PFOA	Copepods/suspended solids	251	Munoz et al. (2019)
PFOA	Mysids/water	316	Munoz et al. (2019)
PFOA	Mysids/suspended solids	3.16 – 79.4	Munoz et al. (2019)

Chemical	Organism	BAF	Reference
PFOA	Shrimp/water	125 – 631	Munoz et al. (2019)
PFOA	Shrimp/suspended solids	2.5 – 31.6	Munoz et al. (2019)
PFHpA	Copepods/water	200 – 500	Munoz et al. (2019)
PFHpA	Copepods/suspended solids	251	Munoz et al. (2019)

Table 2c. Examples of biomagnification factors (BMFs)

Chemical	Organism	BMF	Reference
PFOS	Arctic cod/zooplankton (Western Canadian Arctic)	8.7	Powley et al. (2008)
PFOS	Caribou/lichen (Canada)	2.0 – 9.1	Müller et al. (2011)
PFOS	Wolf/caribou (Canada)	0.8 – 4.5	Müller et al. (2011)
PFOS	Dolphin/seatrout (2 U.S. locations)	0.9	Houde et al. (2006)
PFOS	Seatrout/pinfish (2 U.S. locations)	4.6	Houde et al. (2006)
PFOS	Walrus/clam (Eastern Arctic Food Web)	4.6	Tomy et al. (2004)
PFOS	Narwhal/Arctic cod (Eastern Arctic Food Web)	7.2	Tomy et al. (2004)
PFOS	Beluga/Arctic cod (Eastern Arctic Food Web)	8.4	Tomy et al. (2004)
PFOS	Beluga/redfish (Eastern Arctic Food Web)	4	Tomy et al. (2004)
PFOS	Polar bear/seal (Canadian Arctic)	177	Martin et al. (2004)
PFOA	Water-breathing animals	0.02 – 7.2	ECHA (2014)
PFOA	Caribou/lichen (Canada)	0.9 – 11	Müller et al. (2011)
PFOA	Wolf/caribou (Canada)	0.9 – 3.8	Müller et al. (2011)
PFOA	Walrus/clam (Eastern Arctic Food Web)	1.8	Tomy et al. (2004)
PFOA	Narwhal/Arctic cod (Eastern Arctic Food Web)	1.6	Tomy et al. (2004)
PFOA	Beluga/Arctic cod (Eastern Arctic Food Web)	2.7	Tomy et al. (2004)

Chemical	Organism	BMF	Reference
PFOA	Beluga/redfish (Eastern Arctic Food Web)	0.8	Tomy et al. (2004)
PFOA	Beluga whale/Pacific herring (Western Canadian Arctic Food Web)	1.3	Tomy et al. (2004)
PFOA	Arctic cod/marine Arctic copepod (Western Canadian Arctic Food Web)	2.2	Tomy et al. (2004)
PFOA	Dolphin/seatrout (2 U.S. locations)	1.8	Houde et al. (2006)
PFOA	Seatrout/pinfish (2 U.S. locations)	7.2	Houde et al. (2006)
PFOA	Polar bears/ringed seal (2 U.S. locations)	45 – 125	Butt et al. (2008)
PFOA	Polar bear/seal (Canadian Arctic)	8.6	Martin et al. (2004)

Degradation, reaction, or metabolic products of concern

Reference: CAL. CODE REGS. tit. 22, § 69503.3(a)(1)(G).

A Candidate Chemical may degrade, form reaction products, or metabolize into other chemicals that have one or more hazard traits. These metabolites, degradation products, and reaction products (which may or may not be Candidate Chemicals) may cause different adverse impacts from those of the parent chemical. In some cases, a Candidate Chemical's degradation or reaction products or metabolites may have the same hazard trait, and may be more potent or more environmentally persistent, or both, than the parent chemical. In such cases, adverse impacts may be more severe, or may continue long after, the Candidate Chemical's release to the environment.

Of the 4,730 individual PFASs or commercial PFAS mixtures identified by OECD on the global market, 4,186 likely degrade to PFAAs in the environment or biota (OECD 2018). PFASs that degrade into PFAAs are termed PFAA precursors and provide a significant indirect source of PFAAs to the environment (see examples of PFAA precursors in Chapter 4). Degradation to PFAAs can occur, for instance, in the atmosphere, consumer products, landfills, or WWTPs, mainly via hydroxylation or photo-oxidation (Nielsen 2014; Prevedouros et al. 2006). The degradation half-life of PFAA precursors can range from days to centuries (Li et al. 2017; Nielsen 2014; Washington et al. 2015).

PFAA precursors can persist in the environment and in biological compartments for various lengths of time, but eventually degrade through biotic and abiotic transformation to PFAAs, via intermediate metabolites suspected of high acute toxicity (Butt et al. 2014; D'Eon and Mabury 2007; Li and Guo 2016; Rand and Mabury 2012a, b, 2013, 2014, 2017). Fluorotelomer-based compounds degrade

(Phillips et al. 2007; Young et al. 2007) or undergo biotransformation into PFCAs (Chen et al. 2017b; Dasu and Lee 2016; Kim et al. 2012a, 2014b; Lee et al. 2010; Liu et al. 2010a, b; Rand and Mabury 2014; Royer et al. 2015; Ruan et al. 2014; Russell et al. 2015; Zhao and Zhu 2017). The intermediates in this process include saturated and unsaturated fluorotelomer aldehydes (FTALs and FTUALs, respectively), and saturated and unsaturated fluorotelomer carboxylic acids (FTCAs and FTUCAs, respectively) (Frömel et al. 2016; Nilsson et al. 2013; Rand et al. 2014). FTUCAs were detected in all 81 indoor air samples collected in 2015 during a study in China, suggesting the transformation of PFAA precursors such as FTOHs in the indoor environment (Yao et al. 2018). Pharmacokinetic data from rat and human studies on 6:2 FTOH revealed that, of its main metabolites, the 5:3 fluorotelomer carboxylic acid has the highest internal exposure and slowest clearance, which decreases with increasing exposure (Kabadi et al. 2018). The study also found that 5:3 FTCA may reach steady state following repeated exposure to 6:2 FTOH, therefore resulting in biopersistence, with potential for systemic toxicity (Kabadi et al. 2018).

Laboratory studies of fluorotelomer carboxylic acid and aldehyde intermediates indicate they are more acutely toxic than PFCAs, including PFOA. One study found that FTCAs and FTUCAs are one to five orders of magnitude more toxic to freshwater organisms than PFCAs (Phillips et al. 2007). In a human liver cell toxicity assay, FTUALs were found to be up to 200 times more toxic than FTUCAs, FTCAs, and PFCAs (Rand et al. 2014).

Side-chain fluorinated polymers, which are fluorotelomer-based, can also degrade to FTOHs or other fluorotelomer compounds, with PFCAs as terminal degradation products (Washington et al. 2015, 2019). Because of their potential to be mobilized, interact with their surrounding environment, and degrade (KEMI 2015; Russell et al. 2010), side-chain fluorinated polymers are sometimes referred to as “functionalized oligomers” to distinguish them from true polymers.

Intentional or accidental combustion of PFASs also forms hazardous chemicals. A literature survey regarding the combustion of various fluorinated polymers reported emissions of C3-C14 PFCAs, ozone depleting substances such as chlorofluorocarbons (CFCs), and greenhouse gases such as fluorocarbons when fluoropolymers are combusted at temperatures representative of municipal incinerators (Huber et al. 2009). In a review of how fluoropolymers differ from nonpolymeric PFASs, the authors acknowledged that end-of-life considerations should be further investigated because hazardous substances may arise from the combustion of fluoropolymers (Henry et al. 2018). For instance, during incineration at temperatures above 450°C, PTFE forms additional hazardous substances including hydrofluoric acid (HF) (Henry et al. 2018), which is on DTSC’s Candidate Chemicals list. An industry-sponsored experimental study in a rotary kiln test facility simulating municipal incinerators found that PTFE polymer pellets begin to decompose at around 500°C, and by around 650°C they completely convert to HF gas and F-containing ash, with no significant PFAA emissions (Aleksandrov et al. 2019). Other authors report that at lower temperatures, as could occur during accidental landfill fires,

fluoropolymers such as PTFE can break down into PFCAs, including PFOA (Feng et al. 2015; Schlummer et al. 2015).

Thus, while PFAAs constitute a small subset (approximately 1 percent) of PFASs, they are terminal degradation products, manufacturing aids/feedstocks, or impurities of other PFAS class members. This makes their hazard traits relevant to the entire class (see Figure 2).




Figure 2: Use of all PFAS subclasses can lead to exposure to PFAAs at some point in the chemicals' life cycle; they either degrade to form PFAAs, release PFAAs if incinerated, or require PFAAs for their manufacture, often resulting in PFAA impurities in the final product. Specifically:

- PFPEs and fluoropolymers are made using PFAAs
- PFAAs can be released from PFPEs and fluoropolymers during combustion
- PFAA precursors such as fluorotelomer-based substances, including side-chain fluorinated polymers, degrade to PFAAs

Hazard Traits and Environmental or Toxicological Endpoints

Reference: CAL. CODE REGS. tit. 22, § 69503.3(a)(1)(A).

The hazard traits and environmental or toxicological endpoints summarized in this section are defined in the SCP regulations sections 69501.1(a)(36) and (33), respectively, both of which refer to the Office of Environmental Health Hazard Assessment's (OEHHA) Green Chemistry Hazard Trait regulations (California Code of Regulations, Title 22, Chapter 54).¹¹ These include exposure potential, toxicological, and environmental hazard traits.

PFASs display multiple hazard traits according to OEHHA's Green Chemistry Hazard Traits regulations (CAL. CODE REGS. tit. 22, § 69401 et seq.). These include toxicological hazard traits, (Articles 2 and 3), environmental hazard traits (Article 4), and exposure potential hazard traits (Article 5).

¹¹ http://oehha.ca.gov/multimedia/green/pdf/GC_Regtext011912.pdf

Exposure potential hazard traits

Environmental persistence

Environmental persistence is one of the hazard traits identified in the Green Chemistry Hazard Traits Regulations. According to OEHHA's Statement of Reasons for these regulations (OEHHA 2012):

"Persistence of a chemical in the environment promotes sustained exposure and contributes to accumulation in the environment. Because persistence is an inherent property of a chemical in the environment that results in increased exposure to the chemical and consequently potential for health risks, it can appropriately be identified as a hazard trait. Legacy chemicals such as DDT and PCBs remain public health concerns decades after their production was banned because of their ability to persist in the environment."

PFAAs are extremely persistent in the environment, with the exception of PFPiAs, which degrade into PFPAs and potentially PFCAs (Scheringer et al. 2014; Wang et al. 2016). All other PFAAs degrade significantly only under environmentally irrelevant conditions. PFOS, its salts, and perfluorooctane sulfonyl fluoride are designated as persistent organic pollutants (POPs) under the Stockholm Convention (SSCCH 2017). PFHxS and C11 through C14 PFCAs are listed as very persistent and very bioaccumulative (vPvB) on the European Chemicals Agency (ECHA)'s Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) Candidate List of Substances of Very High Concern (SVHC); PFOA, ammonium perfluorooctanoate (APFO), PFNA, and PFDA are listed as persistent, bioaccumulative, and toxic (PBT) (ECHA 2017). Limited data on some PFECAs and PFESAs suggest resistance to biodegradation (Beekman et al. 2016). PFPEs and fluoropolymers are similarly persistent, whereas PFAA precursors degrade into extremely persistent PFAAs.

Mobility in environmental media

As discussed in Chapter 4, PFAAs are capable of long-range transport. Depending on their physicochemical properties, such as water solubility, K_{oc} , and volatility, PFASs can be transported in the dissolved phase (e.g., through soil water, by rivers and oceanic currents), in the particulate phase (e.g., via suspended sediments or airborne particles), or in the atmosphere. Shorter-chain PFAAs have higher aquatic persistence and mobility than the longer-chains, and thus are more frequently detected in the aquatic environment (Li et al. 2020).

Bioaccumulation

PFAAs tend to bioaccumulate, with elimination half-lives increasing with perfluoroalkyl chain length. In humans, known serum elimination half-lives range from a few days to several years (Bartell et al. 2010; Chang et al. 2008; Olsen et al. 2007a; Nilsson et al. 2010; Seals et al. 2011). PFPiAs may be an exception, since they can biotransform to PFPAs (Joudan et al. 2017; Yeung and Mabury 2016). ECHA's

REACH Candidate List of SVHCs designates PFHxS and C11 through C14 PFCAs as very persistent and very bioaccumulative (vPvB), and PFOA, APFO, PFNA, and PFDA as bioaccumulative (PBT) (ECHA 2017). PFAAs differ in their uptake and accumulation in human tissues (Burkemper et al. 2017; Pérez et al. 2013). PFAAs cross the brain blood barrier and are present in animal brain tissue (Greaves et al. 2013). Biomagnification, which is the increase in contaminant concentration higher in the food chain, has been reported for PFAAs in freshwater and marine organisms (Houde et al. 2006). Less is known about the potential for bioaccumulation or biomagnification of PFECAs and PFESAs (Beekman et al. 2016). To our knowledge, the bioaccumulation potential of other PFASs besides PFAAs has not been evaluated.

Lactational or transplacental transfer

Transplacental and lactational transfer can be significant routes of exposure for newborns (Cariou et al. 2015; Papadopoulou et al. 2016; Winkens et al. 2017b). PFASs, including PFAAs, nonpolymeric PFAA precursors, and chlorinated polyfluorinated ether sulfonates, undergo transplacental transfer in humans (Midasch et al. 2007; Zhang et al. 2013; Yang et al. 2016; Chen et al. 2017a; Zhao et al. 2017; Eryasa et al. 2019). The placenta-to-maternal serum ratios of PFOS, PFOA, and PFNA increased during gestation, more so in pregnancies with male fetuses compared to female ones, suggesting bioaccumulation in the placenta and increasing exposure with fetal age (Mamsen et al. 2019). Lactational transfer has also been documented for PFAAs in humans (Kärman et al. 2007; Llorca et al. 2010; Mogensen et al. 2015; Mondal et al. 2014, p. 2; Tao et al. 2008b), but to our knowledge has not been studied for PFAA precursors or other PFASs.

Global warming potential

Some PFPEs, which are increasingly used as replacements for PFAAs or their precursors, may have high global warming potential.¹² The global warming potential of perfluoropolymethylisopropyl ether, a type of PFPE, ranges from 7,620 over 20 years to 12,400 over 500 years, relative to carbon dioxide (CO₂) (IPCC 2007). Some nonpolymeric fluorinated ethers, which can be used in the production of PFPEs and released to the environment, also have high global warming potential (ranging, relative to CO₂, from 207 to 13,800 over 20 years, from 59 to 14,900 over 100 years, and from 18 to 8,490 over 500 years) (IPCC 2007). Additionally, fluoroform (CHF₃), with a global warming potential of 11,700 to 14,800 relative to CO₂ over 100 years (IPCC 2007), is a degradation product of fluorinated polymers from incomplete combustion (Huber et al. 2009).

¹² Global warming potential (GWP) is defined by the California Air Resources Board (CARB) at <https://ww3.arb.ca.gov/cc/inventory/background/gwp.htm>. The GWP value represents the climate forcing of a kilogram of emissions of a substance relative to the same mass of carbon dioxide (CO₂), which has a GWP value of 1. In California, GWPs are calculated over a 100-year time frame.

Other PFASs ubiquitous in the ocean and transferred onto sea spray aerosols can significantly increase the concentration of cloud condensation nuclei, thus increasing cloud reflectance. This increases the efficiency of solar radiation absorption in the atmosphere, contributing to global warming (MacLeod et al. 2014).

Toxicological hazard traits

Toxicological data are limited to a few PFASs, mostly PFAAs. The toxicities of fluorinated polymers, their intermediate degradation products, and processing aids are poorly understood. Appendix 2 lists toxicological hazard traits for the PFASs with publicly available data (mostly PFAAs and some precursors). Most toxicological studies evaluate exposure to single PFASs for 90 days or less, even though humans and animals are chronically exposed to PFAS mixtures over a lifetime.

Potential adverse human health effects and risk factors from longer-chain PFAS exposure include:

- Increased serum cholesterol (Skuladottir et al. 2015; Winqvist and Steenland 2014)
- Thyroid disease (Winqvist and Steenland 2014);
- Immune dysregulation (Grandjean and Clapp 2014; Grandjean et al. 2016, 2017)
- Pregnancy-induced hypertension (C8 Science Panel 2011); and
- Kidney and testicular cancers (C8 Science Panel 2012).

Epidemiological studies suggest a link to adverse health effects but often entail uncertainties in exposure or confounding factors. A systematic review of 64 epidemiological studies that assessed children's exposure to PFASs and associations with specific health outcomes found positive correlations between PFAS exposure and suppressed immune system response, dyslipidemia, impaired kidney function, and delayed first occurrence of menstruation (Rappazzo et al. 2017). PFOA also appears to be associated with low birth weight in humans (Malits et al. 2017).

A systematic review of 64 epidemiological studies that assessed children's exposure to PFASs and associations with specific health outcomes found positive correlations between PFAS exposure and suppressed immune system response, dyslipidemia, impaired kidney function, and delayed first occurrence of menstruation.

The health effects listed above were observed in studies focused on longer-chain PFASs. Possibly due to their shorter observed biological half-lives, shorter-chain PFASs were expected to have lower toxicity compared to the longer chains (DeWitt 2015; Stahl et al. 2011; Wolf et al. 2008). Notable exceptions are PFHxS (C6) (considered a longer-chain due to its high bioaccumulation potential) and PFBS (C4) in some toxicokinetic studies and receptor binding assays (Wolf et al. 2008; Danish Environmental Protection Agency 2015a). PFBS was reported to be developmentally toxic in zebrafish (Hagenaars et

al. 2011). Delayed pupil response (Butenhoff et al. 2012) and retinal degeneration¹³ in rodents (3M 2006) indicate a potential for ocular toxicity in humans. Rodent studies have also identified reproductive and developmental (fetal resorption, delayed eye opening) (Das et al. 2008), hematotoxic (reduced red blood cell count, hemoglobin, and hematocrit) (Lieder et al. 2009), and respiratory (nasal degeneration) (Loveless et al. 2009) hazards. Neurodevelopmental toxicity has been demonstrated *in vitro* by dose-dependent suppression of neuronal differentiation (Slotkin et al. 2008).

Scientific review of the limited data on the PFECA GenX (a PFOA replacement in some applications) and related perfluorinated ether substances from Chemours (Beekman et al. 2016) indicates potential associations with adverse health effects in laboratory animals, including cancer, body weight gain, changes to the immune system and cholesterol levels, increased weights of kidneys and livers, and liver cell changes (Rae et al. 2015).

When differences in rodent toxicokinetics are taken into consideration, PFECAs and shorter-chain PFAAs may have similar or higher toxic potency than the longer-chain PFAAs they are replacing. Using a toxicokinetic model and existing toxicity data sets, a recent study found that PFBA, PFHxA, and PFOA have the same potency to induce increased liver weight, whereas GenX is more potent (Gomis et al. 2018). The authors concluded that previous findings of lower toxicity of fluorinated alternatives in rats were primarily due to the faster elimination rates and lower distribution to the liver compared to PFOA and other longer-chain PFAAs.

As shown in Appendix 2, the evidence of adverse health effects from PFAS exposure in humans is sometimes conflicting (DeWitt 2015)(DeWitt 2015)(DeWitt 2015). Studies that examine associations between maternal serum, blood, or breast milk PFAS concentrations and child health outcomes sometimes show mixed results (Rappazzo et al. 2017). For example, one study found an association between increased maternal serum PFOS concentrations and increased hyperactivity in children (Høyer et al. 2015), while another reported a null association between PFOS in cord blood and the neurobehavioral symptoms of attention deficit/hyperactivity disorder (Lien et al. 2013). Studies have also suggested detrimental, beneficial, or no effects on metabolism. For instance, PFAS mechanisms of action can both exacerbate insulin resistance and improve insulin sensitivity (Fleisch et al. 2017).

Less is known about the toxicity of most PFAA precursors. Laboratory studies of fluorotelomer carboxylic intermediates indicate that they are more acutely toxic than PFCAs in nonmammalian systems (Phillips et al. 2007; Rand et al. 2014). Industry studies report increased mortality for 6:2 FTOH (O'Connor et al. 2014), and liver and kidney toxicity for 8:2 FTOH (Ladics et al. 2008), at the highest doses only. Other FTOH studies have found a potential for endocrine disruption, interfering with

¹³ Unpublished data mentioned in the cover letter accompanying a 3M submission to the TSCA 8(e) docket.

thyroid and sex steroid hormone systems (Ishibashi et al. 2008; Rosenmai et al. 2016; Weiss et al. 2009).

The mode of action of PFAAs has not been fully characterized. Activation of the nuclear peroxisome proliferation-activated receptor alpha (PPAR-alpha) has been associated with some of the hepatic effects of PFOA and PFOS, although other biological interactions associated with PFAAs' neurodevelopmental and reproductive toxicity have not yet been identified (Guyton et al. 2009; Rappazzo et al. 2017). PFASs have been shown to bind competitively to the human thyroid hormone transport protein (Weiss et al. 2009), and thus may affect thyroid hormone levels and early life brain development (Winkens et al. 2017b).

Environmental hazard traits

Appendix 2 lists several studies documenting the following environmental hazard traits:

- Phytotoxicity;
- Wildlife developmental impairment;
- Wildlife reproductive impairment; and
- Wildlife survival impairment

PFAS toxicity has been reported across a broad range of aquatic organisms (microorganisms, algae, plants, invertebrates, amphibians, fish, marine mammals) and adverse impacts (e.g., impaired growth, mortality, developmental effects, and reproductive effects). However, the environmental effects on terrestrial wildlife are not as well studied. See Appendix 2 for details.

A 2011 study testing the structure-activity relationship of PFOA, PFOS, PFBS, and PFBA in fish development found that all four PFASs were teratogens, leading to malformations of the tail and an uninflated swim bladder, causing fish to swim abnormally; exposure to PFBS and PFOS also resulted in fish head malformations (Hagenaars et al. 2011). Swimming and buoyancy are necessary for fish to compete for food and escape predators. Developmental toxicity was also observed in zebrafish embryos following exposure to 6:2 FTCA (Shi et al. 2017). Studies of zebrafish embryos found higher potential for adverse developmental effects for PFASs with sulfonic groups, such as PFBS and PFOS (Ulhaq et al. 2013), and potential endocrine-disrupting effects for 6:2 FTOH (Liu et al. 2009).

Studies show increased acute toxicity in microalgae and zebrafish embryos with increasing chain length (Latala et al. 2009; Ulhaq et al. 2013). A review of fish toxicity data concluded that shorter-chain PFASs have moderate to low acute toxicity at exposures of less than seven days; data on longer exposures were unavailable (Danish Environmental Protection Agency 2015a). A 2015 study found that levels of PFCAs and PFSA, including PFHxA and PFBS, found in various brain regions affected neurochemical markers in Greenland polar bears (Eggers Pedersen et al. 2015).

Structural or mechanistic similarity to chemicals with known adverse impacts

Reference: CAL. CODE REGS. tit. 22, § 69503.3(a)(3).

Some chemicals may lack sufficient data to definitively establish presence or absence of harm. In such cases, DTSC may also consider data from other chemicals closely related structurally to the Candidate Chemical to identify potential public health and environmental impacts.

While physicochemical properties and hazard traits for many PFASs are either uncharacterized or unavailable in the public domain, the data available for structurally similar PFASs raise sufficient concerns for DTSC to consider PFASs as a class. In particular, longer- and shorter-chain PFAAs share three key structural and mechanistic properties: their structural similarity to fatty acids (DeWitt et al. 2015), their potential to activate PPAR-alpha (Rosenmai et al. 2018; Wolf et al. 2008, 2014), and their persistence in the environment (Krafft and Riess 2015a). The carbon-fluorine bonds characteristic of the structure of all PFASs endow their extreme environmental persistence.

5. POTENTIAL FOR EXPOSURES TO THE CANDIDATE CHEMICAL IN THE PRIORITY PRODUCT

Reference: CAL. CODE REGS. tit. 22, § 69503.3(b).

The SCP regulations direct the Department to evaluate the potential for public or aquatic, avian, or terrestrial animal or plant organism exposure to the Candidate Chemical(s) in the product by considering one or more factors for which information is reasonably available.

Market presence and trends

Reference: CAL. CODE REGS. tit. 22, § 69503.3(b)(1)(A-C) and § 69503.3(b)(3).

Product market presence information may be used as a surrogate to assess potential exposures to the Candidate Chemical in the product. This information may include statewide sales by volume, the number of units sold or amount of sales generated, or information on the targeted customer base.

Tests in the 1940s found that fluorochemicals performed best in imparting resistance to the spreading of liquids on solid surfaces. This led 3M to develop the Scotchgard brand of oil and water repellents (Audenaert et al. 1999). The first commercial Scotchgard rain and stain repellent, called FC-208, was released in 1960. Subsequent formulations with increased ability to withstand laundering fall under three major categories: esters, polyurethanes, and acrylate copolymers (Audenaert et al. 1999). But scientific research highlighting the dangers posed by treatments containing PFASs challenged the industry. As a result, 3M announced in 2000 that it was discontinuing the manufacture and sale of Scotchgard, which at that time was based on C8 chemistry; however, the company reintroduced a reformulated version 16 months later (Singh et al. 2005).

According to data on two broad North American Industry Classification System (NAICS) codes,¹⁴ the number of U.S. manufacturers and sales revenues for these industry sectors have steadily declined from 2007 to 2017, by 17.2 percent and 32.5 percent, respectively (U.S. Census Bureau 2007, 2012, 2017a). Foreign imports of treatment products, however, may meet any unsatisfied demand for these products. Moreover, market data gaps remain substantial. Accessible government, nongovernmental, academic, and corporate data sources reveal only scarce sales information for specific treatment products.

A September 2019 search of the Mintel Global New Products Database identified approximately 50 new treatment product releases over the previous five-year span (Mintel 2019). These products included stain removers and protectors and leather protectors. Detecting the presence of PFASs in

¹⁴ These two NAICS codes – 325612 and 325613 – comprise ‘polish and other sanitation good’ and ‘surface active agent’ manufacturing. Some nontreatment manufacturers fall within these codes.

these types of products, however, remains difficult. Many product labels and Safety Data Sheets fail to provide complete lists of ingredients. In some cases, companies provide no ingredient information whatsoever. Governmental and academic research offers some level of insight into the presence of PFASs in treatments. For example, a 2008 Danish report estimated that at most 10 percent of the treatments for textiles and leather contain PFASs (Jensen et al. 2008).

Treatment products containing PFASs often exist in homes and places of work throughout California. Even a person who chooses not to purchase a PFAS-containing upholstery protector may own several spray bottles of PFAS-containing stain remover. A wide range of occupations rely on treatment products that contain PFASs; these include shoe and leather workers and repairers, vehicle cleaners, and building cleaning workers.¹⁵ The vehicle cleaning sector (Standard Occupational Code 53-7061) in California employed nearly 53,000 workers in 2016. While nationally, trends in employment for most of these occupations are projected to move downward through 2028, employment levels in California for nearly all of these occupations are rising substantially (U.S. Bureau of Labor Statistics 2019). For instance, government analysts estimate that building cleaning worker jobs in California will increase by over 15 percent between 2016 and 2026 (California Employment Development Department 2019).

Potential exposures to the Candidate Chemical during the product's life cycle

Reference: CAL. CODE REGS. tit. 22, § 69503.3(b)(4)(A-H) and § 69503.3(b)(3).

Potential exposures to the Candidate Chemical or its degradation products may occur during various product life cycle stages, including manufacturing, use, storage, transportation, waste, and end-of-life management practices. Information on existing regulatory restrictions, product warnings, or other product use precautions designed to reduce potential exposures during the product's life cycle may also be discussed here.

During their life cycle, products used to treat converted textiles or leathers can expose humans and other living organisms to PFASs via several pathways and routes (Figure 3). PFASs are released into air, water, and soil during the manufacture, consumer use, and landfilling, incineration, or recycling of converted textile or leather products (Ahrens et al. 2011b; Oliaei et al. 2013), including via evaporation during the product use phase and when washing treated textiles or leathers (Favreau et. al. 2017). Administrative and engineering controls are unlikely to address exposure concerns during product use and disposal. PFASs released during the manufacture and use of treatments for converted textiles or leathers can enter waterways via direct discharge from manufacturers, release of WWTP effluent, or

¹⁵ Additional occupations that risk exposure to treatments containing PFASs include carpet installers and automotive service technicians and mechanics. The U.S. Bureau of Labor Statistics reported in 2016 that California led the nation in the number of carpet installers employed in that sector, with 5,270 workers (U.S. Bureau of Labor Statistics 2016).

following land application of contaminated biosolids (Sepulvado et al. 2011), resulting in PFAS-contaminated surface water and groundwater (Hoffman et al. 2011; Lindstrom et al. 2011a). Most WWTPs cannot adequately remove PFASs, making them a common source of PFASs to surface waters (Dauchy et al. 2012; Guo et al. 2010; Rayne and Forest 2009).

PFASs are released into air, water, and soil during stain- and soil-repellent manufacturing, consumer use, and landfilling of treatments used in converted textile or leather products.




Figure 3: Key PFAS exposure pathways from treatments for converted textiles or leathers.

Manufacturing

Volatile PFASs can be released into the environment during the manufacture of treatments for textiles or leathers (Stock et al. 2004). Manufacturing wastewater discharges into WWTPs can contaminate rivers and other waterways (Konwick et al. 2008). Once released into air and water, PFAAs and some of their precursors are highly mobile and undergo long-range transport to distant regions via atmospheric circulation and oceanic currents (Prevedouros et al. 2006; Ahrens et al. 2011a; Muir et al. 2019).

PFAAs and their precursors can contaminate soil and sediments via deposition of contaminated particulate matter, use of contaminated irrigation water, and land application of contaminated biosolids (Armitage et al. 2009; Blaine et al. 2014a; Lindstrom et al. 2011a; Sepulvado et al. 2011; Zhang et al. 2016). One study of 262 surface sediment samples from 48 lakes and two reservoirs in China (Qi et al. 2016) found high levels of PFASs (C4 through C14) in sediment correlated with proximity of the sampling locations to industrial activities in food packaging, textile, electroplating, firefighting, semiconductor, precious metals processing, and coating industries.

Use

A Canadian family that routinely treated its carpets with a Scotchguard formulation was found to have very high PFAS levels in household dust and serum. The family's young children had the highest detected PFAS concentration, which is likely attributed to the increased opportunity for exposure via dust ingestion or inhalation (Beesoon et al. 2012). The inhalation exposure pathway to treatment products containing PFASs is particularly significant due to the aerosolization of the product formulations during application (Trudel et al. 2008).

Due to the use of treated textiles and leathers, PFASs are found in home and office air samples, and in the blood of residents and office workers (Beesoon et al. 2012; Fraser et al. 2012, 2013). Human exposure to PFASs occurs via inhalation of volatile PFASs and PFAS-containing fine particles, as well as via incidental ingestion of household and office dust, which concentrates PFASs (Fraser et al. 2012; Harrad et al. 2010; Haug et al. 2011; Tian et al. 2016; Rosati et al. 2008; Shoeib et al. 2005).

PFAAs from converted textile or leather treatments can also adsorb to house dust or be released as tiny particles from surface abrasion during normal use (Rosati et al. 2008; Tian et al. 2016). Knobloch et al. (2012) measured PFASs in vacuum dust samples from 39 homes, finding correlations between the amount of PFNA and PFUnA in dust and the presence of carpeting. A study of young adults in North Carolina found a statistically significant 57 percent increase in PFHxS levels in participants who reported vacuuming less frequently (Siebenaler et al. 2017). Another study of Canadian pregnant women found that levels of PFAA precursors in indoor air and dust correlated with participants' PFOA, PFNA, and PFOS serum levels (Makey et al. 2017). Although incidental dust ingestion is generally considered a minor pathway of exposure to PFASs in adults, it can be significantly higher in toddlers

and young children (Tian et al. 2016; Wu et al. 2015). Karásková et al. (2016) measured 20 PFASs in dust from homes and found that exposure doses via incidental dust ingestion were significantly higher in toddlers than adults. Also, children's mouthing of treated textile and leather articles could lead to PFAS exposure, particularly to the shorter-chain PFAAs (CEC 2017). This indicates an increased potential for adverse impacts in this vulnerable subgroup.

The air in homes, offices, and retail businesses can become contaminated with volatilized FTOHs, which are impurities or intermediate degradation products of the stain and soil repellents used on converted textiles or leathers (Herzke et al. 2012; Bohlin-Nizzetto et al. 2015; Vestergren et al. 2015; Schlummer et al. 2015; Kotthoff et al. 2015). Compared to outdoor air, indoor air can have more than 1,000 times higher levels of FTOHs (Fraser et al. 2012; Müller et al. 2012). Fraser et al. (2012) found that FTOH concentrations in office air predicted PFOA concentrations in the blood of workers.

An air and dust sampling study in several indoor locations in China estimated the daily intake of PFASs for adults (Yao et al. 2018). The study found that exposure via inhalation ranges between 1.04 and 14.1 ng/kg b.w./day, while exposure via incidental dust ingestion ranges between 0.10 and 8.17 ng/kg b.w./day. Biota can also experience potential exposures via environmental media and diet.

Cleaning of treated textile and leather may dispose of PFAS-contaminated wastewater in drains, entering WWTPs in areas served by public water systems. Migration studies have been conducted for textile articles treated with side-chain fluorinated polymers. Shorter-chain PFAAs, due to their greater solubility, migrate more readily from textiles during simulated laundering and when exposed to artificial saliva (CEC 2017). While polymeric PFASs form a chemical bond with the surfaces they are applied to, they have shown a decrease in water repellency as treated products are put through repeated wash cycles; concurrently, higher concentrations of PFASs have been measured in wash water (Knepper et al. 2014). When contaminated wastewater is discharged into residential septic tanks, water soluble organic chemicals like PFAAs may migrate into nearby soil and waterways adjacent to the leach field (Schaidler et al. 2014, 2016). WWTP discharge can have PFOA concentrations up to hundreds of ng/L (Hamid and Li 2016). Discharge of PFAA-contaminated effluent into surface waters can lead to contamination of streams, rivers, and sediments (Ahrens et al. 2011b; Sun et al. 2012).

End-of-life

In landfills, the side-chain fluorinated polymers currently used in most converted textile and leather and their stain- and soil-resistant treatments can degrade to form first nonpolymeric fluorotelomer-based PFASs such as FTOHs, which then further degrade to intermediates such as FTCAs and FTUCAs, and finally to PFAAs (Hamid et al. 2018; Washington et al. 2015). The degradation of these side-chain fluorinated polymers in landfills and other waste stocks can be a significant long-term source (years to centuries) of PFAAs to the environment (Lang et al. 2017; Li et al. 2017; Washington et al. 2019).

Estimated release rates from waste stocks are almost two orders of magnitude higher for shorter-chain PFCA, due to their greater mobility in water, compared to longer-chain PFCA (Li et al. 2017, 2020).

Landfills have been identified as a source of PFAS contamination to nearby air and plant leaves (Tian et al. 2018). PFAAs and their precursors from discarded treatment containers and treated textile and leather products can be released into the atmosphere during decomposition in landfills, and become widely dispersed (Ahrens et al. 2011b). PFAAs are also mobilized in landfill leachate (Fuertes et al. 2017; Lang et al. 2016; Shoaieioskouei 2012). Contaminated leachate collected from lined landfills for treatment at WWTPs can result in contaminated waterways, as wastewater treatment does not completely remove PFAAs from the effluent (Hamid and Li 2016; Huset et al. 2011).

PFAAs and their precursors have been detected in landfill leachate all over the world, including Germany (Busch et al. 2010), Norway (Knutsen et al. 2019), Canada (Benskin et al. 2012a), Uganda (Dalahmeh et al. 2018) and Australia (Gallen et al. 2017; Hepburn et al. 2019). In the United States, an estimated 61.1 million cubic meters of leachate was generated in 2013, containing approximately 600 kg total of 70 PFASs measured (Lang et al. 2017). 5:3 FTCA, an intermediate in the degradation of fluorotelomer-based substances to PFAAs, showed the highest mean concentration in leachate from surveyed landfills, followed by PFHxA; the following PFASs were detected in similar amounts: PFBA, PFPeA, PFHpA, PFOA, and 6:2 FTCA (Lang et al. 2017). An evaluation of PFASs leaching from carpet under simulated landfill conditions found increased concentrations leaching over time (Kim et al. 2015). The relatively slow release of PFASs from landfills compared to input rates means that landfilled waste continues to release PFASs for more than a decade or longer (Lang et al. 2017; Washington et al. 2019).

The relatively slow release of PFASs from landfills compared to input rates means that landfilled waste continues to release PFASs for more than a decade or longer.

Furthermore, biosolids from WWTPs can be heavily contaminated with PFAAs and precursors that adsorb to organic matter (Arvaniti et al. 2012; Gallen et al. 2018; Kim Lazcano et al. 2019; Sun et al. 2011). When these materials are land-applied or landfilled, shorter-chain PFASs can migrate from biosolids into soil and leach into adjacent waterways (Sepulvado et al. 2011), potentially leading to widespread contamination and ecological impacts. Land application of biosolids contaminated with PFAAs and their precursors can lead to bioaccumulation of PFAAs in grasses (Yoo et al. 2011) and food crops (Blaine et al. 2014a; Lee et al. 2014). PFAA precursors such as those used in treatments for converted textiles or leathers can also undergo transformation to PFAAs in soils (Washington et al. 2015).

Waste incineration may also release PFASs and other hazardous chemicals to the environment. A literature survey regarding the combustion of various fluorinated polymers reported emissions of C3-C14 PFCAs, ozone depleting substances such as CFCs, and greenhouse gases such as fluorocarbons (Huber et al. 2009). A 2015 study found that, even at temperatures above 400°C and in the presence of calcium hydroxide, PTFE does not undergo complete mineralization (Wang et al. 2015).

Recycling of postconsumer converted textiles or leathers such as carpet, upholstery, and clothing is a preferred alternative to landfilling or incineration, but can perpetuate the presence of PFASs in products made from the recycled material (Herzke et al. 2012). Carpet recycling includes several processing steps that generate high amounts of dust, such as hammer milling, shearing, and shredding (CARE 2012). Workers engaged in recycling activities, including waste handlers, could be exposed to PFASs in dust, while nearby residents could be exposed via off-site dust migration (HBN 2017).

Aggregate effects

Reference: CAL. CODE REGS. tit. 22, § 69503.3(a)(1)(B) and § 69503.3(b)(3).

Multiple sources of exposure to the Candidate Chemical may increase the potential for significant or widespread adverse impacts.

PFASs are found in many frequently used consumer products in Californian households and workplaces, including carpets, rugs, flooring, furniture, clothing, cookware, food packaging, cleaning products, personal care products, and electronics (KEMI 2015), as well as in food (Christensen et al. 2017) and drinking water (Schwanz et al. 2016; Boone et al. 2019; Domingo and Nadal 2019). This can result in exposure to complex PFAS mixtures from multiple consumer products, including treatments for converted textiles or leathers (Bečanová et al. 2016). DTSC considers all these potential sources of PFAS exposure under its mission to protect the people and biota of California.

Because of the prevalence of stain- and soil-repellent carpets and rugs in residential and commercial spaces, all Californians are potentially exposed, directly or indirectly, to PFASs from these products. U.S. Census data from 2012 identifies more than 850 stores in California selling carpets and rugs, representing a potential source of increased exposure to PFASs for thousands of retail workers (U.S. Census Bureau 2017b).¹⁶ The Chinese textile industry is estimated to use about 10,000 metric tons of finishing treatments containing fluorinated chemicals (Lim et al. 2011). These materials and products

¹⁶ We performed an advanced search by Industry Codes using the 2012 NAICS code “442210: Floor covering stores,” then refined the search results by limiting them to California, and selecting the document ID EC1244SLLS11 (“Retail Trade: Subject Series – Product Lines: Product Lines Statistics by Industry for the U.S. and States: 2012”). The table lists 852 establishments under Products and Services Codes 20365 (soft-surface (textile) floor coverings to be installed),” and 641 establishments under Products and Services Code 20366 (carpets and rugs not requiring installation).

are then exported globally. High PFAS concentrations have been detected in textile products originating from China, including cotton and leather clothing, bedding, curtains, furniture coverings, and most notably carpeting (Vestergren et al. 2015). Paul et al. (2009) estimated that 14 to 48 percent of the PFASs produced globally are used by the carpet industry. The amount of PFASs in treated carpets and rugs varies widely, depending on the carpet and its age. Stain and soil repellents can contribute approximately 15 percent of fiber weight in synthetic carpets (KEMI 2015).

Despite extensive documentation of the widespread environmental contamination with PFASs from consumer products such as treatments for converted textiles or leathers, longitudinal studies that evaluate aggregate exposures to these chemicals from all sources through different life stages are still lacking. A lack of comprehensive data on source emissions and the distribution of PFASs through all environmental compartments makes it challenging to accurately estimate aggregate exposures over a lifetime. The frequency, level, and duration of individual or population-level PFAS exposures associated with the use of treatments for converted textiles or leathers have not been estimated using either deterministic or probabilistic exposure modeling.

Indicators of potential exposures to the Candidate Chemical in the product

Reference: CAL. CODE REGS. tit. 22, § 69503.3(b)(2).

The SCP regulations consider various data that indicate potential for exposure to the Candidate Chemical or its degradation products, including: the Candidate Chemical's presence in and release from the product; monitoring data indicating the Candidate Chemical's presence in the indoor and outdoor environment, biota, humans (e.g., biomonitoring studies), human food, drinking water, and other media; and evidence of persistence, bioaccumulation, lactational and transplacental transfer.

PFASs are found ubiquitously in the environment, plants, animals, and humans (Campo et al. 2016; Lindstrom et al. 2011a, b). Treatments for converted textiles or leathers contribute to widespread environmental contamination and exposures, as do other consumer products such as carpets, rugs, furniture, food packaging, cosmetics, and waterproof clothing. Once released to the environment during product manufacture, use, or disposal, PFASs become part of a virtually closed cycle leading to chronic human and ecological exposures (Figure 3). Because PFAAs and other persistent PFASs lack a natural degradation route, their levels in the environment, humans, and biota may continue to rise for as long as PFASs are used in consumer products.

Between 2009 and 2017, 455 new PFASs have been detected in environmental media and commercial products. However, most of the PFASs that contribute to total organic fluorine in the environment, wildlife, and human blood samples remain unidentified due to analytical limitations (Xiao 2017). Thus, only a small fraction (sometimes less than 5 percent) of the PFASs in these media are likely reflected in the data summarized in this section. Nakayama et al. (2019) discussed the available analytical methods

for measuring PFASs in air, water, sediments, soil, sludge, dust, and biological samples and identified a great need for new methods that expand the suite of PFASs studied, especially in humans and wildlife. As of now, the full extent of the contamination, despite extensive research, remains poorly understood.

Once released to the environment during product manufacture, use, or disposal, PFASs become part of a virtually closed cycle leading to chronic human and ecological exposures. Because PFAAs and other persistent PFASs lack a natural degradation route, their levels in the environment, humans, and biota may continue to rise for as long as PFASs are used in consumer products.

Evidence that the Candidate Chemical is present in or released from the product

Accurate information on the current production volume of PFASs used in domestically produced and imported treatments is lacking. Using data from the Danish Product Register, the Danish EPA calculated the total content of fluorinated compounds in treatments for shoes to be between 0.03 and 60 g/kg (Jensen et al. 2008). Another study conducted on several product categories detected PFAS in sprays for fabrics and textiles (Ye et al. 2015).

Although only polymeric PFASs are currently used in treatments for converted textiles or leathers (Bowman 2017; FluoroCouncil 2017), nonpolymeric PFASs can be present as manufacturing impurities or degradation products (Kotthoff et al. 2015; Vestergren et al. 2015). FTOHs, which are used as manufacturing intermediates, are also present as impurities or degradation intermediates of side-chain fluorinated polymers and can off-gas from fluorotelomer-based products (Dinglasan-Panlilio and Mabury 2006; Sinclair et al. 2007). Studies in Europe and Canada found FTOH levels as high as 9,420 ng/L, and the sum of PFCAs as high as 8.4 ng/L in treatments for converted textiles or leathers (Jensen et al. 2008). A sampling of textiles used as materials for outdoor products showed a PFHxA concentration of up to 17,100 ng/m², while leather samples exhibited up to 241,800 ng/m² PFBA and 143,000 ng/m² PFBS (Kotthoff et al. 2015). Carpet/fabric care liquids and foams in the United States in 2011 and 2013 were found to contain up to 105,000 ng/g 6:2 FTOH, 2,950 ng/g 8:2 FTOH, and 1,460 ng/g 10:2 FTOH (Liu et al. 2015c). Treated home textiles and upholstery were found to contain up to 40,900 ng/g 6:2 FTOH, 21,200 ng/g 8:2 FTOH, and 12,100 ng/g 10:2 FTOH (Liu et al. 2015c).

Until relatively recently, the side-chain fluorinated polymers used in treatments for converted textiles or leathers were made with longer-chain PFASs (more than seven carbon atoms) and included impurities such as PFOS, PFOA, and 8:2 FTOH (Jensen et al. 2008). Because of these side chains' potential to cause adverse health effects, PFOS-based substances were phased out by major U.S.

producers over a decade ago (Prevedouros et al. 2006). Currently, domestically produced treatments are made with shorter-chain side-chain fluorinated polymers, mainly urethane-based polymers and some (meth)acrylates. These degrade to short-chain FTOHs, such as 6:2 FTOH, which are more volatile than the longer-chain ones and thus more likely to be released from products (U.S. EPA 2017a).

Monitoring data showing that the Candidate Chemical is widespread in the indoor and outdoor environment and accumulates in terrestrial and aquatic organisms

Numerous published studies have found that PFASs are ubiquitous in various types of environmental media, in biota, and in the indoor environment. A whole suite of PFAAs and precursors, including FTOHs, FTOs, FTCAs, FTUCAs, FASAs, FASEs, FTSs, fluorotelomer phosphate diesters (diPAPs), perfluoroalkyl iodides (PFAIs), FTIs, and perfluoroalkyl sulfonates (PFSIs) were found in various matrices such as air, WWTP effluent and sludge, landfill leachate, seawater, rivers, lakes, rain, snow, ice, and soil [see (Frömel et al. 2016) and Table 3 for examples]. Despite the phaseout of longer-chain PFASs started more than a decade ago, concentrations in environmental samples and biota have not declined (Land et al. 2018). This indicates that ecosystems continue to be exposed to previously released PFASs for a long time.

Table 3: Monitoring studies found varying levels of PFASs in the environment and biota.

Sample Type	PFAS Type: Concentration	Reference
Indoor air (homes)	6:2 FTOH: 1.8 ng/m ³ (median) 8:2 FTOH: 8.7 ng/m ³ (median) 10:2 FTOH: 2.5 ng/m ³ (median) 8:2 FTAC: 0.27 ng/m ³ (median) 10:2 FTAC: 0.12 ng/m ³ (median)	Fromme et al. (2015)
Indoor air (homes)	Sum of FTOHs: 3.21 ng/m ³ (median) Sum of longer-chain PFCAs: 0.34 ng/m ³ (median) Sum of shorter-chain PFCAs: 0.35 ng/m ³ (median) Sum of PFSAs: 0.23 ng/m ³ (median) Sum of diPAPs: 0.0012 ng/m ³ (median)	Yao et al. (2018)
Indoor air (hotels)	Sum of FTOHs: 3.33 ng/m ³ (median) Sum of longer-chain PFCAs: 0.21 ng/m ³ (median) Sum of shorter-chain PFCAs: 0.26 ng/m ³ (median) Sum of PFSAs: 0.12 ng/m ³ (median) Sum of FOSE/FOSAs: 0.0053 ng/m ³ (median) Sum of diPAPs: 0.0011 ng/m ³ (median)	Yao et al. (2018)

Sample Type	PFAS Type: Concentration	Reference
Indoor air (schools)	6:2 FTOH: 3.3 ng/m ³ (median) 8:2 FTOH: 4.4 ng/m ³ (median) 10:2 FTOH: 1.8 ng/m ³ (median) 8:2 FTAC: 0.45 ng/m ³ (median) 10:2 FTAC: 0.31 ng/m ³ (median)	Fromme et al. (2015)
Indoor air (offices)	6:2 FTOH: 1.3 ng/m ³ (geometric mean) 8:2 FTOH: 9.9 ng/m ³ (geometric mean) 10:2 FTOH: 2.9 ng/m ³ (geometric mean)	Fraser et al. (2012)
Indoor dust (homes)	Sum of nine PFCAs (C6 to C14): <1.0 – 37,400 µg/kg	Liu et al. (2011b)
Indoor dust (homes)	Sum of longer-chain PFCAs: 59.5 µg/kg (median) Sum of shorter-chain PFCAs: 87.4 µg/kg (median) Sum of PFSAAs: 17.1 µg/kg (median) Sum of diPAPs: 1.13 µg/kg (median)	Yao et al. (2018)
Indoor dust (hotels)	Sum of FTOHs: 149 µg/kg (median) Sum of longer-chain PFCAs: 25.8 µg/kg (median) Sum of shorter-chain PFCAs: 3.25 µg/kg (median) Sum of PFSAAs: 10.3 µg/kg (median) Sum of FOSE/FOSAs: 2.1 µg/kg (median) Sum of diPAPs: 10.2 µg/kg (median)	Yao et al. (2018)
Indoor dust (day care center)	PFOA: 31 – 110 µg/kg PFOS: 23 – 65 µg/kg	Björklund et al. (2009)
Outdoor air	FTOHs: 0.06 – 0.19 ng/m ³	Müller et al. (2012)
North American soils	Sum of PFCAs: 145 – 6,080 ng/kg Sum of PFSAAs: 35 – 1,990 ng/kg	Rankin et al. (2016)
Sediments (South Bay Area, California)	PFOS: 0.60 – 2.61 ng/g dry weight	Sedlak et al. (2017)
Sediments (Cape Fear River, North Carolina)	GenX (PFPrOPrA): 3.1 – 21.6 ng/g dry weight	UNCW (2018)
U.S. river (Cape Fear, North Carolina), used for drinking water	GenX (PFPrOPrA): 631 ng/L (mean)	Sun et al. (2016)
WWTP effluent (San Francisco Bay Area, California)	PFBA: 16 ng/L (mean) PFPeA: 12 ng/L (mean) PFHxA: 26 ng/L (mean) PFBS: 2.7 ng/L (mean) PFHxS: 4.8 ng/L (mean) PFOS: 13 ng/L (mean)	Houtz et al. (2016)

Sample Type	PFAS Type: Concentration	Reference
WWTP effluent (Germany)	PFBA: <4.2 ng/L PFPeA: <254 ng/L	Frömel et al. (2016)
Sewage sludge	Sum of fourteen PFAAs: 126 – 809 µg/kg	Yan et al. (2012)
Fish (whole fish from the Ohio, Missouri, and Upper Mississippi Rivers)	Sum of ten PFAAs: 5.9 – 1,270 µg/kg; 53.4 µg/kg (median wet weight)	Ye et al. (2008)
Prey fish (San Francisco Bay, California)	PFOS: 11.8 µg/kg (geometric mean wet weight)	Sedlak et al. (2017)
Cormorant eggs (San Francisco Bay, California)	PFOS: 36.1 – 466 µg/kg (wet weight)	Sedlak et al. (2017)
Harbor seals serum (San Francisco Bay, California)	PFOS: 12.6 – 796 µg/kg	Sedlak et al. (2017)
Arctic lichen and plants	Sum of six PFCAs (C8 to C13) ¹⁷ : 0.02 – 0.26 µg/kg	Müller et al. (2011)
Arctic caribou (liver)	Sum of six PFCAs (C8 to C13): 6 – 10 µg/kg	Müller et al. (2011)
Arctic wolf (liver)	Sum of six PFCAs (C8 to C13): 10 – 18 µg/kg	Müller et al. (2011)
Arctic reindeer	PFBA: 0.079-14 µg/kg	Ahrens et al. (2016b)

Air and dust: PFASs are ubiquitous in air and dust. All indoor air and dust samples collected in 2015 from residences, hotels, outdoor equipment stores, curtain stores, and carpet stores in China contained at least one PFAA or precursor (FTOHs, FOSE/FOSAs, or diPAPs) (Yao et al. 2018). A recent study identified 34 emerging PFASs (including PFECAs and short-chain PFAA precursors) in indoor and outdoor airborne particulate matter from five Chinese cities (Yu et al. 2018). PFASs were found in indoor dust in homes sampled on four continents, as well as in British cars, classrooms, and offices (Goosey and Harrad 2011). A review and analysis of literature data published after 2010 concluded that neutral PFASs, mainly FTOHs, FASAs, and FASEs, are most dominant in indoor air due to their low water solubility and high vapor pressure (Jian et al. 2017). 6:2 FTAC and 6:2 FTMAC were also frequently

¹⁷ C8 to C13 refers to the number of carbon atoms in the PFCA molecule. This range thus covers: PFOA, PFNA, PFDA, PFUnA, PFDoA, PFTrDA.

detected, sometimes at high concentrations, in children's bedroom air sampled in Finland during 2014/2015 (Winkens et al. 2017a). 6:2, 8:2, and 10:2 FTOHs have been detected in indoor and outdoor ambient air (Jahnke et al. 2007; Karásková et al. 2016; Kim and Kannan 2007; Liu et al. 2013; Shoeib et al. 2011; Tian et al. 2016). The estimated atmospheric lifetime of FTOHs indicates that atmospheric transport can contribute to widespread human exposures and environmental burdens (Ellis et al. 2003). Human exposure to PFAAs, FASAs, FOSEs, and other PFASs can also occur through house dust (Jian et al. 2017), which can have significantly higher PFAS levels than background concentrations in urban soils (Tian et al. 2016; Xu et al. 2013). PFAA precursors in air and dust were observed to correlate with PFAA levels in human serum (Makey et al. 2017). Household dust can contain hundreds of chemicals (Moschet et al. 2018), including some with hazard traits similar to those of PFASs, such as flame retardants, phthalates, and environmental phenols (Mitro et al. 2016). Because Americans spend, on average, more than 90 percent of their time indoors (Klepeis et al. 2001), incidental ingestion and inhalation of indoor dust represent potentially significant exposure pathways to multiple contaminants along with PFASs.

Surface water and groundwater: PFASs have been detected in creeks and rivers (D'eon et al. 2009), lakes (De Silva et al. 2011), glacial meltwater and snow (Skaar et al. 2018), oceans (Benskin et al. 2012b; González-Gaya et al. 2019), and groundwater (Eschauzier et al. 2013; Xiao et al. 2015). Surface waters can contain high levels of PFASs, especially downstream of industrial facilities such as manufacturing sites (Konwick et al. 2008). In fresh water and coastal water, PFHxA and PFHpA have been reported at levels comparable to or higher than PFOA (Cousins et al. 2011). PFPAAs have also been detected in surface water (D'eon et al. 2009). In the San Francisco Bay Area of California, PFASs were detected in bay water (Sedlak et al. 2017) and groundwater, with concentrations ranging from 19 to 192 ng/L (PFOS), <LOD to 22 ng/L (PFOA), and <20 ng/L (PFHxS, PFDS, PFDA, and two PFOS precursors) (Plumlee et al. 2008). In the raw water of a drinking water treatment plant in the Cape Fear River watershed, North Carolina, Sun et al. (2016) found the PFECA GenX, a PFOA replacement in fluoropolymer manufacturing, at high levels (631 ng/L on average, with a maximum of approximately 4,500 ng/L). In a recent U.S. EPA – U.S. Geological Survey study, PFBS and PFOA were detected in the source waters of all 25 drinking water treatment plants tested from across the U.S., while PFHxS, PFOS, PFBA, PFPeA, PFHxA, PFHpA, PFNA, and PFDA were detected at over 90 percent of the sites (Boone et al. 2019). Certain PFECAs and PFESAs were detected in surface waters in all countries on three continents surveyed recently, indicating global distribution and contamination (Pan et al. 2018). The ultra-short-chain PFAS TFA was also detected in seawater, rain, and melted snow on several continents (Ateia et al. 2019b). Contaminated surface waters can be ingested by livestock and wild animals, leading to exposure up the food chain.

WWTP effluent: PFASs are commonly found in municipal and industrial wastewaters (Appleman et al. 2014; Arvaniti et al. 2015; Chen et al. 2017b; Clara et al. 2008; Sun et al. 2012; Coggan et al. 2019). Industrial wastewater can have PFOA and PFOS levels above 1,000 ng/L (Kim et al. 2012b; Lin et al.

2014a). PFPAs have also been detected in WWTP effluent (D'eon et al. 2009). WWTPs are major point sources for PFAS contamination of aquatic environments (Ahrens 2011; Dalahmeh et al. 2018; Frömel et al. 2016; Gallen et al. 2018; Guerra et al. 2014). Degradation of PFAA precursors, such as FTOHs, within WWTPs can lead to increases in effluent PFAA concentrations (Ahrens 2011; Dalahmeh et al. 2018; Frömel et al. 2016; Gallen et al. 2018; Guerra et al. 2014). Despite the phaseout of PFOS- and PFOA-containing stain- and soil-repellent treatments for converted textiles or leathers, these PFASs are frequently detected in wastewater, indicating their ongoing release into the environment (Gallen et al. 2018; Guerra et al. 2014). A study of effluent from WWTPs in the San Francisco Bay Area detected several PFAAs including PFHxA, PFOA, PFBA, and PFOS, with highest median effluent concentrations of 24, 23, 19 and 15 ng/L, respectively (Houtz et al. 2016). A subsequent study (Sedlak et al. 2017) of samples collected in 2012 from three San Francisco Bay WWTPs found mean PFOA, PFOS, PFHxA, and PFPeA concentrations as high as 39.7, 42.3, 30.1, and 21.2 ng/L, respectively.

Landfill leachate: PFASs are commonly detected in landfill leachate around the world (Benskin et al. 2012a; Fuertes et al. 2017; Gallen et al. 2017; Hamid et al. 2018; Knutsen et al. 2019). A study of 95 leachate samples from 18 U.S. landfills estimated the annual release of PFCAs, FTCAs, and PFASs and precursors to be 291 kg/yr, 285 kg/yr, and 84 kg/yr, respectively (Lang et al. 2017). A Swedish study of 26 PFASs in samples from groundwater, surface water, WWTP effluent, and landfill leachate found that landfill leachates had the highest average total PFAS concentrations (487 ng/L) (Ahrens et al. 2016a). Shorter-chain PFAAs have become the most abundant PFASs in landfill leachate, reflecting the relatively recent market shift from longer- to shorter-chain compounds (Hamid et al. 2018; Knutsen et al. 2019). A study of leachate from municipal and industrial landfills in Sweden detected also ultra-short-chain PFASs (with three or fewer carbon atoms) such as trifluoroacetic acid (TFA), perfluoropropanoic acid (PFPrA), and trifluoromethane sulfonic acid (TFMS) (Björnsdotter et al. 2019).

Soils: PFASs are ubiquitous in soils. A survey of soils around the world, including remote locations such as Antarctica, found PFASs in all samples (Llorca et al. 2012; Rankin et al. 2016). Side-chain fluorinated polymers were detected in all biosolid-treated soil samples measured in a Canadian study (Chu and Letcher 2017). Soil and water contaminated with PFASs can lead to plant and soil organism uptake and transfer to grazing animals potentially consumed by humans (Navarro et al. 2017; Yoo et al. 2011). An evaluation of agricultural lands in Kampala, Uganda, found PFASs in the terrestrial food chain, including in plants such as maize, yam, and sugar cane (Dalahmeh et al. 2018). Some plant species show a very high PFAS accumulation potential, with higher BCFs for shorter-chain PFASs. For example, the BCF (plant/soil ratio) of 6:2 FTSA in birch leaves can be up to 143,700 (Gobelius 2016). Based on their ability to uptake and sequester PFASs in aboveground biomass, some plants have been proposed for use in the phytoremediation of PFAS-contaminated soil (Gobelius 2016).

Sediments: Sediments can also become contaminated. (White et al. 2015) analyzed PFASs in estuarine sediments from the Charleston Harbor and the Ashley and Cooper rivers (n=36) in South Carolina,

finding higher concentrations of 11 PFASs than previously reported for U.S. urban areas. The PFASs were dominated by PFOS, followed by PFDA and PFOA (White et al. 2015). Eighteen PFCAs and PFSAs were detected in the archived surface sediments of five major rivers in China (Pan et al. 2014a). Consistent with other studies of sediments from lakes, reservoirs, and estuaries, the most frequently detected PFASs were PFOA and PFOS (detection frequency of 100 and 83 percent, respectively), with greater concentrations in urban areas (Pan et al. 2014b). PFPAs and PFPIAs have been detected in lake sediments, but at lower concentrations compared to other PFAAs (Guo et al. 2016). A recent study found that Cape Fear River sediments appear to act as a repository for GenX, potentially releasing it into the water column (UNCW 2018). Concentrations of PFAAs, including PFHxA and PFBA, have been found at higher concentrations at ocean depths than at the surface, indicating the potential for deep ocean sediments to serve as the ultimate sink for PFAAs that adsorb to particulate matter (Prevedouros et al. 2006; Sanchez-Vidal et al. 2015).

PFAAs are found in numerous marine species, including plankton, sea turtles, seals, California sea otters, whales, fish, sharks, polar bears, dolphins, and marine bird eggs; this indicates potential for widespread adverse impacts throughout the marine food web.

Biota: PFAAs are found in numerous marine species, including plankton (Casal et al. 2017), sea turtles (Keller et al. 2005, 2012; O’Connell et al. 2010), seals (Routti et al. 2016), California sea otters (Kannan et al. 2006), whales (Hart et al. 2008), fish (Wong et al. 2017), penguins (Llorca et al. 2012) sharks (Kumar et al. 2009), polar bears (Dietz et al. 2008; Liu et al. 2018b; Tartu et al. 2018), dolphins (Adams et al. 2008; Houde et al. 2005), seabirds (Escoruela et al. 2018; Roscales et al. 2019) and marine bird eggs (Verreault et al. 2007), including PFPIAs in the blood of fish, birds, and dolphins (De Silva et al. 2016). This indicates potential for widespread adverse impacts throughout the marine food web. The Washington State Department of Ecology surveyed regional lakes and analyzed water, fish tissue, and osprey eggs for PFAS contamination, finding PFOS biomagnification through trophic levels by orders of magnitude. In this same report, bioaccumulation of shorter-chain PFASs was not observed (State of Washington 2017).

As shorter-chain PFAA precursors have gained market share, the concentrations of shorter-chain PFAAs in the marine environment have increased, as evidenced by increasing PFBS levels in cetaceans from 2002-2014 (Lam et al. 2016). The PFBS precursor perfluorobutane sulfonamide (FBSA) also bioaccumulates in fish (Chu et al. 2016). 6:2 Cl-PFESA, currently used as a PFOS substitute in the chrome plating industry, has been detected in a wide range of marine organisms, including several species of gastropods, bivalves, crabs, shrimps, cephalopods, and fish, and tends to magnify along the food chain (Liu et al. 2017). 6:2 Cl-PFESA and hexafluoropropylene oxide trimer acid (a PFOA replacement) accumulate in frogs, particularly in male frog skin, liver, and muscle, and in female frog ovaries, raising concerns regarding potential developmental toxicity (Cui et al. 2018).

PFCA are found in Arctic seals and Antarctic seabirds, presumably due to long-range transport of precursors such as FTOHs and FASAs via oceanic and atmospheric currents (Llorca et al. 2012; Routti et al. 2016; Roscales et al. 2019). Increased coastal PFAS concentrations can follow ice and snow melt in the Arctic, resulting in an increased exposure potential for marine biota in Arctic coastal regions (Zhao et al. 2012). Numerous studies have documented high concentrations of PFASs in Arctic mammals. For instance, up to 17 PFASs were found in the livers of marine mammals off the coast of Greenland, including killer whales, polar bears, and ringed seals (Gebbinck et al. 2016). PFASs can be transported across the brain blood barrier and accumulate in animal brain tissue, leading to concerns about potential neurotoxicity (Greaves et al. 2013). A study of 128 young polar bears (between 3 and 5 years old) sampled between 1984 and 2006 estimated that longer-chain PFASs in polar bears will reach levels associated with adverse impacts in lab animals by 2014-2024 (Dietz et al. 2008). PFASs were found to biomagnify throughout the Arctic food chain, with increasing concentrations in wolves that consume caribou contaminated with PFASs via their lichen-based diet (Müller et al. 2011). Researchers also evaluated the effect of cyclic seasonal emaciation on the distribution and composition of PFASs in Arctic foxes, finding significant differences that could indicate increased potential for health effects from PFASs when foxes have reduced access to food (Aas et al. 2014).

Monitoring data showing that the Candidate Chemical is present in human food

Exposure modeling studies have concluded that the ingestion of contaminated food and drinking water is the largest contributor of PFAAs to the overall human body burden (Trudel et al. 2008). A study of 2013-2014 National Health and Nutrition Examination Survey (NHANES) data concluded that diet is the dominant and most important source of PFAA exposure for children aged 3 to 11, even more so than for adolescents and adults (Jain 2018).

Contamination of food with PFASs occurs via contact with PFAS-treated food packaging materials (Trier et al. 2011), as well as from plant and animal uptake from contaminated soil and water (Dalahmeh et al. 2018). Animal feed can also become contaminated when grown in soil treated with PFAS-containing irrigation water or biosolids (Gobelius 2016; Lee et al. 2014; Yoo et al. 2011).

The use of PFAS-contaminated irrigation water has been linked to elevated PFAA concentrations in fruits and vegetables (Blaine et al. 2014b; Zhang et al. 2016; Scher et al. 2018). The application of PFAS-contaminated biosolids to land leads to the bioaccumulation of PFAAs in grasses (Yoo et al. 2011) and food crops (Blaine et al. 2014a; Navarro et al. 2017).

The PFASs currently used in U.S. treatments for converted textiles or leathers degrade into shorter-chain PFAAs that show greater bioaccumulation in vegetable crops than the longer-chain PFAAs from older treatments (Blaine et al. 2013). Composted biosolid samples from the U.S. and Canada had three times higher levels of PFHxA in 2018 compared to 2016, while PFOA and PFOS remained nearly the

same (Kim Lazcano et al. 2019). The study authors interpreted this finding as a reflection of the recent market shift to shorter-chain compounds.

Studies of PFASs in foods (see examples in Table 4) found them in wide-ranging concentrations (Domingo and Nadal 2017; EFSA 2012). Most of the available studies are from Europe and Asia, but levels are expected to be similar in the United States, including in California. From 2006-2012, the European Food Safety Agency (EFSA 2012) tested 7,560 food samples and found PFASs in most of them. The most prevalent PFAS contaminants were PFOS (29 percent) and PFOA (9 percent). PFBS and PFHxA were found to accumulate in food crops, including sugar beets, broccoli, spinach, lettuce, celery, potato, pea, fruit, and green beans (EFSA 2012). A review and analysis of literature data published after 2010 found that shorter-chain PFAAs, including PFBA, PFPeA, and PFHxA, were frequently detected at high concentrations in vegetables, fruits, and beverages (Jian et al. 2017). PFAAs, including longer- and shorter-chains, were found to accumulate in the roots and shoots of wheat (Zhao et al. 2018; Zhang et al. 2019b). Fish appears to be the most frequently PFAS-contaminated food (Yamada et al. 2014a, b), estimated to contribute the most to dietary exposure (Domingo and Nadal 2017; EFSA 2012). One study measured PFASs in 21 foods and beverages in Norway, reporting concentrations of PFOS in cod and cod liver that were much higher than in other foods, including other types of fish (Haug et al. 2010). Nevertheless, terrestrial animals can also have high PFAS concentrations (EFSA 2012; Haug et al. 2010). For instance, several PFAAs were found in the liver and muscle of beef cattle in China (Wang et al. 2017a).

The PFASs currently used in U.S. treatments for converted textiles or leathers degrade into shorter-chain PFAAs that show greater bioaccumulation in vegetable crops than the longer-chain PFAAs from older treatments.

A 2012 European Union (EU) study estimated that PFOA and PFOS dietary exposure doses were much lower than the tolerable daily intake (TDI) established by EFSA in 2008. Dietary exposure to PFOS in toddlers was estimated to contribute almost 20 percent of the TDI in the most exposed group (EFSA 2012). The TDI was based on a sub-chronic study in *Cynomolgus* monkeys, which did not consider exposure sources other than dietary, nor exposure to PFOA and PFOS precursors (EFSA 2008). In 2018, however, EFSA proposed to lower its TDI from 150 ng/kg body weight per day to 13 ng/kg body weight per week for PFOS, and from 1,500 ng/kg body weight per day to 6 ng/kg body weight per week for PFOA (EFSA 2018). According to EFSA, “exposure of a considerable proportion of the population” exceeds the proposed limits for both compounds (Knutsen et al. 2018). TDIs for other PFASs have not been established.

Dietary exposure studies in the United States are generally lacking (Domingo and Nadal 2017), although one study in California found significant positive associations between PFAS serum

concentrations for children and adults and consumption of certain foods including butter/margarine, fish, meat products, and microwave popcorn (Wu et al. 2015). The globalization of the human food chain, combined with widespread contamination of foods, indicates the potential for lifetime exposure to PFASs for the people of California.

Table 4: Monitoring studies found varying levels of PFASs in human food.

Sample Type	PFAS Type	Concentration	Reference
Potatoes	PFOA	0.07 µg/kg	Schechter et al. (2010)
Vegetables	PFHxA	<LOD – 0.42 µg/kg	EFSA (2012)
Fruit	PFHxA	<LOD – 0.17 µg/kg	EFSA (2012)
Fruit	PFBS	<LOD – 0.067 µg/kg	EFSA (2012)
Fruit	PFHxS	<LOD – 0.20 µg/kg	EFSA (2012)
Meat products	PFOA	<LOD – 0.24 µg/kg	Schechter et al. (2010)
Crustaceans	PFOA	<LOD – 8.0 µg/kg	EFSA (2012)
Fish	PFOA	<LOD – 0.30 µg/kg	Schechter et al. (2010)
Fish	PFHxA	<LOD – 23 µg/kg	EFSA (2012)
Fish	PFOA	<LOD – 18.2 µg/kg	EFSA (2012)
Fish	PFOS	<1 ng/g – >100 µg/kg	Berger et al. (2009)
Fish and other seafood ¹⁸	PFOS	<LOD – 310 µg/kg	EFSA (2012)
Eggs	Sum of eleven PFASs	27 – 160 µg/kg	Wang et al. (2008)
Eggs and products	PFOA	<LOD – 25.5 µg/kg	EFSA (2012)
Eggs and products	PFOS	<LOD – 6.4 µg/kg	EFSA (2012)
Home produced eggs (yolks) two km from industrial site	Sum of twelve PFAAs	8.99 – 482 µg/kg	Su et al. (2017)
Butter	PFOA	1.07 µg/kg	Schechter et al. (2010)
Margarine	PFOA	0.19 µg/kg	Schechter et al. (2010)
Olive oil	PFOA	1.80 µg/kg	Schechter et al. (2010)
Honey	PFOA	0.00 – 0.47 µg/kg	EFSA (2012)

¹⁸ PFOS mean concentrations were “constantly higher” in freshwater fish than in marine fish.

Monitoring data showing that the Candidate Chemical is present in drinking water

PFASs are detected in drinking water around the world, typically in the nanogram per liter range (Domingo and Nadal 2019). Drinking water can become contaminated with PFAAs and their precursors from treatments for converted textile or leather via several routes, including direct discharge of industrial effluent into waterways, leaching of land-applied wastewater or biosolids into groundwater, leachate draining from landfills that contain discarded products, and discharge of WWTP effluent into streams and rivers (Lindstrom et al. 2011a). A review and analysis of literature data published after 2010 found that, out of all PFASs measured to date, PFOS, PFHxA, PFHpA, PFNA, PFDA, and PFOA had the highest concentrations and detection frequencies in drinking water (Jian et al. 2017). A study of 133 PFASs in drinking water (bottled and tap) from nine countries, including the United States, found shorter-chain PFAAs in 64 to 92 percent of the samples (Kaboré et al. 2018).

Drinking water can become contaminated with PFAAs and their precursors from treatments for converted textile or leather stain- and soil-repellents via several routes, including direct discharge of industrial effluent into waterways, leaching of land-applied wastewater or biosolids into groundwater, leachate draining from landfills that contain discarded products, and discharge of WWTP effluent into streams and rivers.

As part of the third Unregulated Contaminant Monitoring Rule (UCMR 3) program, U.S. EPA (2017b) monitored public water supplies for six PFAAs from 2013 until 2015, reporting their presence in approximately 3 percent of samples tested (Table 5). PFOA and PFOS concentrations equaled or exceeded the lifetime total health advisory of 70 ng/L, combined, in water serving approximately 16.5 million people, with most detects (28 out of 452 samples, or 6.2 percent detection rate) in California (Hu et al. 2016; Hurley et al. 2016). The most significant predictors of observed PFAS detection frequencies and concentrations were: (1) proximity to industrial sites that manufacture or use PFASs; (2) proximity to military fire training areas; and (3) the number of nearby WWTPs (Hu et al. 2016).

The Minimum Reporting Levels (MRLs) for the six PFASs measured by the UCMR 3 program ranged from 10 to 90 ng/L, whereas the method used has a limit of quantification in the 2-5 ng/L range. Eurofins Eaton Analytical (2017), which was responsible for analyzing approximately 40 percent of the UCMR 3 data, found PFASs in more than a tenth of the samples it analyzed. However, fewer than 2 percent of those samples showed values above the UCMR 3 MRL and were reported to U.S. EPA. This suggests that the UCMR 3 program may have significantly underestimated the prevalence of PFASs in U.S. drinking water. A U.S. EPA-U.S. Geological Survey joint study (Boone et al. 2019) retested 25 of the drinking water treatment plants sampled under UCMR 3. The authors found PFBS, PFHxA, and PFOA in 100 percent of the treated water samples, and PFPeA, PFHpA, PFNA, and PFOS in over 90 percent of those samples. The shorter-chain PFAAs showed no significant decline between the source and the

treated water (Boone et al. 2019). A recent U.S. EPA literature review found that, in addition to the six PFASs measured during UCMR3, 23 other PFASs in drinking water were reported to date (Crone et al. 2019).

Drinking water contamination can linger for decades. In August 2017, the town of Belmont, Michigan, measured PFASs in some of its drinking water at levels 540 times above the U.S. EPA health advisory level (32,000 ng/L PFOS and 5,800 ng/L PFOA). The contamination is linked to a hazardous waste site used in the 1960s by a former tannery. No PFASs have been added to the site since the 1970s (Chawaga 2017; Ellison 2017).

Table 5: Monitoring studies found varying levels of PFASs in drinking water.

Sample Type	PFAS Type: Concentration	Reference
U.S. drinking water ¹⁹	PFOA: <MRL – 349 ng/L PFOS: <MRL – 1,800 ng/L PFNA: >MRL – 56 ng/L	Hu et al. (2016)
U.S. drinking water	PFBS: <0.032 – 11.9 ng/L PFHxS: ND – 38.4 ng/L PFOS: ND – 36.9 ng/L PFBA: ND – 104 ng/L PFPeA: ND – 514 ng/L PFHxA: 0.09 – 60.8 ng/L PFHpA: ND – 177 ng/L PFOA: <0.56 – 104 ng/L PFNA: ND – 38.6 ng/L PFDA: ND – 24.7 ng/L	Boone et al. (2019)
European drinking water	PFBS: <LOD – 0.24 µg/kg PFHxS: <LOD – 0.011 µg/kg PFOS: <LOD – 0.016 µg/kg	EFSA (2012)

Monitoring data showing that the Candidate Chemical is present in California environmental media at levels of concern

Numerous drinking water sources in California are contaminated with PFAAs, according to the UCMR 3 study. Forty percent of the positive UCMR 3 study detects in California are above the U.S. EPA advisory level of 70 ng/L for PFOA and PFOS combined (Hurley et al. 2016). In October 2019, the California State Water Resources Control Board reported data from 600 water system sites adjacent to airports with

¹⁹ The UCM3 MRLs (Minimum Reporting Levels) for PFASs in U.S. drinking water are 10 – 90 ng/L depending on the PFAS being tested.

fire training areas and municipal solid waste landfills within California (California State Water Resources Control Board 2019). In total, 181 of the water system sites sampled exceeded the state's notification level of 6.5 ppt for PFOS, while 150 samples exceeded the notification level of 5.1 ppt for PFOA. The shorter-chain PFHxA and PFBS were detected in 167 and 185 of the samples, respectively.

California sea otters were found to have PFOA and PFOS levels ranging from less than 5 to 147 ng/g and from less than 1 to 884 ng/g, respectively, significantly correlated with incidence of disease (Kannan et al. 2006). Historically, San Francisco Bay wildlife has shown some of the highest PFOS levels ever measured (Sedlak and Greig 2012). Harbor seal serum sampled in 2014 still contained PFOS at concentrations ranging from 12.6 to 796 ng/g, more than a decade after the PFOS phaseout (Sedlak et al. 2017).

In 2010, contaminant monitoring of the San Francisco Bay surface water detected several PFAAs, including: PFBS at concentrations up to 7.89 ng/L (average 1.58 ng/L), PFBA at up to 62.20 ng/L (average 12.96 ng/L), and PFHxA ranging from 1.37 ng/L to 221.0 ng/L (SFEI 2010). A 2016 study of PFAAs and precursors in wastewater effluent discharged to San Francisco Bay highlighted the local impact of aqueous film-forming foam (AFFF),²⁰ and noted that levels of PFBA, PFPeA, and PFHxA in WWTP effluent had increased by 150 to 220 percent since 2009 (Houtz et al. 2016). Due to this widespread contamination, the San Francisco Bay Regional Monitoring Program designated PFOS as "moderate concern" for the San Francisco Bay, and all other PFASs as "possible concerns" (Sutton et al. 2017).

The Centers for Disease Control and Prevention, which monitors Americans' exposure to PFASs as part of the National Health and Nutrition Examination Survey (NHANES), has detected PFASs in the blood of all people tested.

Human biomonitoring data

Nearly all humans show evidence of exposure to some PFASs (Calafat et al. 2007), which can accumulate in human lungs, kidneys, liver, brain, and bone tissue (Pérez et al. 2013). A recent study found PFOS, PFOA, PFNA, PFDA, PFUnA, and PFHxS in human embryonic and fetal organs, with the highest burden in lung tissue in the first trimester samples, and in the liver in the second and third trimester samples (Mamsen et al. 2019). Biomonitoring studies have been limited to certain PFASs with

²⁰ AFFF may be the largest direct point source of PFASs, according to a Swedish Environmental Protection Agency study (Goldenman 2017). Generally, AFFF contains high levels of PFASs and their precursors, but only low levels of PFCAs (Hu et al. 2016).

available analytical methods (see Table 6 for some examples), however humans are exposed to many more PFASs that haven't yet been measured (Yeung and Mabury 2016).

The Centers for Disease Control and Prevention (CDC), which monitors Americans' exposure to PFASs as part of NHANES, has detected PFASs in the blood of all people tested (Calafat et al. 2007). NHANES blood samples collected in 2011-2012 averaged PFHxS, PFOA, and PFOS between 1.3 and 6.3 µg/L (ATSDR 2017). PFBA and PFHxA were found in 13.3 percent and 22.6 percent, respectively, of 2682 urine samples collected from 2013-2014 NHANES participants (Calafat et al. 2019).

A biomonitoring study of over 65,000 people enrolled in the C8 Health Project, which evaluated exposures to PFASs in drinking water contaminated by DuPont, found PFHxA in the serum of more than 50 percent of the people tested (Frisbee et al. 2009). Shorter-chain PFAAs such as PFHxA, PFPeA, and PFBA often fall below the level of detection in human serum and plasma, but can be found at concentrations higher than most longer-chain PFASs in human whole blood samples (Poothong et al. 2017). They have also been detected in human hair samples at similar or higher levels compared to longer-chain PFAAs (Alves et al. 2015; Ruan et al. 2019; Martín et al. 2019). Shorter-chain PFAAs are also detected in human urine at levels comparable to or higher than PFOA (Hartmann et al. 2017; Kim et al. 2014a; Perez et al. 2012). and may accumulate in certain organs at levels exceeding those of longer-chain PFAAs. In human autopsy tissues, out of 21 PFAAs studied, PFBA was the dominant compound in kidney and lung (median of 263 and 807 ng/g, respectively) and PFHxA in liver and brain (median of 68.3 and 141 ng/g, respectively), whereas PFOA dominated in bone (median of 20.9 ng/g) (Pérez et al. 2013). As the unique behaviors of PFAAs in biological systems and their human toxicokinetics become better understood, CDC may revise its biomonitoring protocols to sample tissues most relevant to assessing exposures to shorter-chain PFAAs (Calafat 2017).

Treatments containing PFASs for use on converted textiles or leathers may impact children more than adults due to their frequent hand-to-mouth behavior, more direct contact with the treated products over a larger body surface area, and other physiological differences.

The California Biomonitoring Program, administered jointly by the California Department of Public Health, DTSC, and OEHHA, has measured PFAA levels in several subpopulations (CECBP 2017). The California Teachers Study is a statewide collaborative research project evaluating health-related factors in the development of breast cancer. For more than 20 years, over 133,000 female schoolteachers and employees have participated in the study. As of 2015, the California Biomonitoring Program has found PFHxS, PFOA, and PFOS in more than 99 percent of 1,759 blood samples provided by study participants (CECBP 2015b). Hurley et al. (2016) analyzed blood samples from a subset of 1,333 participants who lived in areas where PFASs in drinking water concentrations were tested. Of these participants, 109 women lived in an area where at least one PFAS was detected in drinking water from public water

systems. The analysis showed an association between detectable levels of PFOA and PFOS in drinking water and the women’s blood levels (Hurley et al. 2016). In a subset of 1,257 women aged 40 to 94 from the California Teachers Study, serum levels of longer-chain PFASs decreased from 2011 to 2015 on average by 10 to 20 percent per year – except for PFHxS, which did not change significantly (Hurley et al. 2018). Another study of 139 California households detected PFASs in all study participants, including young children and adults (Wu et al. 2015). Exposure levels correlated with diet and use of consumer products such as fire extinguishers and stain-repellent or waterproof clothing, and residential dust exposure in children.

In North Carolina, a study of 37 young adults detected PFHxS (1.07 – 12.55 ng/mL), PFOS (0.39 – 31.35 ng/mL), PFOA (0.30 – 4.07 ng/mL), and PFNA (0.23 – 4.02 ng/mL) in all serum samples, and PFHxA (ND – 1.00 ng/mL) and PFDA (ND – 1.60 ng/mL) in 83.8 percent and 97.3 percent of serum samples, respectively (Siebenaler et al. 2017). In Europe, biomonitoring of pregnant and nursing women from 1996 to 2010 found that PFBS, used in the manufacture of the Scotchgard brand stain and soil repellent, doubled its blood concentrations every 6.3 years, concurrent with the phaseout of PFOS (Glynn et al. 2012).

A study of more than 600 children aged 6 to 10 from Eastern Massachusetts found higher blood concentrations of certain PFAAs and PFAA precursors in those children who had a carpet or rug in their bedroom (Harris et al. 2017). Children may be more impacted than adults by the treatments containing PFASs for use on converted textiles or leathers due to their frequent hand-to-mouth behavior, more direct contact with the treated products over a larger body surface area, and other physiological differences (Moya et al. 2004).

Table 6: Human biomonitoring studies found widespread exposure to PFASs.

Population	Sample Type	PFAS Type: Concentration	Reference
Women aged 40 to 94	Serum	PFHpA: 0.084 ng/mL (mean); 1.16 ng/mL (max) PFOA: 3.0 ng/mL (mean); 27.6 ng/mL (max) PFNA: 1.07 ng/mL (mean); 10.4 ng/mL (max) PFDA: 0.28 ng/mL (mean); 3.9 ng/mL (max) PFUnDA: 0.17 ng/mL (mean); 1.3 ng/mL (max) PFHxS: 2.2 ng/mL (mean); 21.8 ng/mL (max) PFOS: 8.5 ng/mL (mean); 99.8 ng/mL (max)	Hurley et al. (2018)
337 California adults	Serum	Me-PFOSA-AcOH: 0.12 ng/mL (geometric mean) PFDeA: 0.19 ng/mL (geometric mean) PFHxS: 1.03 ng/mL (geometric mean) PFNA: 0.79 ng/mL (geometric mean)	CECBP (2018)

Population	Sample Type	PFAS Type: Concentration	Reference
		PFOA: 1.49 ng/mL (geometric mean) PFOS: 5.21 ng/mL (geometric mean)	
Children aged 3 to 11	Serum	PFHxS: 0.84 ng/mL (geometric mean) PFNA: 0.79 ng/mL (geometric mean) PFOA: 1.92 ng/mL (geometric mean) PFOS: 3.88 ng/mL (geometric mean) MeFOSAA: 0.110 ng/mL (50 th percentile) PFDA: 0.100 ng/mL (50 th percentile, girls ages 3-5 years; boys <LOD)	CDC (2018)
Adults	Blood (near contaminated site)	PFOA: 3.09 ng/mL (mean); 32 ng/mL (max) PFOS: 8.59 ng/mL (mean); 95.6 ng/mL (max) PFHxS: 4.12 ng/mL (mean); 116 ng/mL (max)	NHDPH (2016)
Adult men	Serum	PFOA: 4.50 ng/mL (mean) PFOS: 20.80 ng/mL (mean) PFHxS: 2.88 ng/mL (mean) PFNA: 1.52 ng/mL (mean)	He et al. (2018)
Adult women	Serum	PFOA: 3.46 ng/mL (mean) PFOS: 14.51 ng/mL (mean) PFHxS: 1.94 ng/mL (mean) PFNA: 1.30 ng/mL (mean)	He et al. (2018)
Pregnant women	Plasma	PFHxS: 0.37 ng/mL (mean) PFHxA: 0.16 ng/mL (mean) PFHpA: 0.18 ng/mL (mean) PFOS: 5.74 ng/mL (mean) PFOA: 2.64 ng/mL (mean) PFNA: 1.35 ng/mL (mean) PFDA: 0.57 ng/mL (mean) PFUnDA: 1.50 ng/mL (mean) PFDODA: 0.21 ng/mL (mean) PFTrDA: 0.35 ng/mL (mean) PFTeDA: 0.12 ng/mL (mean)	Tsai et al. (2018)
Pregnant women	Cord serum	PFOA: 1.59 ng/mL (mean) PFNA: 0.13 ng/L (mean) PFOS: 1.43 ng/L (mean) Sum of 11 PFASs: 3.67 ng/L mean	Cao et al. (2018)
Newborn	Umbilical cord plasma	PFOA: 7.65 ng/L (mean); 29.97 ng/L (max) PFOS: 2.93 ng/L (mean); 65.61 ng/L (max) PFNA: 0.69 ng/L (mean); 3.29 ng/L (max)	Liu et al. (2018a)

Population	Sample Type	PFAS Type: Concentration	Reference
		PFDA: 0.44 ng/L (mean); 5.73 ng/L (max) PFUA: 0.45 ng/L (mean); 5.27 ng/L (max) PFDoA: 0.10 ng/L (mean); 1.14 ng/L (max) PFHxS: 0.18 ng/L (mean); 0.85 ng/L (max) PFBS: 0.05 ng/L (mean); 0.46 ng/L (max)	
Breastfeeding women	Breast milk	PFPeA: 0.053 ng/mL (geometric mean) PFHxA: 0.047 ng/mL (geometric mean) PFHpA: 0.030 ng/mL (geometric mean) PFOA: 0.071 ng/mL (geometric mean) PFOS: 0.049 ng/mL (geometric mean)	Kang et al. (2016)

Evidence that the Candidate Chemical exhibits certain hazard traits (persistence, bioaccumulation, lactational/transplacental transfer)

PFAAs are highly persistent and tend to accumulate in the environment and biota, due to the extremely strong bond between the carbon and fluorine atoms that does not degrade under typical environmental conditions (Giesy et al. 2010; Liu and Mejia Avendaño 2013).

PFAAs and nonpolymeric PFAA precursors undergo transplacental transfer in (Midasch et al. 2007; Zhang et al. 2013; Yang et al. 2016; Zhao et al. 2017; Eryasa et al. 2019). Fluorinated ethers, such as Cl-PFESA, can also transfer from mother to fetus via the placenta (Chen et al. 2017a; Pan et al. 2017). The placenta-to-maternal serum ratios of PFOS, PFOA, and PFNA were observed to increase during gestation – more so in pregnancies with male fetuses compared to female ones – suggesting bioaccumulation in the placenta and increasing exposure with fetal age (Mamsen et al. 2019). Through transplacental transfer, the developing fetus is exposed to mixtures of PFAAs *in utero* (Bach et al. 2015; Mamsen et al. 2017). Gestational diabetes appears to significantly increase the rate of PFAS transplacental transfer (Eryasa et al. 2019).

PFAAs, including the shorter chains, are found in breast milk (Nyberg et al. 2018), potentially adding to significant cumulative early-life exposures. A 2016 study reported detectable concentrations of PFHxA in over 70 percent of breast milk samples (n=264), and detectable PFOA levels in the breast milk of nearly 99 percent of lactating mothers, associated with exposure to PFASs in consumer products (Kang et al. 2016). A study of lactating mothers in Korea found that PFAS concentrations in breast milk increased significantly after the first month of breastfeeding, likely due to a change in diet (Lee et al. 2018). Breastfed newborns can have higher body burdens of PFAAs compared to those fed only infant formula (Fromme et al. 2010). Kim et al. (2014b) analyzed concentrations of PFASs in the serum and urine of children and found a correlation between longer duration of breastfeeding and serum concentrations.

Transplacental transfer to the fetus and breastfeeding appear to be significant PFAA elimination routes for women. Serum concentrations of PFOA and PFOS are estimated to decrease by 2 to 3 percent per month while breastfeeding (Mondal et al. 2014). Biomonitoring studies of infants and small children suggest that peak concentrations of PFOS and PFOA occur before children turn 2 years old (Winkens et al. 2017b).

6. POTENTIAL FOR SIGNIFICANT OR WIDESPREAD ADVERSE IMPACTS

Reference: CAL. CODE REGS. tit. 22, § 69503.2(a).

This section integrates the information provided in the profile to demonstrate how the key prioritization principles, as identified in the SCP regulations, are met.

PFASs associated with treatments for converted textiles or leathers are released into air, water, and soil during chemical and product manufacturing, product use, and disposal. These extremely persistent chemicals are now global pollutants that can be found in all environmental compartments.

Atmospheric and oceanic currents, as well as international trade, have distributed PFASs to areas of the world far from manufacturing sites, including pristine Arctic lands and lakes (Stock et al. 2004; Muir et al. 2019).

Once PFASs enter the environment, there is no practical way to remove them. Their widespread presence has led to contaminated drinking water and human food chains, resulting in a growing public health concern (APHA 2016). Conventional, aeration, oxidation, and biological treatment are ineffective for removing PFASs from drinking water (Crone et al. 2019). Nanofiltration was the only treatment that efficiently removed PFASs from contaminated drinking water in a recent study (Boiteux et al. 2017). Anion exchange resins and high-pressure membranes such as nanofiltration and reverse osmosis can remove over 90 percent of PFAAs from drinking water, including shorter-chain ones. However, these methods are expensive, perform poorly at removing neutral PFAA precursors, and generate large waste streams with concentrated PFAAs that need to be disposed of (Crone et al. 2019).

Exposing humans and aquatic, avian, or terrestrial animal or plant organisms to PFASs associated with the treatments for converted textiles or leathers has the potential to contribute to or cause significant and widespread adverse impacts according to multiple factors considered by the Safer Consumer Products regulations.

Given (1) the sheer number of PFASs being produced or explored for use worldwide, (2) the potential for irreversible human and ecological exposures to PFASs throughout the life cycle of treatments for converted textiles or leathers, and (3) the potential for these exposures to contribute to or cause significant or widespread adverse impacts, DTSC proposes to list treatments containing PFASs for use on converted textiles or leathers as a Priority Product.

Exposing humans and aquatic, avian, or terrestrial animal or plant organisms to PFASs associated with the treatments for converted textiles or leathers has the potential to contribute to or cause significant and widespread adverse impacts according to multiple factors considered by the Safer Consumer Products regulations.

Adverse impacts linked to the Candidate Chemical's hazard traits

Reference: CAL. CODE REGS. tit. 22, § 69503.3(a).

The SCP regulations direct the Department to evaluate the potential for the Candidate Chemical to contribute to or cause adverse impacts by considering several adverse impact factors for which information is reasonably available.

PFASs associated with treatments for converted textiles or leathers show several exposure potential hazard traits:

- PFASs are either extremely persistent in the environment, or degrade into other extremely persistent PFASs;
- Many PFASs display significant mobility in environmental media, which makes them widespread in the environment and in living organisms;
- Several members of the PFAS class bioaccumulate significantly in animals or plants, including in foods consumed by humans;
- The PFASs evaluated for this trait display significant lactational or transplacental transfer, which appears to be a widespread issue given the ubiquitous presence of PFASs in breast milk and cord blood; and
- Some PFASs have high global warming potential or may contribute to global warming by increasing cloud reflectance.

These exposure potential hazard traits stem from the PFASs' unique physicochemical properties and environmental fate (see Chapter 4. [Candidate Chemical Definition and Properties](#)). Persistence has been called “the most important single criterion affecting chemical exposure and risk via the environment,” because persistent chemicals may travel longer distances and affect a larger population, have increased potential for bioaccumulation across food webs, and are very slow to remove from the environment through natural processes or engineered remediation (Mackay et al. 2014).

Once PFASs enter the environment, there is no practical way to remove them. Their widespread presence in environmental media has led to contaminated food chains and drinking water (Lindstrom et al. 2011b). PFAAs are extremely persistent, with no known natural degradation pathway under environmental conditions; most other PFASs degrade to PFAAs in the environment and within living organisms, i.e., they are PFAA precursors (Buck et al. 2011; D’eon and Mabury 2010; Krafft and Riess 2015a; Wang et al. 2014a; Washington et al. 2015). In particular, shorter-chain PFASs are highly mobile in soil and water, accumulate in edible plant parts, and are difficult and expensive to remove during water treatment (Brendel et al. 2018; Boone et al. 2019; Ateia et al. 2019b; Crone et al. 2019). The extreme environmental persistence of PFAAs, regardless of chain length, leads to their continuous presence in the environment, hence to continuous chronic background exposures for humans and

biota (Brendel et al. 2018). PFAS accumulation in sediments (Xiao 2017) and drinking water sources (Boone et al. 2019; Cousins et al. 2016), may pose significant environmental and human health hazards (Xiao 2017), and increases the potential for significant and widespread adverse impacts.

Long-chain PFAAs, which may be found in imported products, show evidence for the following toxicological hazard traits:

- Carcinogenicity
- Developmental toxicity
- Reproductive toxicity
- Cardiovascular toxicity
- Endocrine toxicity
- Hematotoxicity
- Hepatotoxicity and digestive system toxicity
- Immunotoxicity
- Nephrotoxicity and other urinary system toxicity
- Neurodevelopmental toxicity
- Respiratory toxicity.

Short-chain PFAAs, which are associated with the current PFAS formulations used in domestic treatments for converted textiles or leathers, show evidence for the following hazard traits:

- Developmental toxicity (PFBA, PFBS)
- Endocrine toxicity (PFHxA, PFBA, PFBS)
- Hematotoxicity (PFBS)
- Immunotoxicity (PFBS)
- Neurodevelopmental toxicity (PFBS)
- Reproductive toxicity (PFBA, PFBS)
- Respiratory toxicity (PFHxA)
- Ocular toxicity (PFBS)

Some of the supporting studies merely indicate the potential for shorter-chain PFASs to modulate biological responses, which we recognize is not a robust indicator of toxicological hazard. Nevertheless, increasing reports of toxicity in humans and experimental animals, sometimes at or below current exposure levels, is leading to growing public health concern (APHA 2016) and expanded regulatory interest (Corsini et al. 2014; Zushi et al. 2012). PFASs may also have hazardous degradation products; PFAA precursors degrade to PFAAs, whereas fluorinated polymers form PFCAs, hydrogen fluoride, ozone depleting compounds, and other hazardous substances if undergoing combustion at product end of life (Henry et al. 2018; Huber et al. 2009). Studies have also suggested that PFAAs, including

PFHxA, may contribute to mixture toxicity and enhance the adverse impacts associated with other hazardous compounds (see Cumulative Effects below).

The relationship between exposure and toxicological outcomes, particularly for PFASs other than PFOA and PFOS, is a major data gap (DeWitt 2015). It remains unclear whether serum concentration or other exposure metrics are useful for estimating exposures and potential adverse impacts. Also missing are early-life longitudinal exposure studies measuring internal and external exposure for multiple pathways and PFASs (Winkens et al. 2017b)

Several PFASs associated with the treatments for converted textiles or leathers show the following significant environmental hazard traits:

- Phytotoxicity, especially in green algae, which are the foundation of the aquatic food chain (Casal et al. 2017; Ding et al. 2012a); and
- Impairment of wildlife development, reproduction, or survival.

Discharges of PFAS-contaminated WWTP effluents into streams and rivers have led to contamination of water and sediments at concentrations considered potentially hazardous to wildlife and threatening to biodiversity (Rodriguez-Jorquera et al. 2016). PFAAs bioaccumulate in phyto- and zooplankton and are toxic to green algae, which are the foundation of the aquatic food chain (Casal et al. 2017; Ding et al. 2012b). In general, aquatic invertebrates, which comprise the largest percentage of animal biomass in aquatic ecosystems, are particularly susceptible to adverse impacts because they are often exposed to complex mixtures of waterborne contaminants (Bradley et al. 2017). Water-filtering species of aquatic animals are the most highly exposed to PFAAs (Renzi et al. 2013). The ubiquitous presence of PFASs in aquatic environments and the limited toxicological knowledge about cumulative impacts to aquatic species have led to increasing concern about the potential for irreversible adverse aquatic impacts (Scheringer et al. 2014).

Laboratory studies of environmentally relevant species of algae, aquatic plants, terrestrial plants, fish, amphibians, mammals, birds, and important pollinating insects, as well as studies of birds exposed in the wild, show an association between PFAS exposure and several adverse environmental impacts, including developmental, reproductive, or survival impairment and death (see Appendix 2) (Annunziato et al. 2019; Pedersen et al. 2016; Liu et al. 2015b; Sanchez et al. 2015; Soloff et al. 2017).

Cumulative effects

Reference: CAL. CODE REGS. tit. 22, § 69503.3(a)(1)(C).

Cumulative effects occur from exposures to the Candidate Chemical and other chemicals with similar hazard traits or endpoints.

Individual PFASs rarely occur in isolation. People and biota are exposed to mixtures of PFASs and other chemicals, potentially leading to cumulative adverse impacts, yet toxicological endpoints are typically assessed using single PFASs. The adverse human and environmental health effects resulting from exposures to PFAS mixtures have not been well-characterized, and toxicological reference values for PFAS mixtures are lacking. Some have argued that exposure to PFAS mixtures may exert greater toxicity than exposure to a single PFAS (Wang et al. 2011). Although exposure to multiple PFASs may result in additive effects, only a few studies have investigated the toxicity of PFAS mixtures, with varying results. For instance, a cumulative health risk assessment of 17 PFASs in the Swedish population concluded there was a threat of hepatotoxicity or reproductive toxicity in a subpopulation that routinely consumed PFOS-contaminated fish, but not in the general public (Borg et al. 2013).

Evidence of increased mixture toxicity includes results from an *in vitro* study in which binary combinations of PFAAs at low concentrations behaved additively in activating PPAR-alpha (Wolf et al. 2014). A cytotoxicity assay found that a mixture of PFASs altered the cellular lipid pattern of human placental cells at levels below those that induce toxic effects (Gorrochategui et al. 2014). An *in vitro* study of PFOA, PFOS, and PFNA found that toxicity in a human macrophage cell line and acute toxicity in zebrafish were greater for mixtures than individual compounds (Rainieri et al. 2017). The cytotoxicity of individual PFASs and of binary mixtures using an amphibian fibroblast cell line showed mixtures were approximately additive, except with PFOS and PFOA, which were weakly synergistic (Hoover et al. 2019).

Mixture toxicity studies of PFASs and other toxicants are limited. The results include:

- Effects on gene expression patterns in zebrafish embryos coexposed to a mixture of PFHxA, PFOS, and PCB126; lowered oxidative stress response from exposure to PCB126 and PFHxA, and to PCB126, PFOS, and PFHxA (but not from the individual compounds, nor from PCB126 and PFOS), suggesting PFHxA plays a synergistic role, although the exact mechanism remains unknown (Blanc et al. 2017);
- Endocrine disruption in developing rats coexposed to a mixture of PFHxS and 12 endocrine-disrupting chemicals (including bisphenol A, dibutyl phthalate, di-2-ethylhexyl phthalate, and butyl paraben), at doses relevant for human exposure, including doses at which PFHxS alone and the mixture of endocrine disruptors alone showed no or only weak effects (Ramhøj et al. 2018);
- Clinically relevant perturbations of normal cell function of benign breast epithelial cells coexposed to a mixture of PFOA, bisphenol A, and methylparaben at concentrations relevant to human exposure; these effects were not observed when cells from the same lines were exposed to the chemicals individually (Dairkee et al. 2018);
- Cumulative adverse effects *in vitro* of PFOS, PFOA, and several other contaminants (Hg²⁺, Cd²⁺, 2,4-D, propylparaben, mitomycin C, and furazolidone) in a bioluminescent cyanobacterial

toxicity test (Rodea-Palomares et al. 2012);

- Increased severity of oxidative stress and apoptosis to zebrafish embryos following exposure to PFOS and zinc oxide nanoparticles (Du et al. 2017); and
- Adverse impacts on mitochondrial function in juvenile Chinook salmon after exposure to a mixture of contaminants of emerging concern, including several PFASs, pharmaceuticals, and personal care products, at environmentally relevant concentrations (Yeh et al. 2017).

While the above data suggest a potential for cumulative harm from exposure to mixtures of PFASs or of PFASs and other chemicals, it is unclear whether the outcomes from the cell lines, zebrafish, or juvenile salmon are associated with adverse phenotypes. In spite of the limitations of these approaches, as described by the National Academy of Sciences (NAS 2017), these studies suggest that exposure to certain combinations of PFASs, or co-exposure to certain PFASs and other chemicals, has the potential to contribute to disease by perturbing various pathways or activating some mechanism. In particular, PFAAs may have cumulative impacts with one another and with other hazardous chemicals, but co-exposures that dramatically affect adverse impacts are not known at this time and require more scientific research.

Adverse waste and end-of-life effects

Reference: CAL. CODE REGS. tit. 22, § 69503.2(b)(1)(B) and 69501.1 (a)(8).

This section summarizes findings related to the waste materials and byproducts generated during the life cycle of the product and their associated adverse effects, as described in the SCP regulations. These considerations can form part of the basis for proposing the product-chemical combination.

Effects on solid waste and wastewater disposal, treatment, and recycling

Conventional wastewater treatment systems such as sedimentation, flocculation, coagulation, and biological degradation are ineffective for PFAS removal (Rahman et al. 2014; Ng et al. 2019). Adsorption and ion-exchange have also been recently explored as methods that could increase the speed and efficiency of PFAS removal (Ji et al. 2018; Ateia et al. 2019a). While adsorption and separation technologies combined with destructive technologies show promise for removing a wider range of PFASs, their large-scale application poses several challenges (Horst et al. 2018; Li et al. 2020). Shorter-chain PFAAs are especially challenging to remove from aquatic environments and to destroy (Li et al. 2020).

Biosolid treatments are also ineffective at reducing the levels of PFAAs. A study of commercial biosolids from the U.S. and Canada found that thermal hydrolysis had no apparent effect on the PFAA concentration, whereas heat treatment and composting increased PFAA concentrations (especially

PFHxA) via the degradation of precursors; only blending with PFAS-free material decreased the concentration of PFAAs in the commercial biosolids, by diluting it (Kim Lazcano et al. 2019).

Removal of PFASs from the waste stream is expensive. For instance, following contamination from carpet manufacturing in Dalton, Georgia water utility Gadsden Water stated that it “has suffered substantial economic and consequential damage, including expenses associated with the future installation and operation of a filtration system capable of removing the chemicals from the water and lost profits and sales” (Chapman 2016). Wastewater effluent can contribute significantly to the PFAA contamination of U.S. drinking water sources (Guelfo and Adamson 2018).

Discharges or disposal to storm drains or sewers that adversely impact the operation of wastewater or stormwater treatment facilities

PFASs are not routinely removed by WWTPs, and removal is expensive (Arvaniti and Stasinakis 2015; Mudumbi et al. 2017). In fact, wastewater treatment typically transforms PFAA precursors into PFAAs, resulting in higher PFAA concentrations in WWTP effluent compared to the influent (Ahrens 2011; Frömel et al. 2016; Guerra et al. 2014).

Releases of the Candidate Chemical during product end-of-life

In California, textiles are one of the top 10 components of the landfill waste stream, accounting for approximately 4 percent of waste by volume, with carpets accounting for approximately 3.2 percent. In 2016, 343 million pounds of carpet were discarded in California, 75 percent (257 million pounds) of which were disposed to landfills.

In California, textiles are one of the top 10 components of the landfill waste stream, accounting for approximately 4 percent of waste by volume, with carpets accounting for approximately 3.2 percent (CalRecycle 2015, 2017; GAIA and Changing Markets 2017). Leather products are grouped within a broad category called “composite organic” waste, and their proportion in the landfill waste stream cannot be easily quantified (CalRecycle 2015). From 2015 to 2016, due to the 2010 California Carpet Stewardship Law, postconsumer carpet collection and carpet tile recycling increased by 4 and 31 percent, respectively. In 2016, 343 million pounds of carpet were discarded in California, 75 percent (257 million pounds) of which were disposed to landfills (CCSP 2016).

The use of incinerators has become popular as a method to reduce landfill volumes while increasing energy recovery (CCSP 2016). Incinerating textile or leather products (i.e., carpets, furniture, upholstery) treated with fluorinated polymers, including shorter-chain-based side-chain fluorinated polymers, can result in air emissions. Potential air toxicants include PFCAs, ozone depleting substances such as CFCs, and potent greenhouse gases such as fluorocarbons, since the mineralization

(defluorination) of PFASs is incomplete even at high temperatures and in the presence of calcium hydroxide (Huber et al. 2009). This combustion occurs in incinerators often located in low-income communities, where emissions burden people already impacted by aggregate chemical exposures from other sources, creating environmental justice concerns (GAIA and Changing Markets 2017).

Empty containers of treatments for converted textiles or leathers may contain PFAS residue, which may enter the environment when the containers are landfilled. Municipal landfill leachate is a known point source of PFAS emissions to the environment (Benskin et al. 2012; Busch et al. 2010; Eggen et al. 2010; Fuertes et al. 2017; Hamid et al. 2018; Huset et al. 2011; Lang et al. 2017; Boucher et al. 2019), with treated carpets, rugs, furnishings, and clothing as significant contributors (Gallen et al. 2017; Lang et al. 2016; Shoaieioskouei 2012). This impacts unlined landfills and those with a leachate management system, but also those that collect and treat the leachate in WWTPs, which often fail to adequately remove PFASs (Arvaniti and Stasinakis 2015; Huset 2007) and discharge PFAS mixtures into receiving waters (Dauchy et al. 2017; Hamid and Li 2016; Pan et al. 2016). WWTPs are also meaningful sources of volatile PFASs, as are landfilled carpets and rugs (Ahrens et al. 2011b; Weinberg et al. 2011). A study of PFASs in digested sewage sludge from 45 WWTPs in Switzerland found that the carpet protection and textile finishing industries were the most relevant point sources of PFCAs (Alder and van der Voet 2015).

PFASs in landfill leachates, land-applied wastewater, and biosolids can contaminate streams, rivers, and other drinking water sources (Hoffman et al. 2011; Lindstrom et al. 2011a). Stormwater runoff containing PFASs released from consumer products may be a significant pathway for PFAS transport and release to surface waters (Houtz and Sedlak 2012).

Populations that may be adversely impacted

Reference: CAL. CODE REGS. tit. 22, § 69503.3(a)(1)(F) and 69503.3(a)(2).

This section identifies specific populations of humans and environmental organisms that may be harmed if exposed to the Candidate Chemical in the product. Sensitive subpopulations, environmentally sensitive habitats, endangered and threatened species, and impaired environments in California have special consideration as they may be more vulnerable.

Exposures to PFASs occur frequently and from multiple sources over a lifetime, including from placental transfer, breastfeeding, drinking water, food, and contact with consumer products and other environmental media. PFASs have the potential to cause significant and widespread adverse impacts to:

- Sensitive subpopulations (which include fetuses, infants, children, pregnant women, some workers, and people with certain preexisting conditions);

- Environmentally sensitive habitats; and
- Threatened and endangered species.

Fetuses, infants, toddlers, and young children experience higher relative exposure levels and are more vulnerable to the effects of toxicants. Workers, including carpet and upholstery cleaners; workers in upholstered furniture, furnishings, clothing, shoe, and carpet stores; auto dealership workers; and auto detailing technicians can experience high PFAS exposures on the job. Individuals with certain preexisting conditions (e.g., elevated cholesterol, high blood pressure, poor kidney function) may be especially sensitive to PFASs. Pregnant women's exposure to PFASs is of concern due to transplacental and lactational transfer to the offspring. Endangered and threatened species in California are vulnerable to contamination of waterways and food webs with PFASs from stain- and soil-repellent treatments for converted textiles or leathers.

The increasing concentrations of PFASs associated with stain and soil repellents found in environmental media, animal tissues, and biota can contribute to or cause significant or widespread adverse impacts in people and the environment.

Due to their widespread presence in the environment and biota, PFASs may contribute to or cause adverse impacts in all humans and other organisms. A recent review paper (Krafft and Riess 2015b) states that:

“Because of ubiquitous PFAS presence, there is no unexposed control population and many studies compare outcome data for high-dose sub-groups with a low-dose sub-group within the same population, a practice that can weaken dose-response correlations.”

The potential for adverse impacts of PFASs on aquatic species has been well-documented, with toxic effects noted in aquatic plants (Ding and Peijnenburg 2013; Ding et al. 2012a; Hoke et al. 2012) and widespread bioaccumulation in aquatic animals. Less is known about potential impacts to terrestrial animals and plants, especially at the population level. Shorter-chain PFASs generally show increased uptake by leafy plants compared to the longer chains (Blaine et al. 2014a), though longer chains may preferentially accumulate in roots (Zhao et al. 2017). PFASs have been shown to bioaccumulate in earthworms, indicating the potential for biomagnification in terrestrial species (Karnjanapiboonwong et al. 2018). Wildlife, particularly apex predators, may suffer adverse effects from chronic exposure to PFASs in food and water (Tartu et al. 2017; Tipton et al. 2017). Fish-eating birds, including eagles and ospreys, are especially vulnerable to PFAS exposure via food chain contamination (Giesy et al. 2010). Known impacts to birds from PFAS exposure include reduced embryo survival in chicken (Nordén et al. 2016).

Infants, toddlers, and small children comprise a sensitive subpopulation because of their increased ingestion and inhalation rates per unit of body weight, rapid development, immature physiological ability to detoxify environmental contaminants, and behavioral characteristics that predispose them to increased exposures to environmental contaminants (U.S. EPA 2011). This results in a higher body burden of PFASs as compared to adults (Rappazzo et al. 2017). Pregnant women and fetuses are also sensitive subpopulations because of transplacental migration and the vulnerability of the rapidly developing fetus (Slotkin et al. 2008). Breastfed infants are susceptible to increased exposures to PFASs in breast milk, because breastfeeding is a route of PFAS excretion for lactating women (Kang et al. 2016; Kärrman et al. 2007; Mogensen et al. 2015; Mondal et al. 2014). Infants, toddlers, and small children often have increased exposures due to hand-mouth behaviors that can lead to increased incidental ingestion of dust and soil with environmental contaminants, and higher doses relative to body weight compared to adults (U.S. EPA 2011). Infants who crawl over carpets and upholstery are at increased risk of inhaling dust (Wu et al. 2018), which may contain PFASs. A study involving over 1,000 children up to seven years old found a negative correlation between plasma concentrations of most PFAAs and age, indicating higher exposures in younger children (Zhang et al. 2018). Table 7 lists the main routes of PFAS exposure for children.

With more than 3.9 million births in the U.S. in 2016, fetuses and infants constitute a large subpopulation highly vulnerable to exposures to the PFASs released during the treatment of converted textiles or leathers. Because PFASs undergo transplacental migration (Midasch et al. 2007; Zhang et al. 2013; Yang et al. 2016; Chen et al. 2017a; Zhao et al. 2017; Eryasa et al. 2019), fetuses are continuously exposed from conception until birth. PFASs are also frequently detected in breast milk (Nyberg et al. 2018), with nursing infants exposed daily over many months. The CDC reported that 79 percent of infants born in 2011 were breastfed, with 27 percent still breastfeeding at 12 months of age. The amount of breast milk ingested by newborns averages 76 g, six to eighteen times per day (CDC 2014).

Toddlers can experience increased exposures to PFASs due to behavioral characteristics. Children between the ages of 1 and 3 spend a lot of time in contact with the floor, where dust settles, and engage in indoor hand-mouth behaviors an average of 16 times per day (U.S. EPA 2011). Thus, incidental dust ingestion can be a significant route of exposure for young children (Mercier et al. 2011; Tian et al. 2016). While adults are estimated to ingest an average of 50 mg/day of soil and dust, children under 6 may ingest 100 mg/day (U.S. EPA 2011). PFAS doses via incidental dust ingestion could be approximately an order of magnitude higher for toddlers than for adults (Karásková et al. 2016). A study of toddlers' cumulative exposure to PFOA and PFOS estimated a daily intake of 53.6 ng/day of PFOA and 14.8 ng/day of PFOS (Tian et al. 2016).

Table 7: Prenatal and early-childhood PFAS exposure routes²¹

Exposure Pathway	Basis
Transplacental migration	Exposure to PFASs begins before birth. Transplacental passage is a significant route of human exposure to PFAAs (Kim et al. 2011) and their precursors (Yang et al. 2016). Transfer to the fetus is one of the major PFAS elimination routes for women, particularly for PFOA (Lee et al. 2013).
Ingestion – breast milk	Breast milk ingestion is a significant route of PFAS exposure for infants (Mondal et al. 2014). Infants fed breast milk may have higher PFAS dietary exposure than those fed infant formula (Fromme et al. 2010).
Ingestion – food	Food ingestion is considered a primary route of exposure for the general population, with higher relative exposures in infants, toddlers, and small children because of their low body weight (Egeghy and Lorber 2011). Due to widespread contamination, PFASs are found in a variety of animal and plant foods (Blaine et al. 2014a; Pérez et al. 2014). PFASs are readily absorbed after ingestion (ATSDR 2015, 2018a; Danish Environmental Protection Agency 2015a).
Ingestion – drinking water	Numerous studies conclude that drinking water is a major source of PFAS intake (DeWitt 2015; Trudel et al. 2008). In 2016, U.S. EPA issued drinking water health advisories for PFOA and PFOS of 70 ng/L (combined concentrations), to protect the most sensitive populations – fetuses during pregnancy and breastfed infants (U.S. EPA 2016b). Some states have adopted, or are considering adopting, lower limits.
Inhalation and ingestion of dust	PFASs are widely found in house dust, with higher concentrations in homes with treated carpets (Haug et al. 2011). Exposure via dust is higher in toddlers and small children than adults and can be significant due to children’s lower body weights, increased inhalation rate, higher incidental dust ingestion rates, increased floor contact and hand-to-mouth behavior (Mercier et al. 2011; Tian et al. 2016). PFASs are readily absorbed following inhalation and ingestion (ATSDR 2018a).

²¹ Protecting children, a sensitive subpopulation, is one of the goals identified in SCP’s 2018-2020 Priority Product Work Plan. Childhood is a life stage that all members of a population experience, although children constitute a population subgroup when evaluating exposures at a specific time.

Office workers are another large group of Californians possibly experiencing chronic exposures to PFASs while on the job. Office air can be contaminated with FTOHs from textile or leather treatments, leading to involuntary, chronic inhalation exposures in workers, which has not been well-characterized (Bohlin-Nizzetto et al. 2015; Schlummer et al. 2015). PFASs can also adsorb to office dust or be released from surface abrasion as tiny particles, which office workers may ingest (Fraser et al. 2013). Due to their long usage life, commercial carpets could be a near-daily source of exposure to PFOA, PFOS, and other phased-out longer-chain PFASs for office workers.

Workers in the textile or leather industry, including retail workers and recyclers, also comprise a sensitive subpopulation because of their potentially higher exposure, particularly to volatile and semi-volatile PFASs emitted from these products. The highest levels of volatile PFASs, such as FTOHs, have been found in stores selling outdoor clothing, carpets, or furniture (Langer et al. 2010; Schlummer et al. 2013). One study estimated that for people living or working in environments with high levels of FTOHs in indoor air, such as carpet stores, exposure to PFOA resulting from FTOH degradation is in the same order of magnitude as dietary PFOA exposure (Schlummer et al. 2013).

Studies demonstrate substantial human exposure in workers at fluorochemical manufacturing facilities (Gao et al. 2018) and communities relying on contaminated drinking water sources (Heydebreck et al. 2016). People living near PFAS chemical manufacturers, or in other areas contaminated by PFASs, have higher-than-average levels of PFASs (Fromme et al. 2009). Individuals with biomarkers of susceptibility for certain health conditions may be unusually sensitive to PFAS exposure (ATSDR 2018a).

Residents of low-income communities may be disproportionately impacted by chronic exposures to multiple hazardous industrial chemicals, increasing the likelihood of cumulative adverse health effects (U.S. EPA 2016a). Manufacturing facilities, including those that release PFASs, are often located in these communities. Environmental justice concerns also arise regarding contamination in areas far from manufacturing sites, such as the remote regions of the Arctic. For instance, the traditional diet of the Inuit in Nunavut, Canada, the Nuuk Inuit in Greenland, and the Faroese includes foods heavily contaminated with PFASs that originated from thousands of miles away (Grandjean et al. 2012, 2016; Long et al. 2012; Ostertag et al. 2009).

California's endangered and threatened species could be adversely affected by exposure to PFASs associated with treatments for converted textiles or leathers, especially considering the adverse effects on reproduction and development demonstrated for some PFAAs. This could contribute to the current biodiversity crisis in aquatic ecosystems (Abell 2002; Mora and Sale 2011; Valentini et al. 2016). Threatened and endangered species of fish and marine mammals are particularly vulnerable to population-level adverse impacts if they cannot obtain food or avoid predators. Mammals and birds occupying the highest trophic level, including orcas, wolves, grizzly bears, eagles and condors, are

vulnerable to adverse health effects from ingestion of food and water contaminated with PFASs that bioaccumulate and biomagnify (Kannan et al. 2006; Kelly et al. 2009).

Environmentally sensitive habitats in California, including estuaries and other wetlands, can receive surface water contaminated with PFASs via wastewater plant effluent or surface runoff, leading to contaminated water and sediments. These ecosystems are important breeding, spawning, and nesting sites, and feeding grounds for millions of migratory birds – including threatened and endangered species – that transit California during their annual migration.

7. OTHER REGULATORY PROGRAMS

Reference: CAL. CODE REGS. tit. 22, § 69503.2(b)(2).

DTSC has assessed all applicable state and federal laws and regulations and international treaties or agreements with the force of domestic law related to the product or the Candidate Chemical in the product. DTSC has determined that these programs do not overlap or conflict with this proposal to list treatments containing members of the class of PFASs for use on converted textiles or leathers, nor with any subsequent regulation that may result from such listing.

No California state laws or regulations currently address the use of PFASs in treatments for converted textiles or leathers. Also, there are no applicable international treaties or agreements with the force of domestic law.

At the federal level, Section 5 of TSCA authorizes U.S. EPA to issue Significant New Use Rules (SNURs) for new chemicals or existing chemicals used in a significant new way. A SNUR requires companies to notify U.S. EPA at least 90 days prior to manufacturing, importing, or processing substances for a significant new use, and to submit a notification including information about the chemical's identity, physical characteristics, processing and use, and available toxicity data. U.S. EPA has 90 days to evaluate the new use and can request more data, prohibit or limit the manufacture, or allow the use. The following SNURs relate to PFASs:

- December 9, 2002 – SNUR regarding any future manufacture (including imports) of 75 PFASs specifically included in the 2000-2002 voluntary phaseout of PFOS by 3M (U.S. EPA 2002a);
- March 11, 2002 – SNUR regarding any future manufacture (including imports) of 13 PFASs specifically included in the 2000-2002 voluntary phaseout of PFOS by 3M (U.S. EPA 2002b);
- October 9, 2007 – SNUR for 183 PFASs believed to no longer be manufactured, imported, or used in the United States (U.S. EPA 2007);
- September 30, 2013 – SNUR requiring companies to report their intent to manufacture certain PFOA-related chemicals to treat carpets, as well as their intent to import carpets containing these PFASs (U.S. EPA 2013); and
- (Proposed) January 21, 2015 – proposed SNUR affecting manufacturers (including importers) of PFOA and PFOA-related chemicals, including as part of articles, and processors of these chemicals (U.S. EPA 2015).

In 2006, U.S. EPA developed a 2010/2015 Stewardship Program for reducing emissions of PFOA, its precursors, and related higher homologues (U.S. EPA 2010). Through a Memorandum of Understanding with U.S. EPA, eight major U.S. manufacturers agreed to voluntarily eliminate PFOA in their emissions and products by 2015. Participating companies include: Arkema, Asahi, BASF, Clariant, Daikin, 3M/Dyneon, DuPont, and Solvay Solexis.

Since 2000, U.S. EPA has reviewed hundreds of substitutes for PFOA, PFOS, and other longer-chain PFASs, particularly regarding their toxicity, fate, and bioaccumulation under the New Chemicals Program (U.S. EPA 2017c). For many PFASs, U.S. EPA has used TSCA section 5(e) Consent Orders to require testing while allowing production and use. U.S. EPA is also investigating substitutes for certain direct uses of PFOA.

On January 27, 2010, U.S. EPA amended the Polymer Exemption Rule for new chemicals under TSCA to exclude certain side-chain fluorinated polymers due to potential risk to human health or the environment (U.S. EPA 2002b).

8. POTENTIAL ALTERNATIVES

Reference: CAL. CODE REGS. tit. 22, § 69503.2(b)(3).

This section summarizes information available to DTSC regarding alternatives that may or may not be safer than the Candidate Chemical. DTSC does not need to ensure that these alternatives are safer and may summarize their associated hazards to illustrate readily available information. The sections below may include information such as how readily available an alternative is, product functions addressed by the alternative, and implications for manufacturers using the alternative (e.g., use limitations, product reformulation, different equipment needs).

Given the potential for exposures and significant or widespread adverse impacts, replacing currently used PFASs in treatments for converted textiles or leathers with other members of the PFAS class could constitute a regrettable substitution. Hence, this proposal covers treatments for converted textiles or leathers containing any member of the class of PFASs.

Given the potential for exposures and significant or widespread adverse impacts, replacing currently used PFASs in treatments for converted textiles or leathers with other members of the PFAS class could constitute a regrettable substitution. Hence, this proposal covers treatments for converted textiles or leathers containing any member of the class of PFASs.

PFAS-free chemical alternatives that prevent soils from adhering to and staining agents from penetrating the fiber surface are becoming available. Examples include TriPlex's Go Green Guard and Go Clean 2X Rinse; Vectra Enterprises, Inc.'s Vectra Spray; and Innovative Chemical Technologies, Inc.'s Flexipel™ HR-100. DTSC does not currently have complete information on the chemical content of these alternatives and has not evaluated their safety. The exact formulation of these products is largely unknown because manufacturers withhold the information as proprietary trade secrets.

Other potential PFAS alternatives have been patented but may not yet be commercially available. A patent from February 2017 describes "fluorine-free compositions rendering textile articles, such as carpets and other textile floor coverings made from synthetic fibres or natural fibres which are water repellent, soils resistant and stain resistant," based on (a) a nanoparticle silicate clay, (b) an anionic acrylic-based copolymer binding agent, and (c) water (Invista North America 2017). Another patent mentions a "superhydrophobic" agent containing nonfluorinated, water-based hydrophobic components that have useful fluid management properties, including stain repellency. It goes on to call this application a viable treatment product that incorporates titanium oxide nanoparticles for maintaining "a translucent-white superhydrophobic surface treatment that does not require subsequent pH-modification" (Qin et al. 2018).

Other potential alternatives to PFASs in treatments for converted textile or leather may include siloxane and silicone polymers, hydrocarbons (e.g., paraffin waxes), derivatives of fatty acids (e.g., long-chain fatty acid esters), polyurethanes, and dendrimers (hydrocarbon- or polyurethane-based) (Danish Environmental Protection Agency 2015b; KEMI 2015; Teli 2018; U.S. EPA 2012; Atav 2018). These are all durable water repellents without oil and alcohol repellency. Some nonfluorinated (meth)acrylates/urethanes have stain release properties for both oil- and water-based stains (Danish Environmental Protection Agency 2015b). These alternatives appear promising if they can be scaled for high throughput manufacturing.

A Danish Environmental Protection Agency review concluded that the main ingredients in paraffin-based repellents are mostly harmless, readily biodegradable, nonbioaccumulative, and nontoxic to aquatic organisms; however, some products may contain potentially hazardous ingredients or impurities such as isocyanates, dipropylene glycol, and metal salts (Danish Environmental Protection Agency 2015b). Similarly, they found that most silicones used in textile impregnation are inert, generally without adverse effects, but toxic manufacturing intermediaries such as D4 (suspected of damaging fertility) and D5 (a potential carcinogen) may be present at trace levels. For dendrimer-based repellents, they lacked sufficient information for an assessment but found that some products may contain potentially toxic siloxanes, cationic polymers, isocyanates, or powerful irritating organic acids.

9. ADDITIONAL CONSIDERATIONS

This section summarizes other relevant information not captured under the adverse impact and exposure factors named in section 69503.3 of the Safer Consumer Products regulations.

Although PFAS manufacturers have been aware of these chemicals' adverse impact potential for decades, national and international action to limit human and ecological exposures has only recently started. In 2015, more than 200 scientists from 40 countries signed The Madrid Statement on Poly- and Perfluoroalkyl Substances (PFASs) (Blum et al. 2015), a scientific consensus on the persistence and potential for harm of this class of chemicals, and a roadmap for international cooperation to prevent further harm.²²

U.S. EPA research and action plan

On December 4, 2017, U.S. EPA announced a cross-agency effort to address PFASs, including filling data gaps, supporting communities impacted by drinking water contamination, and proactively communicating with states, tribes, partners, and the public about the health effects of PFAS exposure (U.S. EPA 2017d). On March 13, 2018, U.S. EPA also announced a partnership with the National Toxicology Program to develop a tiered testing approach to characterize the toxicity and toxicokinetics of 75 PFASs that represent the entire class (U.S. EPA 2018). In February 2019, U.S. EPA revealed its PFAS Action Plan, which includes, among others: (1) setting a maximum contaminant level (MCL) for PFOA and PFOS by the end of 2019; (2) beginning regulatory work to list PFOA and PFOS as hazardous substances under the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA); (3) publishing draft toxicity assessments for PFBA, PFHxA, PFHxS, PFNA, and PFDA in 2020; and (4) continuing to improve analytical methods, monitoring, and risk communication (U.S. EPA 2019). In September 2019, U.S. EPA announced it is awarding approximately \$6 million to eight grantees to research PFASs in waste streams and identify practical approaches to manage their potential impacts (U.S. EPA 2019).

Exposure limits

Academic scientists, environmental consulting firms employed by chemical manufacturers, and government agencies around the world have conducted risk assessments using a variety of methods with the goal of determining whether the levels of PFASs found in different environmental media pose a health risk to humans or animals. Contradictory conclusions have been drawn depending on several factors: the type of model used to calculate risk, the assumptions underlying the assignment of exposure factors in exposure dose calculations, and the uncertainties in the data used in the dose-

²² One of the authors of this document, Simona Bălan, Ph.D. co-authored the Madrid Statement prior to joining DTSC.

response assessment. This has led to different regulatory levels for allowable concentrations in drinking water.

For instance, to address concerns about potential health effects from cumulative exposures to PFOA and PFOS, U.S. EPA established individual chronic reference doses for PFOA and PFOS of 0.00002 mg/kg/day (U.S. EPA 2016c, d). These are estimates of daily human exposure expected to be without adverse health effects over a lifetime. In the fall of 2016, after concerns that more than 16 million Americans are exposed to PFOA and PFOS via contaminated drinking water, U.S. EPA reviewed new studies of developmental effects and reduced its 2009 provisional health advisory of 400 parts per trillion (ppt)²³ to a lifetime drinking water health advisory of 70 ppt for PFOA and PFOS combined (U.S. EPA 2016b). In 2018, ATSDR developed MRLs (ATSDR 2018b) that correspond to a drinking water advisory levels of 11 ng/L for PFOA and 7 ng/L for PFOS (Sunderland et al. 2019).

These health advisories are recommendations only and are not legally binding or enforceable. In the fall of 2016, U.S. Representative Brendan Boyle of Pennsylvania introduced legislation (H.R. 6125) to require an enforceable nationwide standard for PFASs in drinking water to replace the U.S. EPA's voluntary health advisories (Boyle 2016). The bill is still pending in the House Committee on Energy and Commerce. On April 13, 2018, 25 U.S. senators wrote a letter to U.S. EPA Administrator Pruitt asking him to “expeditiously declare an MCL for all PFAS, based on rigorous scientific evidence” (U.S. Senate 2018).

Several U.S. states have set lower health advisories than U.S. EPA, but most of these are also not legally binding. In May 2016, the state of Vermont issued a drinking water health advisory of 20 ppt of PFOA and PFOS combined, which was also based on developmental endpoints, including reduced ossification and accelerated puberty in males and low birth weight (Vermont Department of Health 2016). In May 2017, the Minnesota Department of Health released revised drinking water guidance values of 35 ppt for PFOA and 27 ppt for PFOS (MDH 2017), to “reflect new state-level analysis of the potential for mothers to pass along the chemicals to fetuses and nursing infants.” In November 2017, New Jersey became the first state to set formal MCLs of 14 ppt for PFOA and 13 ppt for PFNA (NJDEP 2017). On December 13, 2017, Michigan Democrats introduced House Bill 5373, which would establish a legal drinking water limit of 5 ppt for PFOA and perfluorooctane (MHR 2017). On June 26, 2018, OEHHA released a recommendation for interim notification levels of 14 ppt for PFOA and 13 ppt for PFOS in drinking water based on evidence for developmental toxicity, immunotoxicity, cancer, and liver toxicity (OEHHA 2018). In August 2019, OEHHA updated its research and recommended that notification levels be set at the lowest levels at which PFOA and PFOS can be reliably detected in drinking water using currently available and appropriate technologies (OEHHA 2019). As a result, the California State Water

²³ 1 ppt is equivalent to 1 ng/L water.

Resources Control Board set its drinking water notification levels to 5.1 ppt for PFOA and 6.5 ppt for PFOS (California State Water Resources Control Board 2019).

Some scientists have argued that the limit should be even lower to protect sensitive subpopulations. For example, (Grandjean and Clapp 2015, p. 2015) derived a limit of 1 ng/L based on evidence of immunotoxicity in children, combined with uncertainties from incomplete toxicity testing and lack of epidemiological studies.

Cleanup at military sites

PFASs have also become a concern for environmental cleanup across the United States due to contamination from firefighting, manufacturing, and other activities. PFAAs and other PFASs used in AFFFs were found to contaminate surface water, groundwater, and soils, and to have numerous adverse impacts.

On August 1, 2017, a bipartisan group of eight Democrats and six Republicans in the U.S. House of Representatives signed a letter asking the White House to address PFAS contamination caused by Department of Defense firefighting activities (Kildee et al. 2017). Also in August 2017, the New York State Department of Health and state officials from Alaska, Michigan, New Hampshire, Pennsylvania, and Vermont asked CDC's Agency for Toxic Substances and Disease Registry to launch a longitudinal study of PFAS-impacted communities (Zucker et al. 2017). On December 12, 2017, the Fiscal Year 2018 National Defense Authorization Act was signed into law including a \$7 million budget for the CDC and the Agency for Toxic Substances and Disease Registry (ATSDR) to run, in consultation with the Department of Defense, a national study on the health impacts of human exposure to PFASs, especially from contaminated drinking water (U.S. Senate 2017). The final federal budget bill passed by lawmakers in March 2018 included \$10 million for the PFAS healthy study, and \$43.8 million for Department of Defense cleanup activities (U.S. House of Representatives 2018). In January 2019, a bipartisan Congressional PFAS Task Force was created to address the PFAS contamination on military bases (Office of Congressman Dan Kildee 2019).

Restrictions on PFASs in product

In March 2018, the San Francisco Department of the Environment adopted regulations regarding environmentally preferable purchasing, which include a requirement that all carpet purchased for city projects contain no intentionally-added PFASs (SFE 2018). Also in March 2018, two new bills were signed into law in Washington state: Engrossed Substitute Senate Bill 6413, which prohibits the use of PFAS-containing firefighting foam for training purposes as of July 1, 2018, and, as of July 1, 2020, prohibits the sale of firefighting foams with intentionally-added PFASs, with exemptions for the military, airports, oil refineries, and chemical plants (State of Washington 2018a) and Engrossed Substitute House Bill 2658 (State of Washington 2018b), which prohibits the sale of food packaging

containing PFASs as of January 1, 2022, provided the state Department of Ecology identifies safer alternatives by January 1, 2020 (State of Washington 2018c). On July 28, 2019, the state of Washington enacted Substitute Senate Bill 5135, which authorizes the state's departments of Ecology and Health to regulate classes of chemicals in consumer products (Washington State Legislature 2019). The law identifies PFASs as one of the five initial priority chemical classes. In September 2019, the state Department of Ecology announced that it will be focusing its initial research on PFASs in carpets and aftermarket carpet treatments (Safer Products for Washington Team 2019).

Major international companies are beginning to remove PFASs from the products they manufacture or sell. Carpet manufacturers Interface and Tandus Centiva have successfully phased out PFASs in their products without sacrificing performance (Davis 2016; Wilkinson 2016). Several apparel manufacturers, such as Levi Strauss & Co., and manufacturers of furniture and furnishings, such as IKEA and Crate and Barrel, have also phased out PFASs from their products (Blum 2016). In 2016, outdoor gear manufacturer Columbia introduced its first PFAS-free rain jacket (Jurries 2016). In January 2017, Target announced that, as part of its new chemical management policy, it will stop selling textile products with added PFASs by 2022 (Target Inc. 2017). In May 2017, 18 major purchasers including Kaiser Permanente, the city and county of San Francisco, LinkedIn Corporation, and the University of California Santa Cruz signed the Center for Environmental Health (CEH)'s pledge to "preferentially purchase furniture made without toxic chemicals," including "fluorinated stain treatments" (i.e., PFASs) (CEH 2017). In September 2019, Home Depot announced that it will stop selling carpets and rugs containing PFASs in the U.S. and Canada beginning in December 2019 (Home Depot 2019). Similarly, in October 2019, Lowe's announced that it will stop purchasing PFAS-containing carpets and rugs in the U.S. and Canada by January 2020 (Floor Daily 2019).

Research and regulatory efforts in other countries

On January 30, 2018, South Australia became the first state to ban PFAS-containing firefighting foams (Government of South Australia 2018). European countries are also taking a closer look at PFASs under REACH and national regulatory frameworks. The German Federal Environment Agency has made the case for designating shorter-chain PFASs as substances of very high concern (SVHCs) and regulating them under REACH as chemicals of equivalent concern to persistent, bioaccumulative, and toxic (PBT) or very persistent and very bioaccumulative (vPvB) substances (Brendel et al. 2018). ECHA is considering a proposed restriction on long-chain PFAAs, their salts, and precursors (the public consultation period ended June 20, 2018) (ECHA 2018). In April 2017, the trade association Swedish Water, representing Sweden's municipal water supply companies, called for a ban on PFASs in all consumer products because of the threat to drinking water supplies (Chemical Watch 2017a). In June 2017, 37 Swedish government agencies and research institutions signed a memorandum of understanding to collaboratively study PFASs and reduce their associated risks (Chemical Watch 2017b). Other EU member countries, including Norway, Denmark, and Germany, have ongoing

programs to study and regulate the use of PFASs. On February 1, 2018, the European Commission adopted a proposal for a revised drinking water directive that adds the entire class of PFASs to its list of regulated compounds (European Commission 2018).

Stockholm Convention on Persistent Organic Pollutants

The Stockholm Convention on Persistent Organic Pollutants (POPs), which the United States has not ratified, is a global treaty to protect human health and the environment from chemicals that remain intact in the environment for long periods, become widely distributed geographically, accumulate in the fatty tissue of humans and wildlife, and have harmful impacts on human health or the environment. In 2009, PFOS, its salts, and perfluorooctane sulfonyl fluoride were added to Annex B of the Stockholm Convention, which restricts the production and use, as well as import and export, of listed POPs (SSCCH 2017). Production and use of these chemicals is still allowed for acceptable purposes after Annex B listing, which included use in carpets until 2015.

The EU Council has also submitted a proposal to list PFOA, its salts, and related compounds in Annex A of the Stockholm Convention (European Commission 2015). Following the recommendation of the POPs Review Committee (UNEP/POPS/POPRC 2017), the Conference of the Parties decided in the spring of 2019 to list PFOA, its salts, and PFOA-related compounds in Annex A or B to the Stockholm Convention, with some time-limited exemptions (UNEP/POPS/POPRC 2019). The POPs Review Committee also agreed in October 2017 with a proposal by Norway to list PFHxS, its salts, and PFHxS-related compounds in Annexes A, B, and/or C to the Convention, and has established a working group to review the proposal further and prepare a draft risk profile.

10. CONCLUSIONS

DTSC has determined that treatments containing any member of the class of PFASs for use on converted textiles or leathers meet the key prioritization criteria²⁴ for listing a Priority Product:

- (1) There must be potential public and/or aquatic, avian, or terrestrial animal or plant organism exposure to the Candidate Chemical(s) in the product; and
- (2) There must be the potential for one or more exposures to contribute to or cause significant or widespread adverse impacts.

Treatments for converted textiles or leathers are widespread in Californian homes and workplaces and can be significant sources of human and ecological PFAS exposures, including human exposures via inhalation during product use and ecological exposures via discharges to wastewater treatment plants.

The degradation products of the PFASs currently used in treatments for converted textiles or leathers have the potential to cause significant and widespread adverse impacts to sensitive subpopulations, including fetuses, infants, young children, and pregnant women; carpet and upholstery cleaners; workers in upholstered furniture, furnishings, clothing, shoes, and carpet stores; auto dealership workers; and auto detailing technicians. Exposure to these degradation products may also adversely impact environmentally sensitive habitats and threatened and endangered species. Given the known hazard traits of all members of the class, replacing currently used PFASs in the treatments for converted textiles or leathers with other PFASs could constitute a regrettable substitution. Hence, this proposal covers treatments containing any member of the class of PFASs intended for use on converted textiles or leathers.

While additional data on product formulations and available alternatives could be helpful in informing a future regulatory proposal, DTSC has sufficient information on potential PFAS exposures from treatments for converted textiles or leathers, and on adverse impacts that could result from these exposures, to support listing this product-chemical combination as a Priority Product.

²⁴ CAL. CODE REGS. tit. 22, § 69503.2(a).

ACRONYMS AND ABBREVIATIONS

PFAS (sub)classes

diPAP	Fluorotelomer phosphate diester
FASA	Perfluoroalkane sulfonamide
FASE	Perfluoroalkane sulfonamide ethanol
FAS(M)AC	Perfluoroalkane sulfonamidoethyl (meth)acrylate
FOSE	Perfluorooctane sulfonamide ethanol
FTAC	Fluorotelomer acrylate
FTAL	Saturated fluorotelomer aldehyde
FTCA	Saturated fluorotelomer carboxylic acid
FTEO	Fluorotelomer ethoxylate
FTEOC	Fluorotelomer ethoxycarboxylate
FTI	Fluorotelomer iodide
FTMAC	Fluorotelomer methacrylate
FTO	Fluorotelomer olefin
FTOH	Fluorotelomer alcohol
FTS	Fluorotelomer sulfonic acid
FTUAL	Unsaturated fluorotelomer aldehyde
FTUCA	Unsaturated fluorotelomer carboxylic acid
PAP	Polyfluoroalkyl phosphate
PFAA	Perfluoroalkyl acid
PFAI	Perfluoroalkyl iodide
PFAS	Perfluoroalkyl and polyfluoroalkyl substance
PFECA	Perfluoroether carboxylic acid
PFESA	Perfluoroether sulfonic acid
PFCA	Perfluoroalkyl carboxylic acid
PFPA	Perfluoroalkyl phosphonic acid
PFPE	Perfluoropolyether
PFPiA	Perfluoroalkyl phosphinic acid
PFSA	Perfluoroalkyl sulfonic acid
PFSI	Perfluoroalkyl sulfinate
PFSiA	Perfluoroalkyl sulfinic acid

Individual PFASs

APFO	Ammonium perfluorooctanoate
FBSA	Perfluorobutane sulfonamide
PFBA	Perfluorobutanoic acid
PFBPA	Perfluorobutyl phosphonic acid
PFBPiA	Perfluorobutyl phosphinic acid
PFBS	Perfluorobutane sulfonic acid
PFDA	Perfluorodecanoic acid
PFDS	Perfluorodecane sulfonic acid
PFDoA	Perfluorododecanoic acid
PFDODPiA	Perfluorododecyl phosphinic acid
PFDPA	Perfluorodecyl phosphonic acid
PFDPiA	Perfluorodecyl phosphinic acid
PFHpA	Perfluoroheptanoic acid
PFHxA	Perfluorohexanoic acid
PFHxDA	Perfluorohexadecanoic acid
PFHxPA	Perfluorohexyl phosphonic acid
PFHxPiA	Perfluorohexyl phosphinic acid
PFHxS	Perfluorohexane sulfonic acid
PFNA	Perfluorononanoic acid
PFOA	Perfluorooctanoate or perfluorooctanoic acid
PFODA	Perfluorooctadecanoic acid
PFOPA	Perfluorooctyl phosphonic acid
PFOPiA	Perfluorooctyl phosphinic acid
PFOS	Perfluorooctane sulfonate or perfluorooctane sulfonic acid
PFPeA	Perfluoropentanoic acid
PFPrA	Perfluoropropanoic acid
PFPrOPrA	Perfluoro-2-propoxypropanoic acid (GenX)
PFTeDA	Perfluorotetradecanoic acid
PFTrDA	Perfluorotridecanoic acid
PFUnA	Perfluoroundecanoic acid
PTFE	Polytetrafluoroethylene
PVDF	Polyvinylidene fluoride
PVF	Polyvinyl fluoride
TFA	Tetrafluoroacetic acid
TFMS	Trifluoromethane sulfonic acid

Other abbreviations used in this document

AA	Alternatives Analysis
AFFF	Aqueous film-forming foam
ATSDR	Agency for Toxic Substances and Disease Registry
BAF	Bioaccumulation factor
BCF	Bioconcentration factor
BMF	Biomagnification factor
C	Carbon
CAS	Chemical Abstracts Service
CCM	Creative Chemical Manufacturers
CDC	Centers for Disease Control and Prevention
CECBP	California Environmental Contaminant Biomonitoring Program
CEH	Center for Environmental Health
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CFCs	Chlorofluorocarbons
DDT	Dichlorodiphenyltrichloroethane
DTSC	Department of Toxic Substances Control
ECHA	European Chemicals Agency
EPA	Environmental Protection Agency
F	Fluorine
HF	Hydrofluoric acid
MCL	Maximum contaminant level
MRL	Minimum Reporting Level
NHANES	National Health and Nutrition Examination Survey
OECD	Organisation for Economic Cooperation and Development
OEHHA	Office of Environmental Health Hazard Assessment
PBT	Persistent, bioaccumulative and toxic
PCB	Polychlorinated biphenyl
POPs	Persistent organic pollutants
PPAR	Peroxisome proliferator-activated receptor
REACH	Registration, Evaluation, Authorisation and Restriction of Chemicals
SCP	Safer Consumer Products
SNUR	Significant New Use Rule
SVHC	Substances of Very High Concern
TDI	Tolerable daily intake
TSCA	Toxic Substances Control Act
UCMR 3	Unregulated Contaminant Monitoring Rule 3

U.S.	United States
vPvB	Very persistent, very bioaccumulative
WWTP	Wastewater treatment plan

Units

°C	Degrees Celsius
µg/kg	Micrograms per kilogram
g	Grams
g/L	Grams per liter
kg	Kilograms
kg/yr	Kilograms per year
kJ/mol	Kilojoules per mol
L/kg	Liters per kilogram
m	Meters
m ³	Meters cubed
mg/day	Milligrams per day
mg/kg/day	Milligrams per kilogram per day
ng/day	Nanograms per day
ng/g	Nanograms per gram
ng/kg	Nanograms per kilogram
ng/L	Nanograms per liter
ng/mL	Nanograms per milliliter
ng/m ²	Nanograms per meter squared
ng/m ³	Nanograms per meter cubed
Pa	Pascal
ppt	Parts per trillion

REFERENCES

- Aas CB, Fuglei E, Herzke D, et al (2014) Effect of body condition on tissue distribution of perfluoroalkyl substances (PFASs) in Arctic fox (*Vulpes lagopus*). *Environmental Science & Technology* 48:11654–11661. <https://doi.org/10.1021/es503147n>
- Abell R (2002) Conservation biology for the biodiversity crisis: a freshwater follow-up. *Conservation Biology* 16:1435–1437. <https://doi.org/10.1046/j.1523-1739.2002.01532.x>
- Adams J, Houde M, Muir D, et al (2008) Land use and the spatial distribution of perfluoroalkyl compounds as measured in the plasma of bottlenose dolphins (*Tursiops truncatus*). *Marine Environmental Research* 66:430–437. <https://doi.org/10.1016/j.marenvres.2008.07.004>
- Ahrens L (2011) Polyfluoroalkyl compounds in the aquatic environment: a review of their occurrence and fate. *Journal of Environmental Monitoring* 13:20–31. <https://doi.org/10.1039/C0EM00373E>
- Ahrens L, Barber JL, Xie Z, Ebinghaus R (2009a) Longitudinal and latitudinal distribution of perfluoroalkyl compounds in the surface water of the Atlantic Ocean. *Environmental Science & Technology* 43:3122–3127. <https://doi.org/10.1021/es803507p>
- Ahrens L, Harner T, Shoeib M, et al (2012) Improved characterization of gas-particle partitioning for per- and polyfluoroalkyl substances in the atmosphere using annular diffusion denuder samplers. *Environmental Science & Technology* 46:7199–7206. <https://doi.org/10.1021/es300898s>
- Ahrens L, Hedlund J, Dürig W, Trö R (2016a) Screening of PFASs in groundwater and surface water. Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden, Vatten och miljö: Rapport 2016:2, Sveriges lantbruksuniversitet, Institutionen för vatten och miljö, ISBN 978-91-576-9386-0. 82
- Ahrens L, Rakovic J, Axelson S, Kallenborn R (2016b) Source tracking and impact of per- and polyfluoroalkyl substances at Svalbard. Report prepared by Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden, Norwegian University of Life Sciences (NMBU), Ås, Norway, and University Centre in Svalbard, Longyearbyen, Norway, April 2016. 17
- Ahrens L, Shoeib M, Del Vento S, et al (2011a) Polyfluoroalkyl compounds in the Canadian Arctic atmosphere. *Environmental Chemistry* 8:399. <https://doi.org/10.1071/EN10131>
- Ahrens L, Shoeib M, Harner T, et al (2011b) Wastewater treatment plant and landfills as sources of polyfluoroalkyl compounds to the atmosphere. *Environmental Science & Technology* 45:8098–8105. <https://doi.org/10.1021/es1036173>
- Ahrens L, Yamashita N, Yeung LWY, et al (2009b) Partitioning behavior of per- and polyfluoroalkyl compounds between pore water and sediment in two sediment cores from Tokyo Bay, Japan. *Environmental Science & Technology* 43:6969–6975. <https://doi.org/10.1021/es901213s>

- Alder AC, van der Voet J (2015) Occurrence and point source characterization of perfluoroalkyl acids in sewage sludge. *Chemosphere* 129:62–73. <https://doi.org/10.1016/j.chemosphere.2014.07.045>
- Aleksandrov K, Gehrmann HJ, Hauser M, et al (2019) Waste incineration of polytetrafluoroethylene (PTFE) to evaluate potential formation of per- and poly-fluorinated alkyl substances (PFAS) in flue gas. *Chemosphere* 226:898–906. <https://doi.org/10.1016/j.chemosphere.2019.03.191>
- Alves A, Jacobs G, Vanermen G, et al (2015) New approach for assessing human perfluoroalkyl exposure via hair. *Talanta* 144:574–583. <https://doi.org/10.1016/j.talanta.2015.07.009>
- Annunziato KM, Jantzen CE, Gronske MC, Cooper KR (2019) Subtle morphometric, behavioral and gene expression effects in larval zebrafish exposed to PFHxA, PFHxS and 6:2 FTOH. *Aquatic Toxicology* 208:126–137. <https://doi.org/10.1016/j.aquatox.2019.01.009>
- APHA (2016) American Public Health Association (APHA): Reducing human exposure to highly fluorinated chemicals to protect public health. APHA Policy Number 20163, November 1, 2016. Available at: <https://www.apha.org/policies-and-advocacy/public-health-policy-statements/policy-database/2016/12/21/reducing-human-exposure-to-highly-fluorinated-chemicals>. Accessed 9 Sep 2017
- Appleman TD, Higgins CP, Quiñones O, et al (2014) Treatment of poly- and perfluoroalkyl substances in U.S. full-scale water treatment systems. *Water Research* 51:246–255. <https://doi.org/10.1016/j.watres.2013.10.067>
- Armitage JM, Schenker U, Scheringer M, et al (2009) Modeling the global fate and transport of perfluorooctane sulfonate (PFOS) and precursor compounds in relation to temporal trends in wildlife exposure. *Environmental Science & Technology* 43:9274–9280
- Arnot JA, Gobas FA (2006) A review of bioconcentration factor (BCF) and bioaccumulation factor (BAF) assessments for organic chemicals in aquatic organisms. *Environmental Reviews* 14:257–297. <https://doi.org/10.1139/a06-005>
- Arvaniti OS, Hwang Y, Andersen HR, et al (2015) Reductive degradation of perfluorinated compounds in water using Mg-aminoclay coated nanoscale zero valent iron. *Chemical Engineering Journal* 262:133–139. <https://doi.org/10.1016/j.cej.2014.09.079>
- Arvaniti OS, Stasinakis AS (2015) Review on the occurrence, fate and removal of perfluorinated compounds during wastewater treatment. *Science of The Total Environment* 524–525:81–92. <https://doi.org/10.1016/j.scitotenv.2015.04.023>
- Arvaniti OS, Ventouri EI, Stasinakis AS, Thomaidis NS (2012) Occurrence of different classes of perfluorinated compounds in Greek wastewater treatment plants and determination of their solid–water distribution coefficients. *Journal of Hazardous Materials* 239–240:24–31. <https://doi.org/10.1016/j.jhazmat.2012.02.015>

- Atav R (2018) Dendritic molecules and their use in water repellency treatments of textile materials. In: *Waterproof and Water Repellent Textiles and Clothing*. Elsevier, pp 191–214
- Ateia M, Arifuzzaman M, Pellizzeri S, et al (2019a) Cationic polymer for selective removal of GenX and short-chain PFAS from surface waters and wastewaters at ng/L levels. *Water Research* 163:114874. <https://doi.org/10.1016/j.watres.2019.114874>
- Ateia M, Maroli A, Tharayil N, Karanfil T (2019b) The overlooked short- and ultrashort-chain poly- and perfluorinated substances: A review. *Chemosphere* 220:866–882. <https://doi.org/10.1016/j.chemosphere.2018.12.186>
- ATSDR (2017) Agency for Toxic Substances and Disease Registry (ATSDR): An overview of perfluoroalkyl and polyfluoroalkyl substances and interim guidance for clinicians responding to patient exposure concerns. Interim Guidance, revised on June 7, 2017. National Center for Environmental Health, Centers for Disease Control and Prevention, Washington, DC
- ATSDR (2018a) Agency for Toxic Substances and Disease Registry (ATSDR): Toxicological profile for perfluoroalkyls. Draft for public comment U.S. Department of Health and Human Services, Public Health Service, June 2018
- ATSDR (2015) Agency for Toxic Substances and Disease Registry (ATSDR): Draft toxicological profile for perfluoroalkyls. U.S. Department of Health and Human Services, Public Health Service, August 2015
- ATSDR (2018b) Agency for Toxic Substances and Disease Registry: Minimal Risk Levels (MRLs), August 2018. Available at: https://www.atsdr.cdc.gov/mrls/pdfs/atsdr_mrls.pdf. Accessed 4 Mar 2019
- Audenaert F, Lens H, Rolly D, Vander Elst P (1999) Fluorochemical textile repellents—synthesis and applications: a 3M perspective. *Journal of the Textile Institute* 90:76–94. <https://doi.org/10.1080/00405009908659480>
- Bach CC, Bech BH, Brix N, et al (2015) Perfluoroalkyl and polyfluoroalkyl substances and human fetal growth: a systematic review. *Critical Reviews in Toxicology* 45:53–67. <https://doi.org/10.3109/10408444.2014.952400>
- Banks RE, Smart BE, Tatlow JC (1994) *Organofluorine chemistry: Principles and commercial applications*. Plenum, New York
- Banzhaf S, Filipovic M, Lewis J, et al (2017) A review of contamination of surface-, ground-, and drinking water in Sweden by perfluoroalkyl and polyfluoroalkyl substances (PFASs). *Ambio* 46:335–346. <https://doi.org/10.1007/s13280-016-0848-8>
- Barber JL, Berger U, Chaemfa C, et al (2007) Analysis of per- and polyfluorinated alkyl substances in air samples from Northwest Europe. *Journal of Environmental Monitoring* 9:530. <https://doi.org/10.1039/b701417a>

- Bartell SM, Calafat AM, Lyu C, et al (2010) Rate of decline in serum PFOA concentrations after granular activated carbon filtration at two public water systems in Ohio and West Virginia. *Environmental Health Perspectives* 118:222–228. <https://doi.org/10.1289/ehp.0901252>
- Bečanová J, Melymuk L, Vojta Š, et al (2016) Screening for perfluoroalkyl acids in consumer products, building materials and wastes. *Chemosphere* 164:322–329. <https://doi.org/10.1016/j.chemosphere.2016.08.112>
- Beekman M, Zweers P, Muller A, et al (2016) Evaluation of substances used in the GenX technology by Chemours. Dordrecht. RIVM Letter Report 2006-0174. National Institute for Public Health and the Environment, Ministry of Health, Welfare and Sport, the Netherlands
- Beesoon S, Genuis SJ, Benskin JP, Martin JW (2012) Exceptionally high serum concentrations of perfluorohexanesulfonate in a Canadian family are linked to home carpet treatment applications. *Environmental Science & Technology* 46:12960–12967. <https://doi.org/10.1021/es3034654>
- Benskin JP, Li B, Ikonomidou MG, et al (2012a) Per- and polyfluoroalkyl substances in landfill leachate: Patterns, time trends, and sources. *Environmental Science & Technology* 46:11532–11540. <https://doi.org/10.1021/es302471n>
- Benskin JP, Muir DC, Scott BF, et al (2012b) Perfluoroalkyl acids in the Atlantic and Canadian Arctic Oceans. *Environmental Science & Technology* 46:5815–5823. <https://doi.org/10.1021/es300578x>
- Berger U, Glynn A, Holmström KE, et al (2009) Fish consumption as a source of human exposure to perfluorinated alkyl substances in Sweden – Analysis of edible fish from Lake Vättern and the Baltic Sea. *Chemosphere* 76:799–804. <https://doi.org/10.1016/j.chemosphere.2009.04.044>
- Björklund JA, Thuresson K, de Wit CA (2009) Perfluoroalkyl compounds (PFCs) in indoor dust: concentrations, human exposure estimates, and sources. *Environmental Science & Technology* 43:2276–2281. <https://doi.org/10.1021/es803201a>
- Björnsdotter MK, Yeung LWY, Kärrman A, Jogsten IE (2019) Ultra-short-chain perfluoroalkyl acids including trifluoromethane sulfonic acid in water connected to known and suspected point sources in Sweden. *Environmental Science & Technology*. <https://doi.org/10.1021/acs.est.9b02211>
- Blaine AC, Rich CD, Hundal LS, et al (2013) Uptake of perfluoroalkyl acids into edible crops via land applied biosolids: field and greenhouse studies. *Environmental Science & Technology* 47:14062–14069. <https://doi.org/10.1021/es403094q>
- Blaine AC, Rich CD, Sedlacko EM, et al (2014a) Perfluoroalkyl acid distribution in various plant compartments of edible crops grown in biosolids-amended soils. *Environmental Science & Technology* 48:7858–7865. <https://doi.org/10.1021/es500016s>

- Blaine AC, Rich CD, Sedlacko EM, et al (2014b) Perfluoroalkyl acid uptake in lettuce (*Lactuca sativa*) and strawberry (*Fragaria ananassa*) irrigated with reclaimed water. *Environ Sci Technol* 48:14361–14368. <https://doi.org/10.1021/es504150h>
- Blanc M, Kärrman A, Kukucka P, et al (2017) Mixture-specific gene expression in zebrafish (*Danio rerio*) embryos exposed to perfluorooctane sulfonic acid (PFOS), perfluorohexanoic acid (PFHxA) and 3,3',4,4',5-pentachlorobiphenyl (PCB126). *Science of The Total Environment* 590–591:249–257. <https://doi.org/10.1016/j.scitotenv.2017.02.232>
- Blum A (2016) Tackling toxics. *Science* 351:1117–1117. <https://doi.org/10.1126/science.aaf5468>
- Blum A, Balan SA, Scheringer M, et al (2015) The Madrid statement on poly- and perfluoroalkyl substances (PFASs). *Environmental Health Perspectives* 123:A107-111. <https://doi.org/10.1289/ehp.1509934>
- Bohlin-Nizzetto P, Hanssen L, Herzke D (2015) PFASs in house dust. Norwegian Institute for Air Research (NILU), Scientific Report NILU OR 29/2015
- Boiteux V, Dauchy X, Bach C, et al (2017) Concentrations and patterns of perfluoroalkyl and polyfluoroalkyl substances in a river and three drinking water treatment plants near and far from a major production source. *Science of The Total Environment* 583:393–400. <https://doi.org/10.1016/j.scitotenv.2017.01.079>
- Boone JS, Vigo C, Boone T, et al (2019) Per- and polyfluoroalkyl substances in source and treated drinking waters of the United States. *Science of The Total Environment* 653:359–369. <https://doi.org/10.1016/j.scitotenv.2018.10.245>
- Borg D, Lund BO, Lindquist NG, Håkansson H (2013) Cumulative health risk assessment of 17 perfluoroalkylated and polyfluoroalkylated substances (PFASs) in the Swedish population. *Environment International* 59:112–123. <https://doi.org/10.1016/j.envint.2013.05.009>
- Bowman J (2017) PFASs in carpet and furniture applications. Presentation at the DTSC Public Workshop on Perfluoroalkyl and Polyfluoroalkyl Substances (PFASs) in Carpets, Rugs, Indoor Upholstered Furniture, and Their Care and Treatment Products, January 31, 2017, Sacramento, CA
- Boyle B (2016) Congressman Brendan Boyle introduces legislation to require enforceable nationwide standard for perfluorinated compounds in drinking water. Press release, September 27, 2016. Available at: <https://boyle.house.gov/media-center/press-releases/congressman-brendan-boyle-introduces-legislation-require-enforceable>. Accessed 15 Aug 2017
- Bradley PM, Journey CA, Romanok KM, et al (2017) Expanded target-chemical analysis reveals extensive mixed-organic-contaminant exposure in U.S. streams. *Environmental Science & Technology* 51:4792–4802. <https://doi.org/10.1021/acs.est.7b00012>

- Brendel S, Fetter É, Staude C, et al (2018) Short-chain perfluoroalkyl acids: environmental concerns and a regulatory strategy under REACH. *Environmental Sciences Europe* 30:9. <https://doi.org/10.1186/s12302-018-0134-4>
- Buck RC, Franklin J, Berger U, et al (2011) Perfluoroalkyl and polyfluoroalkyl substances in the environment: Terminology, classification, and origins. *Integrated Environmental Assessment and Management* 7:513–541. <https://doi.org/10.1002/ieam.258>
- Buck RC, Murphy PM, Pabon M (2012) Chemistry, properties, and uses of commercial fluorinated surfactants. In: Knepper TP, Lange FT (eds) *Handbook of Environmental Chemistry, Volume 17, Polyfluorinated Chemicals and Transformation Products*. Springer, New York, pp 1–24
- Burkemper JL, Aweda TA, Rosenberg AJ, et al (2017) Radiosynthesis and biological distribution of ¹⁸F-labeled perfluorinated alkyl substances. *Environmental Science & Technology Letters* 4:211–215. <https://doi.org/10.1021/acs.estlett.7b00042>
- Busch J, Ahrens L, Sturm R, Ebinghaus R (2010) Polyfluoroalkyl compounds in landfill leachates. *Environmental Pollution* 158:1467–1471. <https://doi.org/10.1016/j.envpol.2009.12.031>
- Butenhoff JL, Bjork JA, Chang SC, et al (2012) Toxicological evaluation of ammonium perfluorobutyrate in rats: twenty-eight-day and ninety-day oral gavage studies. *Reproductive Toxicology* 33:513–530. <https://doi.org/10.1016/j.reprotox.2011.08.004>
- Butenhoff JL, Kennedy GL, Jr., Hinderliter PM (2004) Pharmacokinetics of perfluorooctanoate in cynomolgus monkeys. *Toxicological Sciences* 82:394–406. <https://doi.org/10.1093/toxsci/kfh302>
- Butt CM, Berger U, Bossi R, Tomy GT (2010) Levels and trends of poly- and perfluorinated compounds in the Arctic environment. *Science of The Total Environment* 408:2936–2965. <https://doi.org/10.1016/j.scitotenv.2010.03.015>
- Butt CM, Mabury SA, Kwan M, et al (2008) Spatial trends of perfluoroalkyl compounds in ringed seals (*Phoca hispida*) from the Canadian Arctic. *Environmental Toxicology and Chemistry* 27:542. <https://doi.org/10.1897/07-428.1>
- Butt CM, Muir DCG, Mabury SA (2014) Biotransformation pathways of fluorotelomer-based polyfluoroalkyl substances: A review: Biotransformation pathways of fluorotelomer compounds. *Environ Toxicol Chem* 33:243–267. <https://doi.org/10.1002/etc.2407>
- C8 Science Panel (2011) Probable link evaluation of pregnancy induced hypertension and preeclampsia. December 5, 2011. Available at: http://www.c8sciencepanel.org/pdfs/Probable_Link_C8_PIH_5Dec2011.pdf. Accessed 9 Oct 2017

- C8 Science Panel (2012) Probable link evaluation of cancer. April 15, 2012. Available at: http://www.c8sciencepanel.org/pdfs/Probable_Link_C8_Cancer_16April2012_v2.pdf. Accessed 9 Oct 2017
- Calafat AM (2017) Public Comments to presentations during DTSC Public Workshop on Perfluoroalkyl and Polyfluoroalkyl Substances (PFASs) in Carpets, Rugs, Indoor Upholstered Furniture, and Their Care and Treatment Products. January 31, 2017, Sacramento, CA. The workshop recording is available at: <https://www.youtube.com/watch?v=MryvxUSlxA&feature=youtu.be>. Accessed 29 Aug 2018
- Calafat AM, Kato K, Hubbard K, et al (2019) Legacy and alternative per- and polyfluoroalkyl substances in the U.S. general population: Paired serum-urine data from the 2013–2014 National Health and Nutrition Examination Survey. *Environment International* 131:105048. <https://doi.org/10.1016/j.envint.2019.105048>
- Calafat AM, Kuklennyik Z, Reidy JA, et al (2007) Serum concentrations of 11 polyfluoroalkyl compounds in the U.S. population: Data from the National Health and Nutrition Examination Survey (NHANES). *Environmental Science & Technology* 41:2237–2242. <https://doi.org/10.1021/es062686m>
- California Employment Development Department (2019) Labor Market Information - Long Term Projections (Ten-years). Available at: <https://www.labormarketinfo.edd.ca.gov/data/employment-projections.html>. Accessed 26 Sep 2019
- California State Water Resources Control Board (2019) Per- and Polyfluoroalkyl Substances (PFAS). <https://www.waterboards.ca.gov/pfas/>. Accessed 29 Oct 2019
- CalRecycle (2015) 2014 Disposal-Facility-Based Characterization of Solid Waste in California
- CalRecycle (2017) California Department of Resources Recycling and Recovery (CalRecycle): Carpet Materials Management. Last updated October 3, 2017. Available at: <http://www.calrecycle.ca.gov/carpet/default.htm>. Accessed 24 Sep 2019
- Campo J, Lorenzo M, Pérez F, et al (2016) Analysis of the presence of perfluoroalkyl substances in water, sediment and biota of the Jucar River (E Spain). Sources, partitioning and relationships with water physical characteristics. *Environmental Research* 147:503–512. <https://doi.org/10.1016/j.envres.2016.03.010>
- Cao W, Liu X, Liu X, et al (2018) Perfluoroalkyl substances in umbilical cord serum and gestational and postnatal growth in a Chinese birth cohort. *Environment International* 116:197–205. <https://doi.org/10.1016/j.envint.2018.04.015>
- CARE (2012) CARE operational guidelines for handling and processing post consumer carpet. Available at: <http://www.calrecycle.ca.gov/ConDemo/Carpet/CAREOperat.pdf>. Accessed 18 May 2018

- Cariou R, Veyrand B, Yamada A, et al (2015) Perfluoroalkyl acid (PFAA) levels and profiles in breast milk, maternal and cord serum of French women and their newborns. *Environment International* 84:71–81. <https://doi.org/10.1016/j.envint.2015.07.014>
- Casal P, González-Gaya B, Zhang Y, et al (2017) Accumulation of perfluoroalkylated substances in oceanic plankton. *Environmental Science & Technology* 51:2766–2775. <https://doi.org/10.1021/acs.est.6b05821>
- CCSP (2016) California Carpet Stewardship Program (CCSP): CARE California Carpet Stewardship Program Annual Report, January 2016 – December 2016 Carpet America Recovery Effort (CARE), Dalton, GA. Available at: <http://www.calrecycle.ca.gov/Carpet/AnnualRpts/2016/CARE2016.pdf>. Accessed 3 Oct 2017
- CDC (2018) Centers for Disease Control and Prevention (CDC): Fourth National Report on Human Exposure to Environmental Chemicals Updated Tables, March 2018, Volume One. Centers for Disease Control and Prevention, U.S. Department of Health and Human Services. Available at: https://www.cdc.gov/exposurereport/pdf/FourthReport_UpdatedTables_Volume1_Mar2018.pdf. Accessed 21 May 2018
- CDC (2014) Centers for Disease Control and Prevention (CDC): Breastfeeding Report Card United States, 2014. Available at: <https://www.cdc.gov/breastfeeding/pdf/2014breastfeedingreportcard.pdf>. Accessed 9 Oct 2017
- CEC (2017) Commission for Environmental Cooperation (CEC): Furthering the understanding of the migration of chemicals from consumer products – A study of per- and polyfluoroalkyl substances (PFASs) in clothing, apparel, and children’s items. Montreal, Canada, 201pp
- CECBP (2015a) California Environmental Contaminant Biomonitoring Program (CECBP): Potential Priority Chemicals: Perfluoroalkyl and Polyfluoroalkyl Substances (PFASs). Available at: https://biomonitoring.ca.gov/sites/default/files/downloads/PotentialPriority_PFASs_111815.pdf. Accessed 17 Sep 2019
- CECBP (2017) California Environmental Contaminant Biomonitoring Program (CECBP): Results for perfluorochemicals (PFCs). Available at: <https://biomonitoring.ca.gov/results/chemical/154>. Accessed 8 Dec 2017
- CECBP (2015b) California Environmental Contaminant Biomonitoring Program (CECBP): California Teachers Study (CTS). Sample collection date: 2011 - ongoing. Results last updated on September 22, 2015. Available at: <https://biomonitoring.ca.gov/results/projects/411>. Accessed 20 Oct 2017
- CECBP (2018) California Environmental Contaminant Biomonitoring Program (CECBP): Project results for biomonitoring exposures study (BEST). 2. Expanded. Updated March 1, 2018. Available at: <https://biomonitoring.ca.gov/results/projects/494>. Accessed 16 Apr 2018

- CEH (2017) Center for Environmental Health (CEH): Major companies demanding safer office furniture, without harmful chemicals. CEH Press Release, May 25, 2017. Available at: <http://www.ceh.org/news-events/press-releases/content/major-companies-demanding-safer-office-furniture-without-harmful-chemicals/>. Accessed 14 Aug 2017
- Ceretta F (2013) Synthesis and characterization of fluorinated compounds for industrial applications. Ph.D. Dissertation, Università Degli Studi di Padova. 127
- Chang ET, Adami HO, Boffetta P, et al (2014) A critical review of perfluorooctanoate and perfluorooctanesulfonate exposure and cancer risk in humans. *Critical Reviews in Toxicology* 44:1–81. <https://doi.org/10.3109/10408444.2014.905767>
- Chang SC, Das K, Ehresman DJ, et al (2008) Comparative pharmacokinetics of perfluorobutyrate in rats, mice, monkeys, and humans and relevance to human exposure via drinking water. *Toxicological Sciences* 104:40–53. <https://doi.org/10.1093/toxsci/kfn057>
- Chang SC, Noker PE, Gorman GS, et al (2012) Comparative pharmacokinetics of perfluorohexanesulfonate (PFHxS) in rats, mice, and monkeys. *Reproductive Toxicology* 33:441–451. <https://doi.org/10.1016/j.reprotox.2011.07.004>
- Chapman (2016) Alabama Water Utility sues Dalton carpet makers. *The Atlanta Journal-Constitution*, Sept. 26, 2016. Available at: <http://www.ajc.com/news/state--regional/alabama-water-utility-sues-dalton-carpet-makers/Yh6dnovTsKxe9vMij7ahAN/>. Accessed 15 Aug 2017
- Chawaga P (2017) Michigan town uncovers PFC contamination 400 times EPA-advised limits. News Feature in *Water Online*, September 13, 2017. Available at: <https://www.wateronline.com/doc/michigan-town-the-latest-to-uncover-pfc-contamination-0001>. Accessed 19 Oct 2017
- Chemical Watch (2017a) Swedish water body urges ban on goods potentially containing PFAS. April 24, 2017. Available at: <https://chemicalwatch.com/55406>. Accessed 15 Aug 2017
- Chemical Watch (2017b) Swedish authorities agree to tackle PFAS. June 1, 2017. Available at: <https://chemicalwatch.com/56578/swedish-authorities-agree-to-tackle-pfas>. Accessed 15 Aug 2017
- Chen F, Yin S, Kelly BC, Liu W (2017a) Chlorinated polyfluoroalkyl ether sulfonic acids in matched maternal, cord, and placenta samples: a study of transplacental transfer. *Environmental Science & Technology* 51:6387–6394. <https://doi.org/10.1021/acs.est.6b06049>
- Chen H, Peng H, Yang M, et al (2017b) Detection, occurrence, and fate of fluorotelomer alcohols in municipal wastewater treatment plants. *Environmental Science & Technology* 51:8953–8961. <https://doi.org/10.1021/acs.est.7b00315>

- Chen H, Reinhard M, Nguyen VT, Gin KY-H (2016) Reversible and irreversible sorption of perfluorinated compounds (PFCs) by sediments of an urban reservoir. *Chemosphere* 144:1747–1753. <https://doi.org/10.1016/j.chemosphere.2015.10.055>
- Chen H, Reinhard M, Yin T, et al (2019) Multi-compartment distribution of perfluoroalkyl and polyfluoroalkyl substances (PFASs) in an urban catchment system. *Water Research* 154:227–237. <https://doi.org/10.1016/j.watres.2019.02.009>
- Chen YM, Guo LH (2009) Fluorescence study on site-specific binding of perfluoroalkyl acids to human serum albumin. *Archives of Toxicology* 83:255–261. <https://doi.org/10.1007/s00204-008-0359-x>
- Christensen KY, Raymond M, Blackowicz M, et al (2017) Perfluoroalkyl substances and fish consumption. *Environmental Research* 154:145–151. <https://doi.org/10.1016/j.envres.2016.12.032>
- Chu S, Letcher RJ (2017) Side-chain fluorinated polymer surfactants in aquatic sediment and biosolid-augmented agricultural soil from the Great Lakes basin of North America. *Science of The Total Environment* 607–608:262–270. <https://doi.org/10.1016/j.scitotenv.2017.06.252>
- Chu S, Letcher RJ, McGoldrick DJ, Backus SM (2016) A new fluorinated surfactant contaminant in biota: perfluorobutane sulfonamide in several fish species. *Environmental Science & Technology* 50:669–675. <https://doi.org/10.1021/acs.est.5b05058>
- Clara M, Scharf S, Weiss S, et al (2008) Emissions of perfluorinated alkylated substances (PFAS) from point sources—identification of relevant branches. *Water Science and Technology* 58:59–66. <https://doi.org/10.2166/wst.2008.641>
- Codling G, Vogt A, Jones PD, et al (2014) Historical trends of inorganic and organic fluorine in sediments of Lake Michigan. *Chemosphere* 114:203–209. <https://doi.org/10.1016/j.chemosphere.2014.03.080>
- Coggan TL, Anumol T, Pyke J, et al (2019) A single analytical method for the determination of 53 legacy and emerging per- and polyfluoroalkyl substances (PFAS) in aqueous matrices. *Analytical and Bioanalytical Chemistry* 411:3507–3520. <https://doi.org/10.1007/s00216-019-01829-8>
- Concawe (2016) Environmental fate and effects of poly and perfluoroalkyl substances (PFAS). Concawe Soil and Groundwater Taskforce, Brussels, Belgium. Prepared by ARCADIS: T. Pancras, G. Schrauwen, T. Held, K. Baker, I. Ross, H. Slenders, Report # 8/16. Accessed March 15, 2017
- Corsini E, Luebke RW, Germolec DR, DeWitt JC (2014) Perfluorinated compounds: emerging POPs with potential immunotoxicity. *Toxicology Letters* 230:263–270. <https://doi.org/10.1016/j.toxlet.2014.01.038>

- Cousins IT, Kong D, Vestergren R (2011) Reconciling measurement and modelling studies of the sources and fate of perfluorinated carboxylates. *Environ Chem* 8:339. <https://doi.org/10.1071/EN10144>
- Cousins IT, Vestergren R, Wang Z, et al (2016) The precautionary principle and chemicals management: the example of perfluoroalkyl acids in groundwater. *Environment International* 94:331–340. <https://doi.org/10.1016/j.envint.2016.04.044>
- Crone BC, Speth TF, Wahman DG, et al (2019) Occurrence of per- and polyfluoroalkyl substances (PFAS) in source water and their treatment in drinking water. *Critical Reviews in Environmental Science and Technology* 49:2359–2396. <https://doi.org/10.1080/10643389.2019.1614848>
- Cui Q, Pan Y, Zhang H, et al (2018) Occurrence and tissue distribution of novel perfluoroether carboxylic and sulfonic acids and legacy per/polyfluoroalkyl substances in Black-Spotted Frog (*Pelophylax nigromaculatus*). *Environmental Science & Technology* 52:982–990. <https://doi.org/10.1021/acs.est.7b03662>
- Dairkee SH, Luciani-Torres G, Moore DH, et al (2018) A ternary mixture of common chemicals perturbs benign human breast epithelial cells more than the same chemicals do individually. *Toxicological Sciences* 165:131–144. <https://doi.org/10.1093/toxsci/kfy126>
- Dalahmeh S, Tirgani S, Komakech AJ, et al (2018) Per- and polyfluoroalkyl substances (PFASs) in water, soil and plants in wetlands and agricultural areas in Kampala, Uganda. *Science of The Total Environment* 631–632:660–667. <https://doi.org/10.1016/j.scitotenv.2018.03.024>
- Danish Environmental Protection Agency (2015a) Short-chain polyfluoroalkyl substances (PFAS). A literature review of information on human health effects and environmental fate and effect aspects of short-chain PFAS. Project No. 1707, Copenhagen, Denmark
- Danish Environmental Protection Agency (2015b) Alternatives to perfluoroalkyl and polyfluoroalkyl substances (PFAS) in textiles. Survey of chemical substances in consumer products No. 137. Copenhagen, Denmark
- Das KP, Grey BE, Zehr RD, et al (2008) Effects of perfluorobutyrate exposure during pregnancy in the mouse. *Toxicological Sciences* 105:173–181. <https://doi.org/10.1093/toxsci/kfn099>
- Dasu K, Lee LS (2016) Aerobic biodegradation of toluene-2,4-di(8:2 fluorotelomer urethane) and hexamethylene-1,6-di(8:2 fluorotelomer urethane) monomers in soils. *Chemosphere* 144:2482–2488. <https://doi.org/10.1016/j.chemosphere.2015.11.021>
- Dauchy X, Boiteux V, Bach C, et al (2017) Mass flows and fate of per- and polyfluoroalkyl substances (PFASs) in the wastewater treatment plant of a fluorochemical manufacturing facility. *Science of The Total Environment* 576:549–558. <https://doi.org/10.1016/j.scitotenv.2016.10.130>
- Dauchy X, Boiteux V, Rosin C, Munoz JF (2012) Relationship between industrial discharges and contamination of raw water resources by perfluorinated compounds: part II: Case study of a

fluorotelomer polymer manufacturing plant. *Bulletin of Environmental Contamination and Toxicology* 89:525–530. <https://doi.org/10.1007/s00128-012-0704-x>

Davis M (2016) Personal communication via a meeting with DTSC Safer Consumer Products Branch staff. July 1, 2016, Berkeley, CA

De Silva AO, Spencer C, Ho KCD, et al (2016) Perfluoroalkylphosphinic acids in northern pike (*Esox lucius*), double-crested cormorants (*Phalacrocorax auritus*), and bottlenose dolphins (*Tursiops truncatus*) in relation to other perfluoroalkyl acids. *Environmental Science & Technology* 50:10903–10913. <https://doi.org/10.1021/acs.est.6b03515>

De Silva AO, Spencer C, Scott BF, et al (2011) Detection of a cyclic perfluorinated acid, perfluoroethylcyclohexane sulfonate, in the Great Lakes of North America. *Environmental Science & Technology* 45:8060–8066. <https://doi.org/10.1021/es200135c>

Dean JA (1999) *Lange's Handbook of Chemistry*, 15. ed. McGraw-Hill, New York, NY

Del Vento S, Halsall C, Gioia R, et al (2012) Volatile per- and polyfluoroalkyl compounds in the remote atmosphere of the western Antarctic Peninsula: an indirect source of perfluoroalkyl acids to Antarctic waters. *Atmospheric Pollution Research* 3:450–455

D'eon JC, Crozier PW, Furdui VI, et al (2009) Perfluorinated phosphonic acids in Canadian surface waters and wastewater treatment plant effluent: discovery of a new class of perfluorinated acids. *Environmental Toxicology and Chemistry* 28:2101. <https://doi.org/10.1897/09-048.1>

D'eon JC, Mabury SA (2011) Exploring indirect sources of human exposure to perfluoroalkyl carboxylates (PFCAs): Evaluating uptake, elimination, and biotransformation of polyfluoroalkyl phosphate esters (PAPs) in the rat. *Environmental Health Perspectives* 119:344–350. <https://doi.org/10.1289/ehp.1002409>

D'Eon JC, Mabury SA (2007) Production of perfluorinated carboxylic acids (PFCAs) from the biotransformation of polyfluoroalkyl phosphate surfactants (PAPS): Exploring routes of human contamination. *Environmental Science & Technology* 41:4799–4805. <https://doi.org/10.1021/es070126x>

D'eon JC, Mabury SA (2010) Uptake and elimination of perfluorinated phosphonic acids in the rat. *Environmental Toxicology and Chemistry* n/a-n/a. <https://doi.org/10.1002/etc.167>

DeWitt JC (2015) Toxicological Effects of Perfluoroalkyl and Polyfluoroalkyl Substances. In: *Molecular and Integrative Toxicology*. Available at SpringerLink (Online service): <http://link.springer.com/10.1007/978-3-319-15518-0>. Accessed 9 Oct 2017

DeWitt JC, Peden-Adams MM, Keil DE (2015) Immunotoxic effects of perfluoroalkylated compounds: mechanisms of action. Wiley-VCH Verlag GmbH, Weinheim. In: Corsini E, Loveren H van (eds) *Molecular Immunotoxicology*. pp 263–284

- Dietz R, Bossi R, Rigét FF, et al (2008) Increasing perfluoroalkyl contaminants in East Greenland polar bears (*Ursus maritimus*): a new toxic threat to the Arctic bears. *Environmental Science & Technology* 42:2701–2707. <https://doi.org/10.1021/es7025938>
- Ding G, Peijnenburg WJGM (2013) Physicochemical properties and aquatic toxicity of poly- and perfluorinated compounds. *Critical Reviews in Environmental Science and Technology* 43:598–678. <https://doi.org/10.1080/10643389.2011.627016>
- Ding G, Wouterse M, Baerselman R, Peijnenburg WJGM (2012a) Toxicity of polyfluorinated and perfluorinated compounds to lettuce (*Lactuca sativa*) and green algae (*Pseudokirchneriella subcapitata*). *Archives of Environmental Contamination and Toxicology* 62:49–55. <https://doi.org/10.1007/s00244-011-9684-9>
- Ding GH, Frömel T, van den Brandhof EJ, et al (2012b) Acute toxicity of poly- and perfluorinated compounds to two cladocerans, *Daphnia magna* and *Chydorus sphaericus*. *Environmental Toxicology and Chemistry* 31:605–610. <https://doi.org/10.1002/etc.1713>
- Dinglasan-Panlilio MJ, Mabury SA (2006) Significant residual fluorinated alcohols present in various fluorinated materials. *Environmental Science & Technology* 40:1447–1453
- Domingo JL, Nadal M (2019) Human exposure to per- and polyfluoroalkyl substances (PFAS) through drinking water: a review of the recent scientific literature. *Environmental Research* 177:108648. <https://doi.org/10.1016/j.envres.2019.108648>
- Domingo JL, Nadal M (2017) Per- and Polyfluoroalkyl Substances (PFASs) in Food and Human Dietary Intake: A Review of the Recent Scientific Literature. *J Agric Food Chem* 65:533–543. <https://doi.org/10.1021/acs.jafc.6b04683>
- Dreyer A, Kirchgeorg T, Weinberg I, Matthias V (2015) Particle-size distribution of airborne poly- and perfluorinated alkyl substances. *Chemosphere* 129:142–149. <https://doi.org/10.1016/j.chemosphere.2014.06.069>
- Drottar KR, Van Hoven RL, Krueger HO (2001) Perfluorooctanesulfonate, potassium salt (PFOS): a flow-through bioconcentration test with the bluegill (*Lepomis macrochirus*). Wildlife International, Ltd., Project No. 454A-134, EPA Docket AR226-1030a042
- Du J, Cai J, Wang S, You H (2017) Oxidative stress and apoptosis to Zebrafish (*Danio rerio*) embryos exposed to perfluorooctane sulfonate (PFOS) and ZnO nanoparticles. *International Journal of Occupational Medicine and Environmental Health*. <https://doi.org/10.13075/ijomeh.1896.00669>
- ECHA (2014) European Chemicals Agency (ECHA): Annex XV restriction report. Proposal for a restriction: Perfluorooctanoic acid (PFOA), PFOA salts and PFOA-related substances. Version 1.0, October 17, 2014. Available at: <https://echa.europa.eu/documents/10162/e9cddee6-3164-473d-b590-8fcf9caa50e7>. Accessed 26 Sep 2019

- ECHA (2017) European Chemicals Agency (ECHA): Candidate list of substances of very high concern for authorisation. Available at: <https://echa.europa.eu/web/guest/candidate-list-table>. Accessed September 1, 2017
- ECHA (2018) European Chemical Agency (ECHA): Submitted restrictions under consideration. Available at: <https://www.echa.europa.eu/web/guest/restrictions-under-consideration/-/substance-rev/18115/term>. Accessed 7 Feb 2018
- EFSA (2011) The European Food Safety Authority (EFSA): Scientific opinion on the safety evaluation of the substance, 3H-perfluoro-3-[(3-methoxy-propoxy)propanoic acid], ammonium salt, CAS No. 958445-44-8, for use in food contact materials. EFSA Panel on food contact materials, enzymes, flavourings and processing aids (CEF). EFSA Journal 9:2182. <https://doi.org/10.2903/j.efsa.2011.2182>
- EFSA (2008) The European Food Safety Authority (EFSA): Perfluorooctane sulfonate (PFOS), perfluorooctanoic acid (PFOA) and their salts: Scientific opinion of the scientific panel on contaminants in the food chain. EFSA Journal 653:1-131
- EFSA (2018) The European Food Safety Authority (EFSA): Risk to human health related to the presence of perfluorooctane sulfonic acid and perfluorooctanoic acid in food. EFSA Journal 16:. <https://doi.org/10.2903/j.efsa.2018.5194>
- EFSA (European Food Safety Authority) (2012) Perfluoroalkylated substances in food: occurrence and dietary exposure. EFSA Journal 10:. <https://doi.org/10.2903/j.efsa.2012.2743>
- Eftaiha AF, Brunet SMK, Paige MF (2012) Influence of film composition on the morphology, mechanical properties, and surfactant recovery of phase-separated phospholipid-perfluorinated fatty acid mixed monolayers. Langmuir 28:15150–15159. <https://doi.org/10.1021/la3026655>
- Egghy PP, Lorber M (2011) An assessment of the exposure of Americans to perfluorooctane sulfonate: A comparison of estimated intake with values inferred from NHANES data. Journal of Exposure Science & Environmental Epidemiology 21:150–168. <https://doi.org/10.1038/jes.2009.73>
- Eggen T, Moeder M, Arukwe A (2010) Municipal landfill leachates: A significant source for new and emerging pollutants. Science of The Total Environment 408:5147–5157. <https://doi.org/10.1016/j.scitotenv.2010.07.049>
- Eggers Pedersen K, Basu N, Letcher R, et al (2015) Brain region-specific perfluoroalkylated sulfonate (PFSA) and carboxylic acid (PFCA) accumulation and neurochemical biomarker responses in east Greenland polar Bears (*Ursus maritimus*). Environmental Research 138:22–31. <https://doi.org/10.1016/j.envres.2015.01.015>
- Ellis DA, Martin JW, De Silva AO, et al (2004) Degradation of fluorotelomer alcohols: a likely atmospheric source of perfluorinated carboxylic acids. Environmental Science & Technology 38:3316–3321. <https://doi.org/10.1021/es049860w>

- Ellis DA, Martin JW, Mabury SA, et al (2003) Atmospheric Lifetime of Fluorotelomer Alcohols. *Environ Sci Technol* 37:3816–3820. <https://doi.org/10.1021/es034136j>
- Ellison G (2017) Tannery chemicals are 540 times above EPA level in Belmont well. *Grand Rapids News*, September 11, 2017. Available at: http://www.mlive.com/news/grand-rapids/index.ssf/2017/09/wolverine_house_street_well.html. Accessed 19 Oct 2017
- Ericson JI, Nadal M, van Bavel B, et al (2012) Per- and polyfluorinated compounds (PFCs) in house dust and indoor air in Catalonia, Spain: implications for human exposure. *Environment International* 39:172–180. <https://doi.org/10.1016/j.envint.2011.09.004>
- Eryasa B, Grandjean P, Nielsen F, et al (2019) Physico-chemical properties and gestational diabetes predict transplacental transfer and partitioning of perfluoroalkyl substances. *Environment International* 130:104874. <https://doi.org/10.1016/j.envint.2019.05.068>
- Eschauzier C, Raat KJ, Stuyfzand PJ, De Voogt P (2013) Perfluorinated alkylated acids in groundwater and drinking water: identification, origin and mobility. *Science of The Total Environment* 458–460:477–485. <https://doi.org/10.1016/j.scitotenv.2013.04.066>
- Escoruela J, Garreta E, Ramos R, et al (2018) Occurrence of per- and polyfluoroalkyl substances in Calonectris shearwaters breeding along the Mediterranean and Atlantic colonies. *Marine Pollution Bulletin* 131:335–340. <https://doi.org/10.1016/j.marpolbul.2018.04.032>
- Eun H, Yamazaki E, Taniyasu S, et al (2020) Evaluation of perfluoroalkyl substances in field-cultivated vegetables. *Chemosphere* 239:124750. <https://doi.org/10.1016/j.chemosphere.2019.124750>
- Eurofins Eaton Analytical (2017) PFAS monitoring in a post health advisory world-what should we be doing? Poly- and perfluoroalkyl substances (PFASs). Available at: <https://assets.documentcloud.org/documents/4165198/EEA-Study-Documents.pdf>. Accessed 15 Nov 2017
- European Commission (2017) Study for the strategy for a non-toxic environment of the 7th Environment Action Programme, Sub-study d-Very Persistent Chemicals. Prepared by G. Goldenman (Milieu Ltd), with contributions from R. Pedersen, H. Bradley, M. Fernandez, R. Weber, M. Scheringer, and P. Fantke, August 2017. The European Commission, Directorate-General for Environment, Brussels, Belgium. Available at:
- European Commission (2018) Directive of the European Parliament and of the Council on the quality of water intended for human consumption (recast). Brussels, 1.2.2018, COM(2017) 753 final. Available at: <http://eur-lex.europa.eu/legal-content/EN/TXT/?qid=1519210589057&uri=CELEX:52017PC0753>. Accessed 11 Apr 2018
- European Commission (2015) Proposal for a Council decision on the submission, on behalf of the European Union, of a proposal for the listing of additional chemicals in Annex A to the Stockholm Convention on Persistent Organic Pollutants. Brussels, 19.3.2015, COM(2015) 133

final 2015/0066 (NLE). Available at: <http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=COM:2015:133:FIN&qid=1427117617364&from=EN>. Accessed 15 Oct 2017

EWG (2003) Environmental Working Group (EWG): PFCs: Global contaminants: PFCs last forever. Available at: <http://www.ewg.org/research/pfcs-global-contaminants/pfcs-last-forever>. Accessed 20 Jun 2017

Feng M, Qu R, Wei Z, et al (2015) Characterization of the thermolysis products of nafion membrane: a potential source of perfluorinated compounds in the environment. *Scientific Reports* 5:. <https://doi.org/10.1038/srep09859>

Fleisch AF, Rifas-Shiman SL, Mora AM, et al (2017) Early-life exposure to perfluoroalkyl substances and childhood metabolic function. *Environmental Health Perspectives* 125:481–487. <https://doi.org/10.1289/EHP303>

Floor Daily (2019) Lowe's Follows Home Depot in Phasing Out Carpet with PFAS. <https://www.floordaily.net/flooring-news/lowes-follows-home-depot-in-phasing-out-carpet-products-with-pfas>. Accessed 5 Nov 2019

FluoroCouncil (2019) Applications - FluoroTechnology: Critical to Modern Life. Available at: <https://fluorocouncil.com/applications/>. Accessed 18 Sep 2019

FluoroCouncil (2017) How carpet treatments have changed since the Battelle study in 2000 and why there is reduced loss of PFAS during the carpet life span. Information provided to DTSC by the FluoroCouncil in May 2017

Fraser AJ, Webster TF, Watkins DJ, et al (2013) Polyfluorinated compounds in dust from homes, offices, and vehicles as predictors of concentrations in office workers' serum. *Environment International* 60:128–136. <https://doi.org/10.1016/j.envint.2013.08.012>

Fraser AJ, Webster TF, Watkins DJ, et al (2012) Polyfluorinated compounds in serum linked to indoor air in office environments. *Environmental Science & Technology* 46:1209–1215. <https://doi.org/10.1021/es2038257>

Frisbee SJ, Brooks AP, Maher A, et al (2009) The C8 health project: design, methods, and participants. *Environmental Health Perspectives* 117:1873–1882. <https://doi.org/10.1289/ehp.0800379>

Frömel DT, Gremmel C, Dimzon DIK, Weil H (2016) Investigations on the presence and behavior of precursors to perfluoroalkyl substances in the environment as a preparation of regulatory measures. German Environment Agency Report No. (UBA-FB) 002223/ENG. Available at: https://www.umweltbundesamt.de/sites/default/files/medien/378/publikationen/texte_08_2016_investigations_on_the_presence_and_behavior.pdf. Accessed December 20, 2016. 272

- Frömel T, Knepper TP (2010) Biodegradation of fluorinated alkyl substances. In: Whitacre DM, de Voogt P (eds). *Reviews of Environmental Contamination and Toxicology 208: Perfluorinated Alkylated Substances*. Springer, New York:161–177
- Fromme H, Dreyer A, Dietrich S, et al (2015) Neutral polyfluorinated compounds in indoor air in Germany – The LUPE 4 study. *Chemosphere* 139:572–578.
<https://doi.org/10.1016/j.chemosphere.2015.07.024>
- Fromme H, Mosch C, Morovitz M, et al (2010) Pre- and postnatal exposure to perfluorinated compounds (PFCs). *Environmental Science & Technology* 44:7123–7129.
<https://doi.org/10.1021/es101184f>
- Fromme H, Tittlemier SA, Völkel W, et al (2009) Perfluorinated compounds – exposure assessment for the general population in western countries. *International Journal of Hygiene and Environmental Health* 212:239–270. <https://doi.org/10.1016/j.ijheh.2008.04.007>
- Fuertes I, Gómez-Lavín S, Elizalde MP, Urtiaga A (2017) Perfluorinated alkyl substances (PFASs) in northern Spain municipal solid waste landfill leachates. *Chemosphere* 168:399–407.
<https://doi.org/10.1016/j.chemosphere.2016.10.072>
- GAIA and Changing Markets (2017) The CAREless carpet industry: A critique of the carpet stewardship program’s reliance on incineration. A report published by the Global Alliance for Incinerator Alternatives (GAIA) and Changing Markets. April 2017. Available at: <http://www.no-burn.org/wp-content/uploads/FINAL-CARPET-INCINERATION-REPORT-.pdf>. Accessed 24 Sep 2019
- Gallen C, Drage D, Eaglesham G, et al (2017) Australia-wide assessment of perfluoroalkyl substances (PFASs) in landfill leachates. *Journal of Hazardous Materials* 331:132–141.
<https://doi.org/10.1016/j.jhazmat.2017.02.006>
- Gallen C, Eaglesham G, Drage D, et al (2018) A mass estimate of perfluoroalkyl substance (PFAS) release from Australian wastewater treatment plants. *Chemosphere* 208:975–983.
<https://doi.org/10.1016/j.chemosphere.2018.06.024>
- Gannon SA, Johnson T, Nabb DL, et al (2011) Absorption, distribution, metabolism, and excretion of [1-¹⁴C]-perfluorohexanoate ([¹⁴C]-PFHx) in rats and mice. *Toxicology* 283:55–62.
<https://doi.org/10.1016/j.tox.2011.02.004>
- Gao K, Fu J, Xue Q, et al (2018) An integrated method for simultaneously determining 10 classes of per- and polyfluoroalkyl substances in one drop of human serum. *Analytica Chimica Acta* 999:76–86.
<https://doi.org/10.1016/j.aca.2017.10.038>
- Gebbink WA, Bossi R, Rigét FF, et al (2016) Observation of emerging per- and polyfluoroalkyl substances (PFASs) in Greenland marine mammals. *Chemosphere* 144:2384–2391.
<https://doi.org/10.1016/j.chemosphere.2015.10.116>

- Gellrich V, Brunn H, Stahl T (2013) Perfluoroalkyl and polyfluoroalkyl substances (PFASs) in mineral water and tap water. *Journal of Environmental Science and Health, Part A* 48:129–135. <https://doi.org/10.1080/10934529.2013.719431>
- Giesy JP, Kannan K (2001) Global distribution of perfluorooctane sulfonate in wildlife. *Environmental Science & Technology* 35:1339–1342. <https://doi.org/10.1021/es001834k>
- Giesy JP, Naile JE, Khim JS, et al (2010) Aquatic toxicology of perfluorinated chemicals. *Reviews of Environmental Contamination and Toxicology* 202:1–52. https://doi.org/10.1007/978-1-4419-1157-5_1
- Glynn A, Berger U, Bignert A, et al (2012) Perfluorinated alkyl acids in blood serum from primiparous women in Sweden: serial sampling during pregnancy and nursing, and temporal trends 1996–2010. *Environmental Science & Technology* 46:9071–9079. <https://doi.org/10.1021/es301168c>
- Gobelius L (2016) Uptake of per- and polyfluoroalkyl substances by plants. Degree project, University of Agricultural Sciences, Uppsala, Sweden and Hohenheim, Germany. 74
- Gobelius L, Lewis J, Ahrens L (2017) Plant uptake of per- and polyfluoroalkyl substances at a contaminated fire training facility to evaluate the phytoremediation potential of various plant species. *Environmental Science & Technology* 51:12602–12610. <https://doi.org/10.1021/acs.est.7b02926>
- Goldenman G (2017) Study for the strategy for a non-toxic environment of the 7th EAP. Sub-study d: very persistent chemicals
- Gomis MI, Vestergren R, Borg D, Cousins IT (2018) Comparing the toxic potency in vivo of long-chain perfluoroalkyl acids and fluorinated alternatives. *Environment International* 113:1–9. <https://doi.org/10.1016/j.envint.2018.01.011>
- Gomis MI, Wang Z, Scheringer M, Cousins IT (2015) A modeling assessment of the physicochemical properties and environmental fate of emerging and novel per- and polyfluoroalkyl substances. *Science of The Total Environment* 505:981–991. <https://doi.org/10.1016/j.scitotenv.2014.10.062>
- González-Gaya B, Casal P, Jurado E, et al (2019) Vertical transport and sinks of perfluoroalkyl substances in the global open ocean. *Environmental Science: Processes & Impacts*. <https://doi.org/10.1039/C9EM00266A>
- González-Gaya B, Dachs J, Roscales JL, et al (2014) Perfluoroalkylated substances in the global tropical and subtropical surface oceans. *Environmental Science & Technology* 48:13076–13084. <https://doi.org/10.1021/es503490z>

- Goosey E, Harrad S (2011) Perfluoroalkyl compounds in dust from Asian, Australian, European, and North American homes and UK cars, classrooms, and offices. *Environment International* 37:86–92. <https://doi.org/10.1016/j.envint.2010.08.001>
- Gorochategui E, Pérez-Albaladejo E, Casas J, et al (2014) Perfluorinated chemicals: differential toxicity, inhibition of aromatase activity and alteration of cellular lipids in human placental cells. *Toxicology and Applied Pharmacology* 277:124–130. <https://doi.org/10.1016/j.taap.2014.03.012>
- Government of South Australia (2018) South Australia Environment Protection (Water Quality) Amendment Policy 2018. Available at: [https://www.legislation.sa.gov.au/LZ/V/PO/2018/ENVIRONMENT%20PROTECTION%20\(WATER%20QUALITY\)%20AMENDMENT%20POLICY%202018_30.1.2018%20P%20521/30.1.2018%20P%20521.UN.PDF](https://www.legislation.sa.gov.au/LZ/V/PO/2018/ENVIRONMENT%20PROTECTION%20(WATER%20QUALITY)%20AMENDMENT%20POLICY%202018_30.1.2018%20P%20521/30.1.2018%20P%20521.UN.PDF). Accessed 7 Feb 2018
- Grandjean P, Andersen EW, Budtz-Jørgensen E, et al (2012) Serum vaccine antibody concentrations in children exposed to perfluorinated compounds. *JAMA* 307:. <https://doi.org/10.1001/jama.2011.2034>
- Grandjean P, Clapp R (2014) Changing interpretation of human health risks from perfluorinated compounds. *Public Health Reports* 129:482–485. <https://doi.org/10.1177/003335491412900605>
- Grandjean P, Clapp R (2015) Perfluorinated alkyl substances: emerging insights into health risks. *New Solutions* 25:147–163. <https://doi.org/10.1177/1048291115590506>
- Grandjean P, Heilmann C, Weihe P, et al (2016) Serum vaccine antibody concentrations in adolescents exposed to perfluorinated compounds. *Environmental Health Perspectives* 125:077018. <https://doi.org/10.1289/EHP275>
- Grandjean P, Heilmann C, Weihe P, et al (2017) Estimated exposures to perfluorinated compounds in infancy predict attenuated vaccine antibody concentrations at age 5-years. *Journal of Immunotoxicology* 14:188–195. <https://doi.org/10.1080/1547691X.2017.1360968>
- Greaves AK, Letcher RJ, Sonne C, et al (2012) Tissue-specific concentrations and patterns of perfluoroalkyl carboxylates and sulfonates in East Greenland polar bears. *Environmental Science & Technology* 46:11575–11583. <https://doi.org/10.1021/es303400f>
- Greaves AK, Letcher RJ, Sonne C, Dietz R (2013) Brain region distribution and patterns of bioaccumulative perfluoroalkyl carboxylates and sulfonates in East Greenland polar bears (*Ursus maritimus*). *Environmental Toxicology and Chemistry* 32:713–722. <https://doi.org/10.1002/etc.2107>

- Guelfo JL, Adamson DT (2018) Evaluation of a national data set for insights into sources, composition, and concentrations of per- and polyfluoroalkyl substances (PFASs) in U.S. drinking water. *Environmental Pollution* 236:505–513. <https://doi.org/10.1016/j.envpol.2018.01.066>
- Guerra P, Kim M, Kinsman L, et al (2014) Parameters affecting the formation of perfluoroalkyl acids during wastewater treatment. *Journal of Hazardous Materials* 272:148–154. <https://doi.org/10.1016/j.jhazmat.2014.03.016>
- Guo C, Zhang Y, Zhao X, et al (2015) Distribution, source characterization and inventory of perfluoroalkyl substances in Taihu Lake, China. *Chemosphere* 127:201–207. <https://doi.org/10.1016/j.chemosphere.2015.01.053>
- Guo R, Megson D, Myers AL, et al (2016) Application of a comprehensive extraction technique for the determination of poly- and perfluoroalkyl substances (PFASs) in Great Lakes Region sediments. *Chemosphere* 164:535–546. <https://doi.org/10.1016/j.chemosphere.2016.08.126>
- Guo R, Sim W-J, Lee E-S, et al (2010) Evaluation of the fate of perfluoroalkyl compounds in wastewater treatment plants. *Water Research* 44:3476–3486. <https://doi.org/10.1016/j.watres.2010.03.028>
- Guyton KZ, Chiu WA, Bateson TF, et al (2009) A reexamination of the PPAR-alpha activation mode of action as a basis for assessing human cancer risks of environmental contaminants. *Environmental Health Perspectives* 117:1664–1672. <https://doi.org/10.1289/ehp.0900758>
- Hagenaars A, Vergauwen L, De Coen W, Knapen D (2011) Structure–activity relationship assessment of four perfluorinated chemicals using a prolonged zebrafish early life stage test. *Chemosphere* 82:764–772. <https://doi.org/10.1016/j.chemosphere.2010.10.076>
- Hamid H, Li L (2016) Role of wastewater treatment plant (WWTP) in environmental cycling of poly- and perfluoroalkyl (PFAS) compounds. *Ecocycles* 2:. <https://doi.org/10.19040/ecocycles.v2i2.62>
- Hamid H, Li LY, Grace JR (2018) Review of the fate and transformation of per- and polyfluoroalkyl substances (PFASs) in landfills. *Environmental Pollution* 235:74–84. <https://doi.org/10.1016/j.envpol.2017.12.030>
- Harrad S, de Wit CA, Abdallah MA, et al (2010) Indoor contamination with hexabromocyclododecanes, polybrominated diphenyl ethers, and perfluoroalkyl compounds: an important exposure pathway for people? *Environmental Science & Technology* 44:3221–3231. <https://doi.org/10.1021/es903476t>
- Harris MH, Rifas-Shiman SL, Calafat AM, et al (2017) Predictors of per- and polyfluoroalkyl substance (PFAS) plasma concentrations in 6-10 year old American children. *Environmental Science & Technology* 51:5193–5204. <https://doi.org/10.1021/acs.est.6b05811>

- Hart K, Kannan K, Tao L, et al (2008) Skipjack tuna as a bioindicator of contamination by perfluorinated compounds in the oceans. *Science of The Total Environment* 403:215–221. <https://doi.org/10.1016/j.scitotenv.2008.05.035>
- Hartmann C, Raffesberg W, Scharf S, Uhl M (2017) Perfluoroalkylated substances in human urine: results of a biomonitoring pilot study. *Biomonitoring* 4:1–10. <https://doi.org/10.1515/bimo-2017-0001>
- Haug LS, Huber S, Schlabach M, et al (2011) Investigation on per- and polyfluorinated compounds in paired samples of house dust and indoor air from Norwegian homes. *Environmental Science & Technology* 45:7991–7998. <https://doi.org/10.1021/es103456h>
- Haug LS, Salihovic S, Jogsten IE, et al (2010) Levels in food and beverages and daily intake of perfluorinated compounds in Norway. *Chemosphere* 80:1137–1143. <https://doi.org/10.1016/j.chemosphere.2010.06.023>
- HBN (2017) Healthy Building Network (HBN): Eliminating toxics in carpet: Lessons for the future of recycling. An optimizing recycling report by the Healthy Building Network with support from Changing Markets and the Global Alliance for Incinerator Alternatives, October 2017. Available at: <https://healthybuilding.net/uploads/files/eliminating-toxics-in-carpet-lessons-for-the-future-of-recycling.pdf>. Accessed 16 Apr 2018
- He X, Liu Y, Xu B, et al (2018) PFOA is associated with diabetes and metabolic alteration in US men: National Health and Nutrition Examination Survey 2003–2012. *Science of The Total Environment* 625:566–574. <https://doi.org/10.1016/j.scitotenv.2017.12.186>
- Hebert PC, MacManus-Spencer LA (2010) Development of a fluorescence model for the binding of medium- to long-chain perfluoroalkyl acids to human serum albumin through a mechanistic evaluation of spectroscopic evidence. *Analytical Chemistry* 82:6463–6471. <https://doi.org/10.1021/ac100721e>
- Henry BJ, Carlin JP, Hammerschmidt JA, et al (2018) A critical review of the application of polymer of low concern and regulatory criteria to fluoropolymers: fluoropolymers PLC. *Integrated Environmental Assessment and Management* 14:316–334. <https://doi.org/10.1002/ieam.4035>
- Hepburn E, Madden C, Szabo D, et al (2019) Contamination of groundwater with per- and polyfluoroalkyl substances (PFAS) from legacy landfills in an urban re-development precinct. *Environmental Pollution* 248:101–113. <https://doi.org/10.1016/j.envpol.2019.02.018>
- Herzke D, Olsson E, Posner S (2012) Perfluoroalkyl and polyfluoroalkyl substances (PFASs) in consumer products in Norway – A pilot study. *Chemosphere* 88:980–987. <https://doi.org/10.1016/j.chemosphere.2012.03.035>

- Heydebreck F, Tang J, Xie Z, Ebinghaus R (2016) Emissions of per- and polyfluoroalkyl substances in a textile manufacturing plant in China and their relevance for workers' exposure. *Environmental Science & Technology* 50:10386–10396. <https://doi.org/10.1021/acs.est.6b03213>
- Higgins CP, Luthy RG (2006) Sorption of perfluorinated surfactants on sediments. *Environmental Science & Technology* 40:7251–7256. <https://doi.org/10.1021/es061000n>
- Hoffman K, Webster TF, Bartell SM, et al (2011) Private drinking water wells as a source of exposure to perfluorooctanoic acid (PFOA) in communities surrounding a fluoropolymer production facility. *Environmental Health Perspectives* 119:92–97. <https://doi.org/10.1289/ehp.1002503>
- Hoke RA, Bouchelle LD, Ferrell BD, Buck RC (2012) Comparative acute freshwater hazard assessment and preliminary PNEC development for eight fluorinated acids. *Chemosphere* 87:725–733. <https://doi.org/10.1016/j.chemosphere.2011.12.066>
- Home Depot (2019) Phasing Out Products Containing PFAS. Available at: <https://corporate.homedepot.com/newsroom/phasing-out-products-containing-pfas>. Accessed 24 Sep 2019
- Hong S, Khim JS, Wang T, et al (2015) Bioaccumulation characteristics of perfluoroalkyl acids (PFAAs) in coastal organisms from the west coast of South Korea. *Chemosphere* 129:157–163. <https://doi.org/10.1016/j.chemosphere.2014.06.023>
- Hoover G, Kar S, Guffey S, et al (2019) In vitro and in silico modeling of perfluoroalkyl substances mixture toxicity in an amphibian fibroblast cell line. *Chemosphere* 233:25–33. <https://doi.org/10.1016/j.chemosphere.2019.05.065>
- Horst J, McDonough J, Ross I, et al (2018) Water treatment technologies for PFAS: the next generation. *Groundwater Monitoring & Remediation* 38:13–23. <https://doi.org/10.1111/gwmr.12281>
- Houde M, Czub G, Small JM, et al (2008) Fractionation and bioaccumulation of perfluorooctane sulfonate (PFOS) isomers in a Lake Ontario food web. *Environmental Science & Technology* 42:9397–9403. <https://doi.org/10.1021/es800906r>
- Houde M, De Silva AO, Muir DCG, Letcher RJ (2011) Monitoring of perfluorinated compounds in aquatic biota: an updated review. *Environmental Science & Technology* 45:7962–7973. <https://doi.org/10.1021/es104326w>
- Houde M, Martin JW, Letcher RJ, et al (2006) Biological monitoring of polyfluoroalkyl substances: a review. *Environmental Science & Technology* 40:3463–3473. <https://doi.org/10.1021/es052580b>
- Houde M, Wells RS, Fair PA, et al (2005) Polyfluoroalkyl compounds in free-ranging bottlenose dolphins (*Tursiops truncatus*) from the Gulf of Mexico and the Atlantic Ocean. *Environmental Science & Technology* 39:6591–6598. <https://doi.org/10.1021/es0506556>

- Houtz EF, Sedlak DL (2012) Oxidative conversion as a means of detecting precursors to perfluoroalkyl acids in urban runoff. *Environmental Science & Technology* 46:9342–9349. <https://doi.org/10.1021/es302274g>
- Houtz EF, Sutton R, Park JS, Sedlak M (2016) Poly- and perfluoroalkyl substances in wastewater: significance of unknown precursors, manufacturing shifts, and likely AFFF impacts. *Water Research* 95:142–149. <https://doi.org/10.1016/j.watres.2016.02.055>
- Høyer BB, Ramlau-Hansen CH, Vrijheid M, et al (2015) Anthropometry in 5- to 9-year-old Greenlandic and Ukrainian children in relation to prenatal exposure to perfluorinated alkyl substances. *Environmental Health Perspectives* 123:841–846. <https://doi.org/10.1289/ehp.1408881>
- Hu XC, Andrews DQ, Lindstrom AB, et al (2016) Detection of poly- and perfluoroalkyl substances (PFASs) in U.S. drinking water linked to industrial sites, military fire training areas, and wastewater treatment plants. *Environmental Science & Technology Letters* 3:344–350. <https://doi.org/10.1021/acs.estlett.6b00260>
- Huber S, Moe MK, Schmidbauer N, et al (2009) Emissions from incineration of fluoropolymer materials: a literature survey
- Hurley S, Goldberg D, Wang M, et al (2018) Time trends in per- and polyfluoroalkyl substances (PFASs) in California women: declining serum levels, 2011–2015. *Environmental Science & Technology* 52:277–287. <https://doi.org/10.1021/acs.est.7b04650>
- Hurley S, Houtz E, Goldberg D, et al (2016) Preliminary associations between the detection of perfluoroalkyl acids (PFAAs) in drinking water and serum concentrations in a sample of California women. *Environmental Science & Technology Letters* 3:264–269. <https://doi.org/10.1021/acs.estlett.6b00154>
- Huset CA (2007) Determination of fluorochemicals in waste-dominated aqueous systems. Ph.D thesis. May 2007, Oregon State University
- Huset CA, Barlaz MA, Barofsky DF, Field JA (2011) Quantitative determination of fluorochemicals in municipal landfill leachates. *Chemosphere* 82:1380–1386. <https://doi.org/10.1016/j.chemosphere.2010.11.072>
- Invista North America (2017) Water repellent, soil resistant, fluorine-free compositions. United States Patent application, Pub No.: US 20170030010 A1. Available at: <https://patentimages.storage.googleapis.com/0f/bb/ce/9582bc9072b90c/US20170030010A1.pdf>. Accessed 9 Nov 2017
- IPCC (2007) Intergovernmental Panel on Climate Change (IPCC): Working Group I: The physical science basis, 2.10.2 Direct global warming potentials. IPCC Fourth Assessment Report: Climate Change 2007. Available at: https://www.ipcc.ch/publications_and_data/ar4/wg1/en/ch2s2-10-2.html. Accessed 23 Aug 2017

- Ishibashi H, Yamauchi R, Matsuoka M, et al (2008) Fluorotelomer alcohols induce hepatic vitellogenin through activation of the estrogen receptor in male medaka (*Oryzias latipes*). *Chemosphere* 71:1853–1859. <https://doi.org/10.1016/j.chemosphere.2008.01.065>
- ITRC (2018) Interstate Technology and Regulatory Council (ITRC): PFAS fact sheet series-Environmental Fate and Transport for Per- and Polyfluoroalkyl Substances (Published March 16, 2018). Available at: https://pfas-1.itrcweb.org/wp-content/uploads/2018/03/pfas_fact_sheet_fate_and_transport__3_16_18.pdf. Accessed 23 Sep 2019
- Jahnke A, Ahrens L, Ebinghaus R, Temme C (2007) Urban versus remote air concentrations of fluorotelomer alcohols and other polyfluorinated alkyl substances in Germany. *Environmental Science & Technology* 41:745–752. <https://doi.org/10.1021/es0619861>
- Jain RB (2018) Contribution of diet and other factors to the observed levels of selected perfluoroalkyl acids in serum among US children aged 3-11 years. *Environmental Research* 161:268–275. <https://doi.org/10.1016/j.envres.2017.11.018>
- Jamieson AJ, Malkocs T, Piertney SB, et al (2017) Bioaccumulation of persistent organic pollutants in the deepest ocean fauna. *Nature Ecology & Evolution* 1:51. <https://doi.org/10.1038/s41559-016-0051>
- Jensen AA, Poulsen PB, Bossi R (2008) Survey and environmental/health assessment of fluorinated substances in impregnated consumer products and impregnating agents. *Survey of Chemical Substances in Consumer Products*, No. 99
- Ji W, Xiao L, Ling Y, et al (2018) Removal of GenX and perfluorinated alkyl substances from water by amine-functionalized covalent organic frameworks. *Journal of the American Chemical Society* 140:12677–12681. <https://doi.org/10.1021/jacs.8b06958>
- Jian J-M, Guo Y, Zeng L, et al (2017) Global distribution of perfluorochemicals (PFCs) in potential human exposure source—A review. *Environment International* 108:51–62. <https://doi.org/10.1016/j.envint.2017.07.024>
- Jones PD, Hu W, De Coen W, et al (2003) Binding of perfluorinated fatty acids to serum proteins. *Environmental Toxicology and Chemistry* 22:2639. <https://doi.org/10.1897/02-553>
- Joudan S, Yeung LWY, Mabury SA (2017) Biological cleavage of the C-P bond in perfluoroalkyl phosphinic acids in male Sprague-Dawley rats and the formation of persistent and reactive metabolites. *Environmental Health Perspectives* 125:117001. <https://doi.org/10.1289/EHP1841>
- Jurries A (2016) Columbia Goes PFC-Free. *TheGearCaster*, June 20, 2016. Available at: <http://www.thegearcaster.com/2016/06/columbia-goes-pfc-free.html>. Accessed 5 Jan 2018

- Kabadi SV, Fisher J, Aungst J, Rice P (2018) Internal exposure-based pharmacokinetic evaluation of potential for biopersistence of 6:2 fluorotelomer alcohol (FTOH) and its metabolites. *Food and Chemical Toxicology* 112:375–382. <https://doi.org/10.1016/j.fct.2018.01.012>
- Kaboré HA, Vo Duy S, Munoz G (2018) Worldwide drinking water occurrence and levels of newly-identified perfluoroalkyl and polyfluoroalkyl substances. *Science of The Total Environment* 616–617:1089–1100. <https://doi.org/10.1016/j.scitotenv.2017.10.210>
- Kang H, Choi K, Lee HS, et al (2016) Elevated levels of short carbon-chain PFCAs in breast milk among Korean women: current status and potential challenges. *Environmental Research* 148:351–359. <https://doi.org/10.1016/j.envres.2016.04.017>
- Kannan K, Perrotta E, Thomas NJ (2006) Association between perfluorinated compounds and pathological conditions in southern sea otters. *Environmental Science & Technology* 40:4943–4948. <https://doi.org/10.1021/es060932o>
- Karášková P, Venier M, Melymuk L, et al (2016) Perfluorinated alkyl substances (PFASs) in household dust in Central Europe and North America. *Environment International* 94:315–324. <https://doi.org/10.1016/j.envint.2016.05.031>
- Karnjanapiboonwong A, Deb SK, Subbiah S, et al (2018) Perfluoroalkylsulfonic and carboxylic acids in earthworms (*Eisenia fetida*): accumulation and effects results from spiked soils at PFAS concentrations bracketing environmental relevance. *Chemosphere* 199:168–173. <https://doi.org/10.1016/j.chemosphere.2018.02.027>
- Kärman A, Ericson I, van Bavel B, et al (2007) Exposure of perfluorinated chemicals through lactation: levels of matched human milk and serum and a temporal trend, 1996-2004, in Sweden. *Environmental Health Perspectives* 115:226–230. <https://doi.org/10.1289/ehp.9491>
- Keller JM, Kannan K, Taniyasu S, et al (2005) Perfluorinated compounds in the plasma of loggerhead and Kemp's ridley sea turtles from the southeastern coast of the United States. *Environmental Science & Technology* 39:9101–9108. <https://doi.org/10.1021/es050690c>
- Keller JM, Ngai L, McNeill JB, et al (2012) Perfluoroalkyl contaminants in plasma of five sea turtle species: comparisons in concentration and potential health risks. *Environmental Toxicology and Chemistry* 31:1223–1230. <https://doi.org/10.1002/etc.1818>
- Kelly BC, Ikonomidou MG, Blair JD, et al (2009) Perfluoroalkyl contaminants in an Arctic marine food web: trophic magnification and wildlife exposure. *Environmental Science & Technology* 43:4037–4043. <https://doi.org/10.1021/es9003894>
- KEMI (2015) Swedish Chemicals Agency (KEMI): Occurrence and use of highly fluorinated substances and alternatives. Report from a government assignment. Report 7/15. Stockholm, Sweden

- Kildee DT, Rochester LB, Boyle BF, et al. (2017) Letter to President Donald J. Trump, August 1, 2017. Available at: <https://dankildee.house.gov/sites/danielkildee.house.gov/files/8-1-17%20-%20PFC%20Contamination%20at%20Military%20Bases%20%28Trump%29.pdf>. Accessed 8 Sep 2017
- Kim DH, Lee MY, Oh JE (2014a) Perfluorinated compounds in serum and urine samples from children aged 5–13 years in South Korea. *Environmental Pollution* 192:171–178. <https://doi.org/10.1016/j.envpol.2014.05.024>
- Kim Lazcano R, Perre C, Mashtare ML, Lee LS (2019) Per- and polyfluoroalkyl substances in commercially available biosolid-based products: the effect of treatment processes. *Water Environment Research*. <https://doi.org/10.1002/wer.1174>
- Kim M, Li LY, Grace JR, et al (2015) Compositional effects on leaching of stain-guarded (perfluoroalkyl and polyfluoroalkyl substance-treated) carpet in landfill leachate. *Environmental Science & Technology* 49:6564–6573. <https://doi.org/10.1021/es505333y>
- Kim MH, Wang N, Chu KH (2014b) 6:2 Fluorotelomer alcohol (6:2 FTOH) biodegradation by multiple microbial species under different physiological conditions. *Applied Microbiology and Biotechnology* 98:1831–1840. <https://doi.org/10.1007/s00253-013-5131-3>
- Kim MH, Wang N, McDonald T, Chu KH (2012a) Biodefluorination and biotransformation of fluorotelomer alcohols by two alkane-degrading *Pseudomonas* strains. *Biotechnology and Bioengineering* 109:3041–3048. <https://doi.org/10.1002/bit.24561>
- Kim S, Choi K, Ji K, et al (2011) Trans-placental transfer of thirteen perfluorinated compounds and relations with fetal thyroid hormones. *Environmental Science & Technology* 45:7465–7472. <https://doi.org/10.1021/es202408a>
- Kim SK, Im JK, Kang YM, et al (2012b) Wastewater treatment plants (WWTPs)-derived national discharge loads of perfluorinated compounds (PFCs). *Journal of Hazardous Materials* 201–202:82–91. <https://doi.org/10.1016/j.jhazmat.2011.11.036>
- Kim SK, Kannan K (2007) Perfluorinated acids in air, rain, snow, surface runoff, and lakes: relative importance of pathways to contamination of urban lakes. *Environmental Science & Technology* 41:8328–8334. <https://doi.org/10.1021/es072107t>
- Kiplinger JL, Richmond TG, Osterberg CE (1994) Activation of carbon-fluorine bonds by metal complexes. *Chemical Reviews* 94:373–431. <https://doi.org/10.1021/cr00026a005>
- Kirchgeorg T, Dreyer A, Gabrielli P, et al (2016) Seasonal accumulation of persistent organic pollutants on a high altitude glacier in the Eastern Alps. *Environmental Pollution* 218:804–812. <https://doi.org/10.1016/j.envpol.2016.08.004>

- Klepeis NE, Nelson WC, Ott WR, et al (2001) The National Human Activity Pattern Survey (NHAPS): a resource for assessing exposure to environmental pollutants. *Journal of Exposure Science & Environmental Epidemiology* 11:231–252. <https://doi.org/10.1038/sj.jea.7500165>
- Knepper TP, Frömel T, Gremmel C, et al (2014) Understanding the exposure pathways of per- and polyfluoroalkyl substances (PFASs) via use of PFASs-containing products - risk estimation for man and environment. Federal Environment Agency (Umweltbundesamt), Germany
- Knobeloch L, Imm P, Anderson H (2012) Perfluoroalkyl chemicals in vacuum cleaner dust from 39 Wisconsin homes. *Chemosphere* 88:779–783. <https://doi.org/10.1016/j.chemosphere.2012.03.082>
- Knutsen H, Mæhlum T, Haarstad K, et al (2019) Leachate emissions of short- and long-chain per- and polyfluoroalkyl substances (PFASs) from various Norwegian landfills. *Environmental Science: Processes & Impacts*. <https://doi.org/10.1039/C9EM00170K>
- Knutsen HK, Alexander J, Barregård L, et al (2018) Risk to human health related to the presence of perfluorooctane sulfonic acid and perfluorooctanoic acid in food. *EFSA Journal* 16:. <https://doi.org/10.2903/j.efsa.2018.5194>
- Konwick BJ, Tomy GT, Ismail N, et al (2008) Concentrations and patterns of perfluoroalkyl acids in Georgia, USA surface waters near and distant to a major use source. *Environmental Toxicology and Chemistry* 27:2011. <https://doi.org/10.1897/07-659.1>
- Kotthoff M, Müller J, Jüriling H, et al (2015) Perfluoroalkyl and polyfluoroalkyl substances in consumer products. *Environmental Science and Pollution Research* 22:14546–14559. <https://doi.org/10.1007/s11356-015-4202-7>
- Krafft MP, Riess JG (2015a) Selected physicochemical aspects of poly- and perfluoroalkylated substances relevant to performance, environment and sustainability—Part one. *Chemosphere* 129:4–19. <https://doi.org/10.1016/j.chemosphere.2014.08.039>
- Krafft MP, Riess JG (2015b) Per- and polyfluorinated substances (PFASs): environmental challenges. *Current Opinion in Colloid & Interface Science* 20:192–212. <https://doi.org/10.1016/j.cocis.2015.07.004>
- Kumar SK, Zushi Y, Masunaga S, et al (2009) Perfluorinated organic contaminants in sediment and aquatic wildlife, including sharks, from Georgia, USA. *Marine Pollution Bulletin* 58:621–629. <https://doi.org/10.1016/j.marpolbul.2008.12.006>
- Ladics GS, Kennedy GL, O'Connor J, et al (2008) 90-day oral gavage toxicity study of 8-2 fluorotelomer alcohol in rats. *Drug and Chemical Toxicology* 31:189–216. <https://doi.org/10.1080/01480540701873103>

- Lam JCW, Lyu J, Kwok KY, Lam PKS (2016) Perfluoroalkyl substances (PFASs) in marine mammals from the South China Sea and their temporal changes 2002-2014: concern for alternatives of PFOS? *Environmental Science & Technology* 50:6728–6736. <https://doi.org/10.1021/acs.est.5b06076>
- Land M, de Wit CA, Bignert A, et al (2018) What is the effect of phasing out long-chain per- and polyfluoroalkyl substances on the concentrations of perfluoroalkyl acids and their precursors in the environment? A systematic review. *Environmental Evidence* 7:.
<https://doi.org/10.1186/s13750-017-0114-y>
- Lang JR, Allred BM, Field JA, et al (2017) National estimate of per- and polyfluoroalkyl substance (PFAS) release to U.S. municipal landfill leachate. *Environmental Science & Technology* 51:2197–2205.
<https://doi.org/10.1021/acs.est.6b05005>
- Lang JR, Allred BM, Peaslee GF, et al (2016) Release of per- and polyfluoroalkyl substances (PFASs) from carpet and clothing in model anaerobic landfill reactors. *Environmental Science & Technology* 50:5024–5032. <https://doi.org/10.1021/acs.est.5b06237>
- Langer V, Dreyer A, Ebinghaus R (2010) Polyfluorinated compounds in residential and nonresidential indoor air. *Environmental Science & Technology* 44:8075–8081.
<https://doi.org/10.1021/es102384z>
- Latala A, Nędzi M, Stepnowski P (2009) Acute toxicity assessment of perfluorinated carboxylic acids towards the Baltic microalgae. *Environmental Toxicology and Pharmacology* 28:167–171.
<https://doi.org/10.1016/j.etap.2009.03.010>
- Lechner M, Knapp H (2011) Carryover of perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS) from soil to plant and distribution to the different plant compartments studied in cultures of carrots (*Daucus carota* ssp. *Sativus*), potatoes (*Solanum tuberosum*), and cucumbers (*Cucumis Sativus*). *Journal of Agricultural and Food Chemistry* 59:11011–11018.
<https://doi.org/10.1021/jf201355y>
- Lee H, D'eon J, Mabury SA (2010) Biodegradation of polyfluoroalkyl phosphates as a source of perfluorinated acids to the environment. *Environmental Science & Technology* 44:3305–3310.
<https://doi.org/10.1021/es9028183>
- Lee H, Tevlin AG, Mabury SA, Mabury SA (2014) Fate of polyfluoroalkyl phosphate diesters and their metabolites in biosolids-applied soil: Biodegradation and plant uptake in greenhouse and field experiments. *Environmental Science & Technology* 48:340–349.
<https://doi.org/10.1021/es403949z>
- Lee S, Kim S, Park J, et al (2018) Perfluoroalkyl substances (PFASs) in breast milk from Korea: time-course trends, influencing factors, and infant exposure. *Science of The Total Environment* 612:286–292. <https://doi.org/10.1016/j.scitotenv.2017.08.094>

- Lee YJ, Kim MK, Bae J, Yang JH (2013) Concentrations of perfluoroalkyl compounds in maternal and umbilical cord sera and birth outcomes in Korea. *Chemosphere* 90:1603–1609. <https://doi.org/10.1016/j.chemosphere.2012.08.035>
- Lemal DM (2004) Perspective on fluorocarbon chemistry. *The Journal of Organic Chemistry* 69:1–11. <https://doi.org/10.1021/jo0302556>
- Li F, Duan J, Tian S, et al (2020) Short-chain per- and polyfluoroalkyl substances in aquatic systems: occurrence, impacts and treatment. *Chemical Engineering Journal* 380:
- Li L, Liu J, Hu J, Wania F (2017) Degradation of fluorotelomer-based polymers contributes to the global occurrence of fluorotelomer alcohol and perfluoroalkyl carboxylates: a combined dynamic substance flow and environmental fate modeling analysis. *Environmental Science & Technology* 51:4461–4470. <https://doi.org/10.1021/acs.est.6b04021>
- Li Y, Oliver DP, Kookana RS (2018) A critical analysis of published data to discern the role of soil and sediment properties in determining sorption of per and polyfluoroalkyl substances (PFASs). *Science of The Total Environment* 628–629:110–120. <https://doi.org/10.1016/j.scitotenv.2018.01.167>
- Li Z, Guo L (2016) Environmental occurrences and toxicology of fluorotelomer alcohols (in Chinese with English abstract). *Progress in Chemistry* 28:993–1005
- Lieder PH, Chang SH, York RG, Butenhoff JL (2009) Toxicological evaluation of potassium perfluorobutanesulfonate in a 90-day oral gavage study with Sprague-Dawley rats. *Toxicology* 255:45–52. <https://doi.org/10.1016/j.tox.2008.10.002>
- Lien GW, Huang CC, Wu KH, et al (2013) Neonatal–maternal factors and perfluoroalkyl substances in cord blood. *Chemosphere* 92:843–850. <https://doi.org/10.1016/j.chemosphere.2013.04.038>
- Lilienthal H, Dieter HH, Hölzer J, Wilhelm M (2017) Recent experimental results of effects of perfluoroalkyl substances in laboratory animals – Relation to current regulations and guidance values. *International Journal of Hygiene and Environmental Health* 220:766–775. <https://doi.org/10.1016/j.ijheh.2017.03.001>
- Lim TC, Wang B, Huang J, et al (2011) Emission inventory for PFOS in China: review of past methodologies and suggestions. *Scientific World Journal* 11:1963–1980. <https://doi.org/10.1100/2011/868156>
- Lin AY., Panchangam SC, Tsai YT, Yu TH (2014a) Occurrence of perfluorinated compounds in the aquatic environment as found in science park effluent, river water, rainwater, sediments, and biotissues. *Environ Monit Assess* 186:3265–3275. <https://doi.org/10.1007/s10661-014-3617-9>

- Lin J, Pan D, Davis SJ, et al (2014b) China's international trade and air pollution in the United States. *Proceedings of the National Academy of Sciences* 111:1736–1741. <https://doi.org/10.1073/pnas.1312860111>
- Lindstrom AB, Strynar MJ, Delinsky AD, et al (2011a) Application of WWTP biosolids and resulting perfluorinated compound contamination of surface and well water in Decatur, Alabama, USA. *Environmental Science & Technology* 45:8015–8021. <https://doi.org/10.1021/es1039425>
- Lindstrom AB, Strynar MJ, Libelo EL (2011b) Polyfluorinated compounds: past, present, and future. *Environmental Science & Technology* 45:7954–7961. <https://doi.org/10.1021/es2011622>
- Liu B, Zhang H, Yao D, et al (2015a) Perfluorinated compounds (PFCs) in the atmosphere of Shenzhen, China: spatial distribution, sources and health risk assessment. *Chemosphere* 138:511–518. <https://doi.org/10.1016/j.chemosphere.2015.07.012>
- Liu C, Gin KY, Chang VW, et al (2011a) Novel perspectives on the bioaccumulation of PFCs – the concentration dependency. *Environmental Science & Technology* 45:9758–9764. <https://doi.org/10.1021/es202078n>
- Liu H, Chen Q, Lei L, et al (2018a) Prenatal exposure to perfluoroalkyl and polyfluoroalkyl substances affects leukocyte telomere length in female newborns. *Environmental Pollution* 235:446–452. <https://doi.org/10.1016/j.envpol.2017.12.095>
- Liu H, Sheng N, Zhang W, Dai J (2015b) Toxic effects of perfluorononanoic acid on the development of Zebrafish (*Danio rerio*) embryos. *Journal of Environmental Sciences* 32:26–34. <https://doi.org/10.1016/j.jes.2014.11.008>
- Liu J, Mejia Avendaño S (2013) Microbial degradation of polyfluoroalkyl chemicals in the environment: a review. *Environment International* 61:98–114. <https://doi.org/10.1016/j.envint.2013.08.022>
- Liu J, Wang N, Buck RC, et al (2010a) Aerobic biodegradation of [14C] 6:2 fluorotelomer alcohol in a flow-through soil incubation system. *Chemosphere* 80:716–723. <https://doi.org/10.1016/j.chemosphere.2010.05.027>
- Liu J, Wang N, Szostek B, et al (2010b) 6-2 Fluorotelomer alcohol aerobic biodegradation in soil and mixed bacterial culture. *Chemosphere* 78:437–444. <https://doi.org/10.1016/j.chemosphere.2009.10.044>
- Liu W, Chen S, Harada KH, Koizumi A (2011b) Analysis of perfluoroalkyl carboxylates in vacuum cleaner dust samples in Japan. *Chemosphere* 85:1734–1741. <https://doi.org/10.1016/j.chemosphere.2011.09.024>
- Liu W, Takahashi S, Sakuramachi Y, et al (2013) Polyfluorinated telomers in indoor air of Japanese houses. *Chemosphere* 90:1672–1677. <https://doi.org/10.1016/j.chemosphere.2012.09.062>

- Liu X, Guo Z, Folk EE, Roache NF (2015c) Determination of fluorotelomer alcohols in selected consumer products and preliminary investigation of their fate in the indoor environment. *Chemosphere* 129:81–86. <https://doi.org/10.1016/j.chemosphere.2014.06.012>
- Liu X, Guo Z, Krebs KA, et al (2014) Concentrations and trends of perfluorinated chemicals in potential indoor sources from 2007 through 2011 in the US. *Chemosphere* 98:51–57. <https://doi.org/10.1016/j.chemosphere.2013.10.001>
- Liu Y, Richardson ES, Derocher AE, et al (2018b) Hundreds of unrecognized halogenated contaminants discovered in polar bear serum. *Angewandte Chemie International Edition* 57:16401–16406. <https://doi.org/10.1002/anie.201809906>
- Liu Y, Ruan T, Lin Y, et al (2017) Chlorinated polyfluoroalkyl ether sulfonic acids in marine organisms from Bohai Sea, China: occurrence, temporal variations, and trophic transfer behavior. *Environmental Science & Technology* 51:4407–4414. <https://doi.org/10.1021/acs.est.6b06593>
- Liu Z, Lu Y, Wang T, et al (2016) Risk assessment and source identification of perfluoroalkyl acids in surface and ground water: spatial distribution around a mega-fluorochemical industrial park, China. *Environment International* 91:69–77. <https://doi.org/10.1016/j.envint.2016.02.020>
- Llorca M, Farré M, Picó Y, et al (2010) Infant exposure of perfluorinated compounds: Levels in breast milk and commercial baby food. *Environment International* 36:584–592. <https://doi.org/10.1016/j.envint.2010.04.016>
- Llorca M, Farré M, Tavano MS, et al (2012) Fate of a broad spectrum of perfluorinated compounds in soils and biota from Tierra del Fuego and Antarctica. *Environmental Pollution* 163:158–166. <https://doi.org/10.1016/j.envpol.2011.10.027>
- Long M, Bossi R, Bonefeld-Jørgensen EC (2012) Level and temporal trend of perfluoroalkyl acids in Greenlandic Inuit. *International Journal of Circumpolar Health* 71:17998. <https://doi.org/10.3402/ijch.v71i0.17998>
- Loveless SE, Slezak B, Serex T, et al (2009) Toxicological evaluation of sodium perfluorohexanoate. *Toxicology* 264:32–44. <https://doi.org/10.1016/j.tox.2009.07.011>
- Mackay D, Fraser A (2000) Bioaccumulation of persistent organic chemicals: mechanisms and models. *Environmental Pollution* 110:375–391. [https://doi.org/10.1016/S0269-7491\(00\)00162-7](https://doi.org/10.1016/S0269-7491(00)00162-7)
- Mackay D, Hughes DM, Romano ML, Bonnell M (2014) The role of persistence in chemical evaluations. *Integrated Environmental Assessment and Management* 10:588–594. <https://doi.org/10.1002/ieam.1545>
- MacLeod M, Breitholtz M, Cousins IT, et al (2014) Identifying chemicals that are planetary boundary threats. *Environmental Science & Technology* 48:11057–11063. <https://doi.org/10.1021/es501893m>

- Makey CM, Webster TF, Martin JW, et al (2017) Airborne precursors predict maternal serum perfluoroalkyl acid concentrations. *Environmental Science & Technology* 51:7667–7675. <https://doi.org/10.1021/acs.est.7b00615>
- Malits J, Blustein J, Trasande L, Attina TM (2017) Perfluorooctanoic acid and low birth weight: estimates of US attributable burden and economic costs from 2003 through 2014. *International Journal of Hygiene and Environmental Health* 221:269–275. <https://doi.org/10.1016/j.ijheh.2017.11.004>
- Mamsen LS, Björvang RD, Mucs D, et al (2019) Concentrations of perfluoroalkyl substances (PFASs) in human embryonic and fetal organs from first, second, and third trimester pregnancies. *Environment International* 124:482–492. <https://doi.org/10.1016/j.envint.2019.01.010>
- Mamsen LS, Jönsson BAG, Lindh CH, et al (2017) Concentration of perfluorinated compounds and cotinine in human foetal organs, placenta, and maternal plasma. *Science of The Total Environment* 596–597:97–105. <https://doi.org/10.1016/j.scitotenv.2017.04.058>
- Martín J, Santos JL, Aparicio I, Alonso E (2019) Exposure assessment to parabens, bisphenol A and perfluoroalkyl compounds in children, women and men by hair analysis. *Science of The Total Environment* 695:133864. <https://doi.org/10.1016/j.scitotenv.2019.133864>
- Martin JW, Mabury SA, Solomon KR, Muir DCG (2003) Bioconcentration and tissue distribution of perfluorinated acids in rainbow trout (*Oncorhynchus mykiss*). *Environmental Toxicology and Chemistry* 22:196–204
- Martin JW, Smithwick M, Braune BM, et al (2004) Identification of long-chain perfluorinated acids in biota from the Canadian Arctic. *Environmental Science & Technology* 38:373–380
- Mason Chemical Company (2017) Specialty Surfactants: Fluorosurfactant - Structure/Function. Available at: <http://archive.li/GthbK>. Accessed 20 Sep 2019
- McLachlan MS, Felizeter S, Klein M, et al (2019) Fate of a perfluoroalkyl acid mixture in an agricultural soil studied in lysimeters. *Chemosphere* 223:180–187. <https://doi.org/10.1016/j.chemosphere.2019.02.012>
- MDH (2017) Minnesota Department of Health (MDH): MDH response to EPA health advisory for PFOS and PFOA. June 2017. Available at: <http://www.health.state.mn.us/divs/eh/hazardous/topics/pfcs/current.html>. Accessed 22 Aug 2017
- Mercier F, Glorennec P, Thomas O, Bot BL (2011) Organic contamination of settled house dust, a review for exposure assessment purposes. *Environmental Science & Technology* 45:6716–6727. <https://doi.org/10.1021/es200925h>

- MHR (2017) Michigan House of Representatives (MHR): House Bill No. 5375. Available at: <http://www.legislature.mi.gov/documents/2017-2018/billintroduced/House/pdf/2017-HIB-5375.pdf>. Accessed 5 Jan 2018
- Midasch O, Drexler H, Hart N, et al (2007) Transplacental exposure of neonates to perfluorooctanesulfonate and perfluorooctanoate: a pilot study. *International Archives of Occupational and Environmental Health* 80:643–648. <https://doi.org/10.1007/s00420-006-0165-9>
- Mintel (2019) Global New Products Database. Available at: www.gnpd.com. Accessed 18 Sep 2019
- Mitro SD, Dodson RE, Singla V, et al (2016) Consumer product chemicals in indoor dust: a quantitative meta-analysis of U.S. studies. *Environmental Science & Technology* 50:10661–10672. <https://doi.org/10.1021/acs.est.6b02023>
- Mogensen UB, Grandjean P, Nielsen F, et al (2015) Breastfeeding as an exposure pathway for perfluorinated alkylates. *Environmental Science & Technology* 49:10466–10473. <https://doi.org/10.1021/acs.est.5b02237>
- Mondal D, Weldon RH, Armstrong BG, et al (2014) Breastfeeding: a potential excretion route for mothers and implications for infant exposure to perfluoroalkyl acids. *Environmental Health Perspectives* 122:187–192. <https://doi.org/10.1289/ehp.1306613>
- Monroy R, Morrison K, Teo K, et al (2008) Serum levels of perfluoroalkyl compounds in human maternal and umbilical cord blood samples. *Environmental Research* 108:56–62. <https://doi.org/10.1016/j.envres.2008.06.001>
- Mora C, Sale P (2011) Ongoing global biodiversity loss and the need to move beyond protected areas: a review of the technical and practical shortcomings of protected areas on land and sea. *Marine Ecology Progress Series* 434:251–266. <https://doi.org/10.3354/meps09214>
- Moschet C, Anumol T, Lew BM, et al (2018) Household dust as a repository of chemical accumulation: new insights from a comprehensive high-resolution mass spectrometric study. *Environmental Science & Technology* 52:2878–2887. <https://doi.org/10.1021/acs.est.7b05767>
- Moya J, Bearer CF, Etzel RA (2004) Children’s behavior and physiology and how it affects exposure to environmental contaminants. *Pediatrics* 113:996–1006
- Mudumbi JBN, Ntwampe SKO, Matsha T, et al (2017) Recent developments in polyfluoroalkyl compounds research: a focus on human/environmental health impact, suggested substitutes and removal strategies. *Environmental Monitoring and Assessment* 189:. <https://doi.org/10.1007/s10661-017-6084-2>

- Muir D, Bossi R, Carlsson P, et al (2019) Levels and trends of poly- and perfluoroalkyl substances in the Arctic environment – An update. *Emerging Contaminants* 5:240–271. <https://doi.org/10.1016/j.emcon.2019.06.002>
- Müller CE, De Silva AO, Small J, et al (2011) Biomagnification of perfluorinated compounds in a remote terrestrial food chain: Lichen-caribou-wolf. *Environmental Science & Technology* 45:8665–8673. <https://doi.org/10.1021/es201353v>
- Müller CE, Gerecke AC, Bogdal C, et al (2012) Atmospheric fate of poly- and perfluorinated alkyl substances (PFASs): I. Day–night patterns of air concentrations in summer in Zurich, Switzerland. *Environmental Pollution* 169:196–203. <https://doi.org/10.1016/j.envpol.2012.04.010>
- Munoz G, Budzinski H, Babut M, et al (2019) Temporal variations of perfluoroalkyl substances partitioning between surface water, suspended sediment, and biota in a macrotidal estuary. *Chemosphere* 233:319–326. <https://doi.org/10.1016/j.chemosphere.2019.05.281>
- Munoz G, Giraudel JL, Botta F, et al (2015) Spatial distribution and partitioning behavior of selected poly- and perfluoroalkyl substances in freshwater ecosystems: A French nationwide survey. *Science of The Total Environment* 517:48–56. <https://doi.org/10.1016/j.scitotenv.2015.02.043>
- Nakata H, Kannan K, Nasu T, et al (2006) Perfluorinated contaminants in sediments and aquatic organisms collected from shallow water and tidal flat areas of the Ariake Sea, Japan: environmental fate of perfluorooctane sulfonate in aquatic ecosystems. *Environmental Science & Technology* 40:4916–4921. <https://doi.org/10.1021/es0603195>
- Nakayama SF, Yoshikane M, Onoda Y, et al (2019) Worldwide trends in tracing poly- and perfluoroalkyl substances (PFAS) in the environment. *TrAC Trends in Analytical Chemistry*. <https://doi.org/10.1016/j.trac.2019.02.011>
- NAS (2017) The National Academies of Sciences, Engineering, and Medicine (NAS): Using 21st Century Science to Improve Risk-Related Evaluations. Division on Earth and Life Studies, Committee on Incorporating 21st Century Science Into Risk-Based Evaluations. February, 2017
- Navarro I, de la Torre A, Sanz P, et al (2017) Uptake of perfluoroalkyl substances and halogenated flame retardants by crop plants grown in biosolids-amended soils. *Environmental Research* 152:199–206. <https://doi.org/10.1016/j.envres.2016.10.018>
- Ng A, Weerakoon D, Lim E, Padhye LP (2019) Fate of environmental pollutants. *Water Environment Research*. <https://doi.org/10.1002/wer.1225>
- NGWA (2017) National Ground Water Association (NGWA): Groundwater and PFAS: State of Knowledge and Practice. Section 4: Fate and Transport. Available at: <http://comments.ngwa.org/wp-content/uploads/2017/06/pfas-doc-section-4-transport-draft.pdf>. Accessed 1 May 2018

- NHDHHS (2016) New Hampshire Department of Health and Human Services (NHDHHS): Pease PFC Blood Testing Program: April 2015 – October 2015, Pease Tradeport, Portsmouth, NH. Division of Public Health Services, June 16, 2016
- Nielsen CJ (2014) Potential PFOA Precursors. Literature study and theoretical assessment of abiotic degradation pathways leading to PFOA. The Centre for Theoretical and Computational Chemistry (CTCC), Department of Chemistry University of Oslo, Oslo, Norway
- Nilsson H, Kärrman A, Rotander A, et al (2010) Inhalation exposure to fluorotelomer alcohols yield perfluorocarboxylates in human blood? *Environmental Science & Technology* 44:7717–7722. <https://doi.org/10.1021/es101951t>
- Nilsson H, Kärrman A, Rotander A, et al (2013) Biotransformation of fluorotelomer compound to perfluorocarboxylates in humans. *Environment International* 51:8–12. <https://doi.org/10.1016/j.envint.2012.09.001>
- NJDEP (2017) New Jersey Department of Environmental Protection (NJDEP): Christie administration takes action to enhance protection of New Jersey’s drinking water, Department of Environmental Protection moving forward with science-based health standards for PFOA and PFNA. News Release, November 1, 2017. Available at: http://www.nj.gov/dep/newsrel/2017/17_0104.html. Accessed 9 Nov 2017
- Nordén M, Berger U, Engwall M (2016) Developmental toxicity of PFOS and PFOA in great cormorant (*Phalacrocorax carbo sinensis*), herring gull (*Larus argentatus*) and chicken (*Gallus gallus domesticus*). *Environmental Science and Pollution Research* 23:10855–10862. <https://doi.org/10.1007/s11356-016-6285-1>
- Nyberg E, Awad R, Bignert A, et al (2018) Inter-individual, inter-city, and temporal trends of per- and polyfluoroalkyl substances in human milk from Swedish mothers between 1972 and 2016. *Environmental Science: Processes & Impacts* 20:1136–1147. <https://doi.org/10.1039/C8EM00174J>
- O’Connell SG, Arendt M, Segars A, et al (2010) Temporal and spatial trends of perfluorinated compounds in juvenile loggerhead sea turtles (*Caretta caretta*) along the East Coast of the United States. *Environmental Science & Technology* 44:5202–5209. <https://doi.org/10.1021/es9036447>
- O’Connor JC, Munley SM, Serex TL, Buck RC (2014) Evaluation of the reproductive and developmental toxicity of 6:2 fluorotelomer alcohol in rats. *Toxicology* 317:6–16. <https://doi.org/10.1016/j.tox.2014.01.002>
- OECD (2018) Organisation for Economic Cooperation and Development (OECD): Toward a new comprehensive global database of per- and polyfluoroalkyl substances (PFASs): Summary report on updating the OECD 2007 list of per- and polyfluoroalkyl substances (PFASs). Available at:

[https://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=ENV-JM-MONO\(2018\)7&doclanguage=en](https://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=ENV-JM-MONO(2018)7&doclanguage=en). Accessed 26 Jul 2019

OECD (2013) Organisation for Economic Cooperation and Development (OECD): Synthesis paper on per- and polyfluorinated chemicals (PFCs). OECD Environment Directorate, Environment, Health and Safety Division, Paris, France

OEHHA (2012) Office of Environmental Health Hazard Assessment (OEHHA): Amended Final Statement of Reasons, Division 4.5, Title 22, Cal. Code of Regulations, Chapter 54, Green Chemistry Hazard Traits. Available at:

OEHHA (2018) Office of Environmental Health Hazard Assessment (OEHHA): Recommendation for interim notification levels for perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS). A memorandum to the Deputy Director, Division of Drinking Water, California State Water Resources Control Board, June 26, 2018 Available at: https://www.waterboards.ca.gov/drinking_water/certlic/drinkingwater/documents/pfos_and_pfoa/OEHHA_Recommended_Int_NL_Jun_26_2018.pdf. Accessed 7 Oct 2019

OEHHA (2019) Office of Environmental Health Hazard Assessment (OEHHA): Notification Level Recommendations for Perfluorooctanoic Acid (PFOA) and Perfluorooctane Sulfonate (PFOS). In: OEHHA. <https://oehha.ca.gov/water/notification-level/notification-level-recommendations-perfluorooctanoic-acid-pfoa>. Accessed 29 Oct 2019

Office of Congressman Dan Kildee (2019) Repres. Kildee, Fitzpatrick launch congressional PFAS task force. January 23, 2019 Press Release. Available at: <https://dankildee.house.gov/media/press-releases/rep-kildee-fitzpatrick-launch-congressional-pfas-task-force>. Accessed 4 Mar 2019

Ohmori K, Kudo N, Katayama K, Kawashima Y (2003) Comparison of the toxicokinetics between perfluorocarboxylic acids with different carbon chain length. *Toxicology* 184:135–140. [https://doi.org/10.1016/S0300-483X\(02\)00573-5](https://doi.org/10.1016/S0300-483X(02)00573-5)

Oliaei F, Kriens D, Weber R, Watson A (2013) PFOS and PFC releases and associated pollution from a PFC production plant in Minnesota (USA). *Environmental Science and Pollution Research* 20:1977–1992. <https://doi.org/10.1007/s11356-012-1275-4>

Olsen GW, Burris JM, Ehresman DJ, et al (2007a) Half-life of serum elimination of perfluorooctanesulfonate, perfluorohexanesulfonate, and perfluorooctanoate in retired fluorochemical production workers. *Environmental Health Perspectives* 115:1298–1305. <https://doi.org/10.1289/ehp.10009>

Olsen GW, Mair DC, Reagen WK, et al (2007b) Preliminary evidence of a decline in perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA) concentrations in American Red Cross blood donors. *Chemosphere* 68:105–111. <https://doi.org/10.1016/j.chemosphere.2006.12.031>

- Ostertag SK, Chan HM, Moisey J, et al (2009) Historic dietary exposure to perfluorooctane sulfonate, perfluorinated carboxylates, and fluorotelomer unsaturated carboxylates from the consumption of store-bought and restaurant foods for the Canadian population. *Journal of Agricultural and Food Chemistry* 57:8534–8544. <https://doi.org/10.1021/jf9014125>
- Pan CG, Liu YS, Ying GC (2016) Perfluoroalkyl substances (PFASs) in wastewater treatment plants and drinking water treatment plants: removal efficiency and exposure risk. *Water Research* 106:562–570. <https://doi.org/10.1016/j.watres.2016.10.045>
- Pan CG, Ying GG, Liu YS, et al (2014a) Contamination profiles of perfluoroalkyl substances in five typical rivers of the Pearl River Delta region, South China. *Chemosphere* 114:16–25. <https://doi.org/10.1016/j.chemosphere.2014.04.005>
- Pan CG, Ying GG, Zhao JL, et al (2015) Spatial distribution of perfluoroalkyl substances in surface sediments of five major rivers in China. *Archives of Environmental Contamination and Toxicology* 68:566–576. <https://doi.org/10.1007/s00244-014-0113-8>
- Pan CG, Ying GG, Zhao JL, et al (2014b) Spatiotemporal distribution and mass loadings of perfluoroalkyl substances in the Yangtze River of China. *Science of The Total Environment* 493:580–587. <https://doi.org/10.1016/j.scitotenv.2014.06.033>
- Pan Y, Zhang H, Cui Q, et al (2018) Worldwide distribution of novel perfluoroether carboxylic and sulfonic acids in surface water. *Environmental Science & Technology* 52:7621–7629. <https://doi.org/10.1021/acs.est.8b00829>
- Pan Y, Zhu Y, Zheng T, et al (2017) Novel chlorinated polyfluorinated ether sulfonates and legacy per-/polyfluoroalkyl substances: placental transfer and relationship with serum albumin and glomerular filtration rate. *Environmental Science & Technology* 51:634–644. <https://doi.org/10.1021/acs.est.6b04590>
- Papadopoulou E, Sabaredzovic A, Namork E, et al (2016) Exposure of Norwegian toddlers to perfluoroalkyl substances (PFAS): the association with breastfeeding and maternal PFAS concentrations. *Environment International* 94:687–694. <https://doi.org/10.1016/j.envint.2016.07.006>
- Paul AG, Jones KC, Sweetman AJ (2009) A first global production, emission, and environmental inventory for perfluorooctane sulfonate. *Environmental Science & Technology* 43:386–392. <https://doi.org/10.1021/es802216n>
- Pedersen KE, Letcher RJ, Sonne C, et al (2016) Per- and polyfluoroalkyl substances (PFASs) – New endocrine disruptors in polar bears (*Ursus maritimus*)? *Environment International* 96:180–189. <https://doi.org/10.1016/j.envint.2016.07.015>
- Perez F, Llorca M, Farré M, Barceló D (2012) Automated analysis of perfluorinated compounds in human hair and urine samples by turbulent flow chromatography coupled to tandem mass

spectrometry. *Analytical and Bioanalytical Chemistry* 402:2369–2378.
<https://doi.org/10.1007/s00216-011-5660-5>

Pérez F, Llorca M, Köck-Schulmeyer M, et al (2014) Assessment of perfluoroalkyl substances in food items at global scale. *Environmental Research* 135:181–189.
<https://doi.org/10.1016/j.envres.2014.08.004>

Pérez F, Nadal M, Navarro-Ortega A, et al (2013) Accumulation of perfluoroalkyl substances in human tissues. *Environment International* 59:354–362. <https://doi.org/10.1016/j.envint.2013.06.004>

Phillips MM, Dinglasan-Panlilio MJA, Mabury SA, et al (2007) Fluorotelomer acids are more toxic than perfluorinated acids. *Environmental Science & Technology* 41:7159–7163.
<https://doi.org/10.1021/es070734c>

Plumlee MH, Larabee J, Reinhard M (2008) Perfluorochemicals in water reuse. *Chemosphere* 72:1541–1547. <https://doi.org/10.1016/j.chemosphere.2008.04.057>

Poothong S, Thomsen C, Padilla-Sanchez JA, et al (2017) Distribution of novel and well-known poly- and perfluoroalkyl substances (PFASs) in human serum, plasma, and whole blood. *Environmental Science & Technology* 51:13388–13396. <https://doi.org/10.1021/acs.est.7b03299>

Powley CR, George SW, Russell MH, et al (2008) Polyfluorinated chemicals in a spatially and temporally integrated food web in the Western Arctic. *Chemosphere* 70:664–672.
<https://doi.org/10.1016/j.chemosphere.2007.06.067>

Prevedouros K, Cousins IT, Buck RC, Korzeniowski SH (2006) Sources, fate and transport of perfluorocarboxylates. *Environmental Science & Technology* 40:32–44.
<https://doi.org/10.1021/es0512475>

Qi Y, Huo S, Xi B, et al (2016) Spatial distribution and source apportionment of PFASs in surface sediments from five lake regions, China. *Scientific Reports* 6:22674.
<https://doi.org/10.1038/srep22674>

Qin J, Waldroup D, Calewatts D, et al (2018) Non-fluorinated water-based superhydrophobic surfaces

Rae CJM, Craig L, Slone TW, et al (2015) Evaluation of chronic toxicity and carcinogenicity of ammonium 2,3,3,3-tetrafluoro-2-(heptafluoropropoxy)-propanoate in Sprague–Dawley rats. *Toxicology Reports* 2:939–949. <https://doi.org/10.1016/j.toxrep.2015.06.001>

Rahman MF, Peldszus S, Anderson WB (2014) Behaviour and fate of perfluoroalkyl and polyfluoroalkyl substances (PFASs) in drinking water treatment: a review. *Water Research* 50:318–340.
<https://doi.org/10.1016/j.watres.2013.10.045>

- Rainieri S, Conlledo N, Langerholc T, et al (2017) Toxic effects of perfluorinated compounds at human cellular level and on a model vertebrate. *Food and Chemical Toxicology* 104:14–25. <https://doi.org/10.1016/j.fct.2017.02.041>
- Ramhøj L, Hass U, Boberg J, et al (2018) Perfluorohexane sulfonate (PFHxS) and a mixture of endocrine disruptors reduce thyroxine levels and cause anti-androgenic effects in rats. *Toxicological Sciences* 163:579–591. <https://doi.org/10.1093/toxsci/kfy055>
- Rand AA, Mabury SA (2012a) In vitro interactions of biological nucleophiles with fluorotelomer unsaturated acids and aldehydes: fate and consequences. *Environmental Science & Technology* 46:7398–7406. <https://doi.org/10.1021/es3008485>
- Rand AA, Mabury SA (2012b) Assessing the structure–activity relationships of fluorotelomer unsaturated acids and aldehydes with glutathione: Reactivity of glutathione with fluorotelomer unsaturated acids and aldehydes. *Cell Biology and Toxicology* 28:115–124. <https://doi.org/10.1007/s10565-012-9211-4>
- Rand AA, Mabury SA (2013) Covalent binding of fluorotelomer unsaturated aldehydes (FTUALs) and carboxylic acids (FTUCAs) to proteins. *Environmental Science & Technology* 130109073401005. <https://doi.org/10.1021/es303760u>
- Rand AA, Mabury SA (2014) Protein binding associated with exposure to fluorotelomer alcohols (FTOHs) and polyfluoroalkyl phosphate esters (PAPs) in rats. *Environmental Science & Technology* 140205115442006. <https://doi.org/10.1021/es404390x>
- Rand AA, Mabury SA (2017) Is there a human health risk associated with indirect exposure to perfluoroalkyl carboxylates (PFCAs)? *Toxicology* 375:28–36. <https://doi.org/10.1016/j.tox.2016.11.011>
- Rand AA, Rooney JP, Butt CM, et al (2014) Cellular toxicity associated with exposure to perfluorinated carboxylates (PFCAs) and their metabolic precursors. *Chemical Research in Toxicology* 27:42–50. <https://doi.org/10.1021/tx400317p>
- Rankin K, Mabury SA, Jenkins TM, Washington JW (2016) A North American and global survey of perfluoroalkyl substances in surface soils: distribution patterns and mode of occurrence. *Chemosphere* 161:333–341. <https://doi.org/10.1016/j.chemosphere.2016.06.109>
- Rappazzo K, Coffman E, Hines E (2017) Exposure to perfluorinated alkyl substances and health outcomes in children: a systematic review of the epidemiologic literature. *International Journal of Environmental Research and Public Health* 14:691. <https://doi.org/10.3390/ijerph14070691>
- Rayne S, Forest K (2009) Perfluoroalkyl sulfonic and carboxylic acids: a critical review of physicochemical properties, levels and patterns in waters and wastewaters, and treatment methods. *Journal of Environmental Science and Health, Part A* 44:1145–1199. <https://doi.org/10.1080/10934520903139811>

- Renzi M, Guerranti C, Giovani A, et al (2013) Perfluorinated compounds: levels, trophic web enrichments and human dietary intakes in transitional water ecosystems. *Marine Pollution Bulletin* 76:146–157. <https://doi.org/10.1016/j.marpolbul.2013.09.014>
- Rodea-Palomares I, Leganés F, Rosal R, Fernández-Piñas F (2012) Toxicological interactions of perfluorooctane sulfonic acid (PFOS) and perfluorooctanoic acid (PFOA) with selected pollutants. *Journal of Hazardous Materials* 201–202:209–218. <https://doi.org/10.1016/j.jhazmat.2011.11.061>
- Rodriguez-Jorquera IA, Silva-Sanchez C, Strynar M, et al (2016) Footprints of urban micro-pollution in protected areas: investigating the longitudinal distribution of perfluoroalkyl acids in wildlife preserves. *Plos One* 11:e0148654. <https://doi.org/10.1371/journal.pone.0148654>
- Rosati JA, Thornburg J, Rodes C (2008) Resuspension of particulate matter from carpet due to human activity. *Aerosol Science and Technology* 42:472–482. <https://doi.org/10.1080/02786820802187069>
- Roscales JL, Vicente A, Ryan PG, et al (2019) Spatial and interspecies heterogeneity in concentrations of perfluoroalkyl substances (PFASs) in seabirds of the Southern Ocean. *Environmental Science & Technology* 53:9855–9865. <https://doi.org/10.1021/acs.est.9b02677>
- Rosenmai AK, Ahrens L, le Godec T, et al (2018) Relationship between peroxisome proliferator-activated receptor alpha activity and cellular concentration of 14 perfluoroalkyl substances in HepG2 cells. *Journal of Applied Toxicology* 38:219–226. <https://doi.org/10.1002/jat.3515>
- Rosenmai AK, Taxvig C, Svingen T, et al (2016) Fluorinated alkyl substances and technical mixtures used in food paper-packaging exhibit endocrine-related activity in vitro. *Andrology* 4:662–672. <https://doi.org/10.1111/andr.12190>
- Routti H, Gabrielsen GW, Herzke D, et al (2016) Spatial and temporal trends in perfluoroalkyl substances (PFASs) in ringed seals (*Pusa hispida*) from Svalbard. *Environmental Pollution* 214:230–238. <https://doi.org/10.1016/j.envpol.2016.04.016>
- Royer LA, Lee LS, Russell MH, et al (2015) Microbial transformation of 8:2 fluorotelomer acrylate and methacrylate in aerobic soils. *Chemosphere* 129:54–61. <https://doi.org/10.1016/j.chemosphere.2014.09.077>
- Ruan T, Sulecki LM, Wolstenholme BW, et al (2014) 6:2 Fluorotelomer iodide in vitro metabolism by rat liver microsomes: Comparison with [1,2-¹⁴C] 6:2 fluorotelomer alcohol. *Chemosphere* 112:34–41. <https://doi.org/10.1016/j.chemosphere.2014.02.068>
- Ruan Y, Lalwani D, Kwok KY, et al (2019) Assessing exposure to legacy and emerging per- and polyfluoroalkyl substances via hair – The first nationwide survey in India. *Chemosphere* 229:366–373. <https://doi.org/10.1016/j.chemosphere.2019.04.195>

- Russell MH, Berti WR, Szostek B, et al (2010) Evaluation of PFO formation from the biodegradation of a fluorotelomer-based urethane polymer product in aerobic soils. *Polymer Degradation and Stability* 95:79–85. <https://doi.org/10.1016/j.polymdegradstab.2009.10.004>
- Russell MH, Berti WR, Szostek B, Buck RC (2008) Investigation of the biodegradation potential of a fluoroacrylate polymer product in aerobic soils. *Environmental Science & Technology* 42:800–807. <https://doi.org/10.1021/es0710499>
- Russell MH, Himmelstein MW, Buck RC (2015) Inhalation and oral toxicokinetics of 6:2 FTOH and its metabolites in mammals. *Chemosphere* 120:328–335. <https://doi.org/10.1016/j.chemosphere.2014.07.092>
- Safer Products for Washington Team (2019) SPW Update- Mid September
- Sammut G, Sinagra E, Helmus R, de Voogt P (2017) Perfluoroalkyl substances in the Maltese environment – (I) surface water and rain water. *Science of The Total Environment* 589:182–190. <https://doi.org/10.1016/j.scitotenv.2017.02.128>
- Sanchez D, Houde M, Douville M, et al (2015) Transcriptional and cellular responses of the green alga *Chlamydomonas reinhardtii* to perfluoroalkyl phosphonic acids. *Aquatic Toxicology* 160:31–38. <https://doi.org/10.1016/j.aquatox.2014.12.002>
- Sanchez-Vidal A, Llorca M, Farré M, et al (2015) Delivery of unprecedented amounts of perfluoroalkyl substances towards the deep-sea. *Science of The Total Environment* 526:41–48. <https://doi.org/10.1016/j.scitotenv.2015.04.080>
- Schaider LA, Ackerman JM, Rudel RA (2016) Septic systems as sources of organic wastewater compounds in domestic drinking water wells in a shallow sand and gravel aquifer. *Science of The Total Environment* 547:470–481. <https://doi.org/10.1016/j.scitotenv.2015.12.081>
- Schaider LA, Rudel RA, Ackerman JM, et al (2014) Pharmaceuticals, perfluorosurfactants, and other organic wastewater compounds in public drinking water wells in a shallow sand and gravel aquifer. *Science of The Total Environment* 468–469:384–393. <https://doi.org/10.1016/j.scitotenv.2013.08.067>
- Schechter A, Colacino J, Haffner D, et al (2010) Perfluorinated compounds, polychlorinated biphenyls, and organochlorine pesticide contamination in composite food samples from Dallas, Texas, USA. *Environmental Health Perspectives* 118:796–802. <https://doi.org/10.1289/ehp.0901347>
- Scher DP, Kelly JE, Huset CA, et al (2018) Occurrence of perfluoroalkyl substances (PFAS) in garden produce at homes with a history of PFAS-contaminated drinking water. *Chemosphere* 196:548–555. <https://doi.org/10.1016/j.chemosphere.2017.12.179>

- Scheringer M, Trier X, Cousins IT, et al (2014) Helsingør Statement on poly- and perfluorinated alkyl substances (PFASs). *Chemosphere* 114:337–339. <https://doi.org/10.1016/j.chemosphere.2014.05.044>
- Schlummer M, Gruber L, Fiedler D, et al (2013) Detection of fluorotelomer alcohols in indoor environments and their relevance for human exposure. *Environment International* 57–58:42–49. <https://doi.org/10.1016/j.envint.2013.03.010>
- Schlummer M, Sölch C, Meisel T, et al (2015) Emission of perfluoroalkyl carboxylic acids (PFCA) from heated surfaces made of polytetrafluoroethylene (PTFE) applied in food contact materials and consumer products. *Chemosphere* 129:46–53. <https://doi.org/10.1016/j.chemosphere.2014.11.036>
- Schwanz TG, Llorca M, Farré M, Barceló D (2016) Perfluoroalkyl substances assessment in drinking waters from Brazil, France and Spain. *Science of The Total Environment* 539:143–152. <https://doi.org/10.1016/j.scitotenv.2015.08.034>
- Seals R, Bartell SM, Steenland K (2011) Accumulation and clearance of perfluorooctanoic acid (PFOA) in current and former residents of an exposed community. *Environmental Health Perspectives* 119:119–124. <https://doi.org/10.1289/ehp.1002346>
- Sedlak MD, Benskin JP, Wong A, et al (2017) Per- and polyfluoroalkyl substances (PFASs) in San Francisco Bay wildlife: temporal trends, exposure pathways, and notable presence of precursor compounds. *Chemosphere* 185:1217–1226. <https://doi.org/10.1016/j.chemosphere.2017.04.096>
- Sedlak MD, Greig DJ (2012) Perfluoroalkyl compounds (PFCs) in wildlife from an urban estuary. *Journal of Environmental Monitoring* 14:146–154. <https://doi.org/10.1039/C1EM10609K>
- Sepulvado JG, Blaine AC, Hundal LS, Higgins CP (2011) Occurrence and fate of perfluorochemicals in soil following the land application of municipal biosolids. *Environmental Science & Technology* 45:8106–8112. <https://doi.org/10.1021/es103903d>
- SFE (2018) San Francisco Department of the Environment (SFE): Regulation #SFE 207 8-01-PPO/GBRCBO-Adopting approved alternative products for sustainable carpet for city departments. Available at: https://sfenvironment.org/sites/default/files/policy/regulation_sfe-2018-01-ppo_gbrco.pdf. Accessed 8 May 2019
- Shan G, Wei M, Zhu L, et al (2014) Concentration profiles and spatial distribution of perfluoroalkyl substances in an industrial center with condensed fluorochemical facilities. *Science of The Total Environment* 490:351–359. <https://doi.org/10.1016/j.scitotenv.2014.05.005>
- Shi G, Cui Q, Pan Y, et al (2017) 6:2 Chlorinated polyfluorinated ether sulfonate, a PFOS alternative, induces embryotoxicity and disrupts cardiac development in zebrafish embryos. *Aquatic Toxicology* 185:67–75. <https://doi.org/10.1016/j.aquatox.2017.02.002>

- Shi Y, Vestergren R, Xu L, et al (2016) Human exposure and elimination kinetics of chlorinated polyfluoroalkyl ether sulfonic acids (Cl-PFESAs). *Environ Sci Technol* 50:2396–2404. <https://doi.org/10.1021/acs.est.5b05849>
- Shoaeioskouei S (2012) Perfluorinated compounds in landfill leachate from discarded carpets. M.S. Thesis, Faculty of Graduate Studies (Civil Engineering), the University of British Columbia, Vancouver. June 2012
- Shoeib M, Harner T, Ikonomou M, Kannan K (2004) Indoor and outdoor air concentrations and phase partitioning of perfluoroalkyl sulfonamides and polybrominated diphenyl ethers. *Environmental Science & Technology* 38:1313–1320. <https://doi.org/10.1021/es0305555>
- Shoeib M, Harner T, M. Webster G, Lee SC (2011) Indoor sources of poly- and perfluorinated compounds (PFCs) in Vancouver, Canada: Implications for human exposure. *Environmental Science & Technology* 45:7999–8005. <https://doi.org/10.1021/es103562v>
- Shoeib M, Harner T, Wilford BH, et al (2005) Perfluorinated sulfonamides in indoor and outdoor air and indoor dust: occurrence, partitioning, and human exposure. *Environmental Science & Technology* 39:6599–6606. <https://doi.org/10.1021/es048340y>
- Shoeib M, Vlahos P, Harner T, et al (2010) Survey of polyfluorinated chemicals (PFCs) in the atmosphere over the northeast Atlantic Ocean. *Atmospheric Environment* 44:2887–2893. <https://doi.org/10.1016/j.atmosenv.2010.04.056>
- Siebenaler R, Cameron R, Butt CM, et al (2017) Serum perfluoroalkyl acids (PFAAs) and associations with behavioral attributes. *Chemosphere* 184:687–693. <https://doi.org/10.1016/j.chemosphere.2017.06.023>
- Sinclair E, Kannan K (2006) Mass loading and fate of perfluoroalkyl surfactants in wastewater treatment plants. *Environmental Science & Technology* 40:1408–1414. <https://doi.org/10.1021/es051798v>
- Sinclair E, Kim SK, Akinleye HB, Kannan K (2007) Quantitation of gas-phase perfluoroalkyl surfactants and fluorotelomer alcohols released from nonstick cookware and microwave popcorn bags. *Environmental Science & Technology* 41:1180–1185. <https://doi.org/10.1021/es062377w>
- Singh J, Jayanti RK, Kilgore JE, et al (2005) What goes around comes around: Understanding trust–value dilemmas of market relationships. *Journal of Public Policy & Marketing* 24:38–62. <https://doi.org/10.1509/jppm.24.1.38.63882>
- Skaar JS, Ræder EM, Lyche JL, et al (2018) Elucidation of contamination sources for poly- and perfluoroalkyl substances (PFASs) on Svalbard (Norwegian Arctic). *Environmental Science and Pollution Research* 26:7356–7363. <https://doi.org/10.1007/s11356-018-2162-4>

- Skuladottir M, Ramel A, Rytter D, et al (2015) Examining confounding by diet in the association between perfluoroalkyl acids and serum cholesterol in pregnancy. *Environmental Research* 143:33–38. <https://doi.org/10.1016/j.envres.2015.09.001>
- Slotkin TA, MacKillop EA, Melnick RL, et al (2008) Developmental neurotoxicity of perfluorinated chemicals modeled in vitro. *Environmental Health Perspectives* 116:716–722. <https://doi.org/10.1289/ehp.11253>
- Soloff AC, Wolf BJ, White ND, et al (2017) Environmental perfluorooctane sulfonate exposure drives T cell activation in bottlenose dolphins: PFOS-induced T cell activation in dolphins. *Journal of Applied Toxicology* 37:1108–1116. <https://doi.org/10.1002/jat.3465>
- SSCCH (2017) Secretariat of the Stockholm Convention Clearing House (SSCCH): All POPs listed in the Stockholm Convention. Available at: <http://chm.pops.int/TheConvention/ThePOPs/ListingofPOPs/tabid/2509/Default.aspx>. Accessed 22 Sep 2017
- Stahl T, Heyn J, Thiele H, et al (2009) Carryover of perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS) from soil to plants. *Archives of Environmental Contamination and Toxicology* 57:289–298. <https://doi.org/10.1007/s00244-008-9272-9>
- Stahl T, Mattern D, Brunn H (2011) Toxicology of perfluorinated compounds. *Environmental Sciences Europe* 23:38. <https://doi.org/10.1186/2190-4715-23-38>
- State of Washington (2017) Survey of per- and poly-fluoroalkyl substances (PFASs) in rivers and lakes, 2016, Publication No. 17-03-021, September 2017. Available at: <https://fortress.wa.gov/ecy/publications/SummaryPages/1703021.html>. Accessed 30 Apr 2018
- State of Washington (2018a) House Bill 2793, 2018 Regular Session, 65th Legislature. Available at:
- State of Washington (2018b) Engrossed Substitute - ESSB 6413. Available at: <http://apps2.leg.wa.gov/billsummary?BillNumber=6413&Year=2017&BillNumber=6413&Year=2017>. Accessed 24 Apr 2018
- State of Washington (2018c) House Bill 2658, 2018 Regular Session, 65th Legislature. Available at: <http://lawfilesexternal.leg.wa.gov/biennium/2017-18/Pdf/Bills/House%20Bills/2658.pdf>. Accessed 7 Feb 2018
- Stock NL, Lau FK, Ellis DA, et al (2004) Polyfluorinated telomer alcohols and sulfonamides in the North American troposphere. *Environmental Science & Technology* 38:991–996. <https://doi.org/10.1021/es034644t>
- Strynar MJ, Lindstrom AB (2008) Perfluorinated compounds in house dust from Ohio and North Carolina, USA. *Environmental Science & Technology* 42:3751–3756. <https://doi.org/10.1021/es7032058>

- Su H, Shi Y, Lu Y, et al (2017) Home produced eggs: an important pathway of human exposure to perfluorobutanoic acid (PFBA) and perfluorooctanoic acid (PFOA) around a fluorochemical industrial park in China. *Environment International* 101:1–6. <https://doi.org/10.1016/j.envint.2017.01.016>
- Sun H, Gerecke AC, Giger W, Alder AC (2011) Long-chain perfluorinated chemicals in digested sewage sludges in Switzerland. *Environmental Pollution* 159:654–662. <https://doi.org/10.1016/j.envpol.2010.09.020>
- Sun H, Zhang X, Wang L, et al (2012) Perfluoroalkyl compounds in municipal WWTPs in Tianjin, China—concentrations, distribution and mass flow. *Environmental Science and Pollution Research* 19:1405–1415. <https://doi.org/10.1007/s11356-011-0727-6>
- Sun M, Arevalo E, Strynar M, et al (2016) Legacy and emerging perfluoroalkyl substances are important drinking water contaminants in the Cape Fear River watershed of North Carolina. *Environmental Science & Technology Letters* 3:415–419. <https://doi.org/10.1021/acs.estlett.6b00398>
- Sunderland EM, Hu XC, Dassuncao C, et al (2019) A review of the pathways of human exposure to poly- and perfluoroalkyl substances (PFASs) and present understanding of health effects. *Journal of Exposure Science & Environmental Epidemiology* 29:131–147. <https://doi.org/10.1038/s41370-018-0094-1>
- Sundström M, Chang SC, Noker PE, et al (2012) Comparative pharmacokinetics of perfluorohexanesulfonate (PFHxS) in rats, mice, and monkeys. *Reproductive Toxicology* 33:441–451. <https://doi.org/10.1016/j.reprotox.2011.07.004>
- Sutton R, Sedlak M, Sun J, Lin D (2017) Contaminants of emerging concern in San Francisco Bay: A strategy for future investigations, San Francisco Estuary Institute, November, 2017 revision. Regional Monitoring Program for Water Quality in San Francisco Bay (RMP), Richmond, CA, RMP Contribution #815. Available at: http://www.sfei.org/sites/default/files/biblio_files/RMP%20CEC%20Strategy%20Nov%202017_0.pdf. Accessed 29 Nov 2017
- Taniyasu S, Yamashita N, Moon HB, et al (2013) Does wet precipitation represent local and regional atmospheric transportation by perfluorinated alkyl substances? *Environment International* 55:25–32. <https://doi.org/10.1016/j.envint.2013.02.005>
- Tao L, Kannan K, Aldous KM, et al (2008a) Biomonitoring of perfluorochemicals in plasma of New York State personnel responding to the World Trade Center disaster. *Environmental Science & Technology* 42:3472–3478. <https://doi.org/10.1021/es8000079>
- Tao L, Kannan K, Wong CM, et al (2008b) Perfluorinated compounds in human milk from Massachusetts, U.S.A. *Environmental Science & Technology* 42:3096–3101. <https://doi.org/10.1021/es702789k>

- Target Inc. (2017) Target announces new chemical strategy including policy and goals for its products and operations. Press release via Target's Bullseye View, Target Brands, Inc., January 25, 2017. Available at: <https://corporate.target.com/article/2017/01/chemical-policy-and-goals>. Accessed 15 Aug 2017
- Tartu S, Aars J, Andersen M, et al (2018) Choose your poison-Space-use strategy influences pollutant exposure in Barents Sea polar bears. *Environmental Science & Technology* 52:3211–3221. <https://doi.org/10.1021/acs.est.7b06137>
- Tartu S, Lille-Langøy R, Størseth TR, et al (2017) Multiple-stressor effects in an apex predator: combined influence of pollutants and sea ice decline on lipid metabolism in polar bears. *Scientific Reports* 7:16487. <https://doi.org/10.1038/s41598-017-16820-5>
- Tatum-Gibbs K, Wambaugh JF, Das KP, et al (2011) Comparative pharmacokinetics of perfluorononanoic acid in rat and mouse. *Toxicology* 281:48–55. <https://doi.org/10.1016/j.tox.2011.01.003>
- Teli MD (2018) 9 – Finishing of carpets for value addition. *Advances in Carpet Manufacture, the Textile Institute Book series*. edited by K.K. Goswami, second edition. , Available at: <https://doi.org/10.1016/B978-0-08-101131-7.00001-0>. Accessed 25 May 2018
- Tian Y, Yao Y, Chang S, et al (2018) Occurrence and phase distribution of neutral and ionizable per- and polyfluoroalkyl substances (PFASs) in the atmosphere and plant leaves around landfills: a case study in Tianjin, China. *Environmental Science & Technology* 52:1301–1310. <https://doi.org/10.1021/acs.est.7b05385>
- Tian Z, Kim SK, Shoeib M, et al (2016) Human exposure to per- and polyfluoroalkyl substances (PFASs) via house dust in Korea: implication to exposure pathway. *Science of The Total Environment* 553:266–275. <https://doi.org/10.1016/j.scitotenv.2016.02.087>
- Tipton JJ, Guillette LJ, Lovelace S, et al (2017) Analysis of PFAAs in American alligators part 1: concentrations in alligators harvested for consumption during South Carolina public hunts. *Journal of Environmental Sciences* 61:24–30. <https://doi.org/10.1016/j.jes.2017.05.045>
- Tomy GT, Budakowski W, Halldorson T, et al (2004) Fluorinated organic compounds in an eastern Arctic marine food web. *Environmental Science & Technology* 38:6475–6481. <https://doi.org/10.1021/es049620g>
- Trier X, Granby K, Christensen JH (2011) Polyfluorinated surfactants (PFS) in paper and board coatings for food packaging. *Environmental Science and Pollution Research* 18:1108–1120
- Trudel D, Horowitz L, Wormuth M, et al (2008) Estimating consumer exposure to PFOS and PFOA. *Risk Analysis* 28:251–269. <https://doi.org/10.1111/j.1539-6924.2008.01017.x>

- Tsai MS, Miyashita C, Araki A, et al (2018) Determinants and temporal trends of perfluoroalkyl substances in pregnant women: the Hokkaido study on environment and children's health. *International Journal of Environmental Research and Public Health* 15:989. <https://doi.org/10.3390/ijerph15050989>
- Ulhaq M, Carlsson G, Örn S, Norrgren L (2013) Comparison of developmental toxicity of seven perfluoroalkyl acids to zebrafish embryos. *Environmental Toxicology and Pharmacology* 36:423–426. <https://doi.org/10.1016/j.etap.2013.05.004>
- UNCW (2018) University of North Carolina at Wilmington (UNCW): Report to the Environmental Review Commission from the University of North Carolina at Wilmington Regarding the Implementation of Section 20.(a)(2) of House Bill 56 (S.L. 2017-209). Available at: <https://www.ncleg.net/documentsites/committees/ERC/ERC%20Reports%20Received/2018/CFPUA%20and%20UNC-W/2018-April%20HB%2056%20UNCW%20Rpt.pdf>. Accessed 9 Apr 2018
- UNEP/POPS/POPRC (2017) United Nations Environment Programme, Stockholm Convention on Persistent Organic Pollutants, Persistent Organic Pollutants Review Committee (UNEP/POPS/POPRC): Highlights of the thirteenth meeting of the Persistent Organic Pollutants Review Committee (POPRC.13), Rome, Italy, from 17 to 20 October 2017. Available at: <http://www.pops.int/Default.aspx?tabid=5965>. Accessed 10 Nov 2017
- UNEP/POPS/POPRC (2019) United Nations Environment Programme, Stockholm Convention on Persistent Organic Pollutants, Persistent Organic Pollutants Review Committee (UNEP/POPS/POPRC): POPRC recommendations for listing chemicals. Available at: <http://chm.pops.int/tabid/243/Default.aspx>. Accessed 15 Jul 2019
- U.S. Bureau of Labor Statistics (2019) Employment Projections. <https://data.bls.gov/projections/occupationProj>. Accessed 26 Sep 2019
- U.S. Census Bureau (2007) Economic census. Available at: <https://www.census.gov/data/tables/2007/econ/census/manufacturing-reports.html>. Accessed 25 Sep 2019
- U.S. Census Bureau (2012) Economic census. Available at: <https://www.census.gov/data/tables/2012/econ/census/manufacturing-reports.html>. Accessed 25 Sep 2019
- U.S. Census Bureau (2017a) Economic census. Available at: <https://data.census.gov/cedsci/table?hidePreview=true&table=EC1731BASIC&tid=ECNBASIC2017.EC1731BASIC&lastDisplayedRow=647&q=EC1731BASIC%3A%20Manufacturing%3A%20Summary%20Statistics%20for%20the%20U.S.,%20States,%20and%20Selected%20Geographies%3A%202017>. Accessed 25 Sep 2019

- U.S. Census Bureau (2017b) American FactFinder: Community Facts. Available at:
<https://factfinder.census.gov/faces/nav/jsf/pages/index.xhtml>. Accessed: 21 Sep 2017.
<https://factfinder.census.gov/faces/nav/jsf/pages/searchresults.xhtml?refresh=t>
- U.S. EPA (2017a) U.S. Environmental Protection Agency (U.S. EPA): Chemistry Dashboard. Available at:
<https://comptox.epa.gov/dashboard>. Data are calculated and updated regularly by OPERA [OPEn (quantitative) Structure-activity Relationship Application], a standalone free and open source command line application in Matlab (Version 8.2) providing QSAR models predictions. Model validation data set may be found here:
<http://esc.syrres.com/interkow/EpiSuiteData.htm>. Accessed 6 Nov 2017
- U.S. EPA (2009) U.S. Environmental Protection Agency (U.S. EPA): Long-chain perfluorinated chemicals (PFCs) action plan. December 30, 2009. Available at:
https://www.epa.gov/sites/production/files/2016-01/documents/pfcs_action_plan1230_09.pdf. Accessed 23 Aug 2017
- U.S. EPA (2017b) U.S. Environmental Protection Agency (U.S. EPA): Occurrence Data for the Unregulated Contaminant Monitoring Rule. Available at:
<https://www.epa.gov/dwucmr/occurrence-data-unregulated-contaminant-monitoring-rule>. Accessed 9 Nov 2017
- U.S. EPA (2011) U.S. Environmental Protection Agency (U.S. EPA): Exposure Factors Handbook, 2011 Edition (Final Report). U.S. Environmental Protection Agency, Washington, DC, EPA/600/R-09/052F, 2011. Available at: <https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=236252>. Accessed 22 Aug 2017
- U.S. EPA (2016a) U.S. Environmental Protection Agency (U.S. EPA): EJ 2020 Action Agenda: The U.S. Environmental Protection Agency's 2016 – 2020 Strategic Plan. Available at:
https://www.epa.gov/sites/production/files/2016-05/documents/052216_ej_2020_strategic_plan_final_0.pdf. Accessed 2 Oct 2017
- U.S. EPA (2016b) U.S. Environmental Protection Agency (U.S. EPA): Drinking water health advisories for PFOA and PFOS. Available at: <https://www.epa.gov/ground-water-and-drinking-water/drinking-water-health-advisories-pfoa-and-pfos>. Accessed 14 Oct 2017
- U.S. EPA (2002a) U.S. Environmental Protection Agency (U.S. EPA): 40 CFR Part 721 Perfluoroalkyl Sulfonates; Significant New Use Rule. Federal Register, Vol. 67, No. 236, December 9, 2002. Available at: <https://www.gpo.gov/fdsys/pkg/FR-2002-12-09/pdf/02-31011.pdf>. Accessed 17 May 2017
- U.S. EPA (2002b) U.S. Environmental Protection Agency (U.S. EPA): 40 CFR Part 721 Perfluoroalkyl Sulfonates; Significant New Use Rule; Final Rule and Supplemental Proposed Rule. Federal Register, Vol. 67, No. 47, March 11, 2002. Available at: <https://www.gpo.gov/fdsys/pkg/FR-2002-03-11/pdf/02-5746.pdf>. Accessed 17 May 2017

- U.S. EPA (2007) U.S. Environmental Protection Agency (U.S. EPA): 40 CFR Part 721. Perfluoroalkyl Sulfonates, Significant New Use Rule. Federal Register, Vol. 72, No. 194, October 9, 2007. Available at: <https://www.gpo.gov/fdsys/pkg/FR-2007-10-09/pdf/E7-19828.pdf>. Accessed 17 May 2017
- U.S. EPA (2013) U.S. Environmental Protection Agency (U.S. EPA): 40 CFR Parts 9 of 721 Perfluoroalkyl Carboxylate Substances; Final Significant New Use Rule. U.S. EPA, Washington, DC. Available at: <https://www.regulations.gov/contentStreamer?documentId=EPA-HQ-OPPT-2012-0268-0034&contentType=pdf>. Accessed 28 Nov 2017
- U.S. EPA (2015) U.S. Environmental Protection Agency (U.S. EPA): Long-chain perfluoroalkyl carboxylate and perfluoroalkyl sulfonate chemical substances; Significant New Use Rule. Available at: <https://www.federalregister.gov/documents/2015/01/21/2015-00636/long-chain-perfluoroalkyl-carboxylate-and-perfluoroalkyl-sulfonate-chemical-substances-significant>. Accessed 14 Oct 2017
- U.S. EPA (2010) U.S. Environmental Protection Agency (EPA): 2010/2015 PFOA Stewardship Program. Available at: <https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/and-polyfluoroalkyl-substances-pfass-under-tsca#tab-3>. Accessed 14 Oct 2017
- U.S. EPA (2017c) U.S. Environmental Protection Agency (U.S. EPA): Reviewing New Chemicals under the Toxic Substances Control Act (TSCA). Available at: <https://www.epa.gov/reviewing-new-chemicals-under-toxic-substances-control-act-tsca>. Accessed 17 May 2017
- U.S. EPA (2012) U.S. Environmental Protection Agency (U.S. EPA): Long-chain perfluorinated chemicals (LCPFCs) used in carpets. July 27, 2012. Economic and Policy Analysis Branch, Economics, Exposure and Technology Division, Office of Pollution, Prevention, and Toxics, U.S. Environmental Protection Agency, Washington, DC. Available at: http://www.fluoridealert.org/wp-content/uploads/pfcs.carpet.report.epa_.2012.pdf. Accessed 2 Aug 2017
- U.S. EPA (2017d) U.S. Environmental Protection Agency (U.S. EPA): EPA launches cross-agency effort to address PFAS. U.S. Environmental Protection Agency (U.S. EPA): News Releases from U.S. EPA Office of Research and Development (ORD), 12/04/2017. Available at: <https://www.epa.gov/newsreleases/epa-launches-cross-agency-effort-address-pfas>. Accessed 5 Jan 2018
- U.S. EPA (2018) U.S. Environmental Protection Agency (U.S. EPA): EPA toxicologists focus innovative research on PFAS compounds. March 13, 2018. Available at: <https://www.epa.gov/sciencematters/epa-toxicologists-focus-innovative-research-pfas-compounds>. Accessed 11 Apr 2018
- U.S. EPA (2019) U.S. Environmental Protection Agency (EPA): EPA Awards \$6 Million to Research Potential Environmental Impacts of PFAS Substances in Waste Streams. Available at: <https://www.pressreleasepoint.com/print/1583225>. Accessed 26 Sep 2019

- U.S. EPA (2016c) U.S. Environmental Protection Agency (U.S. EPA): Health effects support document for perfluorooctanoic acid (PFOA). Office of Water (4304T), Health and Ecological Criteria Division, Washington, DC, EPA Document Number 822-R-16-003, May 2016. Available at: https://www.epa.gov/sites/production/files/2016-05/documents/pfoa_hesd_final_508.pdf. Accessed 28 Nov 2017
- U.S. EPA (2016d) U.S. Environmental Protection Agency (U.S. EPA): Health effects support document for perfluorooctane sulfonate (PFOS). Office of Water (4304T), Health and Ecological Criteria Division, Washington, DC, EPA Document Number 822-R-16-002, May 2016. Available at: https://www.epa.gov/sites/production/files/2016-05/documents/pfos_hesd_final_508.pdf. Accessed 28 Nov 2017
- U.S. House of Representatives (2018) Text of the House Amendment to the Senate Amendment to H.R. 1625, March 2018. Available at: <http://docs.house.gov/billsthisweek/20180319/BILLS-115SAHR1625-RCP115-66.pdf>. Accessed 9 Apr 2018
- U.S. Senate (2018) A letter from 25 Senators to the Honorable Scott Pruitt, Administrator of the U.S. Environmental Protection Agency concerning groundwater contamination from per- and polyfluoroalkyl substances (PFAS) in communities and on Department of Defense (DOD) installations across their states, April 13, 2018. Available at: https://drive.google.com/file/d/1LgpWUUVI-wfvSW90LtTzjymSNm_BAZTj1/view. Accessed 9 May 2018
- U.S. Senate (2017) S.1519 - National Defense Authorization Act for Fiscal Year 2018 (Report No. 115–125). Available at: <https://www.congress.gov/bill/115th-congress/senate-bill/1519/text>. Accessed 5 Jan 2018
- Valentini A, Taberlet P, Miaud C, et al (2016) Next-generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. *Molecular Ecology* 25:929–942. <https://doi.org/10.1111/mec.13428>
- Valsecchi S, Conti D, Crebelli R, et al (2017) Deriving environmental quality standards for perfluorooctanoic acid (PFOA) and related short chain perfluorinated alkyl acids. *Journal of Hazardous Materials* 323:84–98. <https://doi.org/10.1016/j.jhazmat.2016.04.055>
- Venkatesan AK, Halden RU (2014) Loss and in situ production of perfluoroalkyl chemicals in outdoor biosolids–soil mesocosms. *Environmental Research* 132:321–327. <https://doi.org/10.1016/j.envres.2014.04.024>
- Vermont Department of Health (2016) Perfluorooctanoic acid (PFOA) and perfluorooctanesulfonic acid (PFOS) Vermont drinking water health advisory. Memorandum from Sarah Vose, State Toxicologist to Chuck Schwer, Director, Waste Management, DEC. June 22, 2016. Available at: https://anrweb.vt.gov/PubDocs/DEC/PFOA/PFOA%20-%20PFOS%20Health%20Advisories/Vermont/PFOA_PFOS_HealthAdvisory_June_22_2016.pdf. Accessed 14 Aug 2017

- Verreault J, Berger U, Gabrielsen GW (2007) Trends of perfluorinated alkyl substances in herring gull eggs from two coastal colonies in northern Norway: 1983-2003. *Environmental Science & Technology* 41:6671–6677. <https://doi.org/10.1021/es070723j>
- Vestergren R, Herzke D, Wang T, Cousins IT (2015) Are imported consumer products an important diffuse source of PFASs to the Norwegian environment? *Environmental Pollution* 198:223–230. <https://doi.org/10.1016/j.envpol.2014.12.034>
- Wallington TJ, Hurley MD, Xia J, et al (2006) Formation of C7F15COOH (PFOA) and other perfluorocarboxylic acids during the atmospheric oxidation of 8:2 fluorotelomer alcohol. *Environmental Science & Technology* 40:924–930. <https://doi.org/10.1021/es051858x>
- Wang C, Lu Y, Li Q, et al (2019) Assessing the contribution of atmospheric transport and tourism activities to the occurrence of perfluoroalkyl acids (PFAAs) in an Alpine Nature Reserve. *Science of The Total Environment* 697:133851. <https://doi.org/10.1016/j.scitotenv.2019.133851>
- Wang F, Lu X, Li X, Shih K (2015) Effectiveness and mechanisms of defluorination of perfluorinated alkyl substances by calcium compounds during waste thermal treatment. *Environmental Science & Technology* 49:5672–5680. <https://doi.org/10.1021/es506234b>
- Wang G, Lu J, Xing Z, et al (2017a) Occurrence, distribution, and risk assessment of perfluoroalkyl acids (PFAAs) in muscle and liver of cattle in Xinjiang, China. *International Journal of Environmental Research and Public Health* 14:970. <https://doi.org/10.3390/ijerph14090970>
- Wang T, Lin Z, Yin D, et al (2011) Hydrophobicity-dependent QSARs to predict the toxicity of perfluorinated carboxylic acids and their mixtures. *Environmental Toxicology and Pharmacology* 32:259–265. <https://doi.org/10.1016/j.etap.2011.05.011>
- Wang Y, Yeung LWY, Taniyasu S, et al (2008) Perfluorooctane sulfonate and other fluorochemicals in waterbird eggs from south China. *Environmental Science & Technology* 42:8146–8151. <https://doi.org/10.1021/es8006386>
- Wang Z, Cousins IT, Berger U, et al (2016) Comparative assessment of the environmental hazards of and exposure to perfluoroalkyl phosphonic and phosphinic acids (PFPA and PFPIAs): current knowledge, gaps, challenges and research needs. *Environment International* 89–90:235–247. <https://doi.org/10.1016/j.envint.2016.01.023>
- Wang Z, Cousins IT, Scheringer M, et al (2014a) Global emission inventories for C4–C14 perfluoroalkyl carboxylic acid (PFCA) homologues from 1951 to 2030, Part I: Production and emissions from quantifiable sources. *Environment International* 70:62–75. <https://doi.org/10.1016/j.envint.2014.04.013>
- Wang Z, DeWitt JC, Higgins CP, Cousins IT (2017b) A never-ending story of per- and polyfluoroalkyl substances (PFASs)? *Environmental Science & Technology* 51:2508–2518. <https://doi.org/10.1021/acs.est.6b04806>

- Wang Z, Xie Z, Möller A, et al (2014b) Atmospheric concentrations and gas/particle partitioning of neutral poly- and perfluoroalkyl substances in northern German coast. *Atmospheric Environment* 95:207–213. <https://doi.org/10.1016/j.atmosenv.2014.06.036>
- Washington JW, Ellington JJ, Jenkins TM, et al (2009) Degradability of an acrylate-linked, fluorotelomer polymer in soil. *Environmental Science & Technology* 43:6617–6623. <https://doi.org/10.1021/es9002668>
- Washington JW, Jenkins TM, Rankin K, Naile JE (2015) Decades-scale degradation of commercial, side-chain, fluorotelomer-based polymers in soils and water. *Environmental Science & Technology* 49:915–923. <https://doi.org/10.1021/es504347u>
- Washington JW, Rankin K, Libelo EL, et al (2019) Determining global background soil PFAS loads and the fluorotelomer-based polymer degradation rates that can account for these loads. *Science of The Total Environment* 651:2444–2449. <https://doi.org/10.1016/j.scitotenv.2018.10.071>
- Washington State Legislature (2019) Substitute Senate Bill 5135
- Weinberg I, Dreyer A, Ebinghaus R (2011) Landfills as sources of polyfluorinated compounds, polybrominated diphenyl ethers and musk fragrances to ambient air. *Atmospheric Environment* 45:935–941. <https://doi.org/10.1016/j.atmosenv.2010.11.011>
- Weiss JM, Andersson PL, Lamoree MH, et al (2009) Competitive binding of poly- and perfluorinated compounds to the thyroid hormone transport protein transthyretin. *Toxicological Sciences* 109:206–216. <https://doi.org/10.1093/toxsci/kfp055>
- White ND, Balthis L, Kannan K, et al (2015) Elevated levels of perfluoroalkyl substances in estuarine sediments of Charleston, SC. *Science of The Total Environment* 521–522:79–89. <https://doi.org/10.1016/j.scitotenv.2015.03.078>
- Wilkinson D (2016) Letter from Mr. David Wilkinson of Tandus Centiva to DTSC Safer Consumer Products Branch staff via email
- Wilkinson JL, Hooda PS, Swinden J, et al (2017) Spatial distribution of organic contaminants in three rivers of Southern England bound to suspended particulate material and dissolved in water. *Science of The Total Environment* 593–594:487–497. <https://doi.org/10.1016/j.scitotenv.2017.03.167>
- Winkens K, Koponen J, Schuster J, et al (2017a) Perfluoroalkyl acids and their precursors in indoor air sampled in children’s bedrooms. *Environmental Pollution* 222:423–432. <https://doi.org/10.1016/j.envpol.2016.12.010>
- Winkens K, Vestergren R, Berger U, Cousins IT (2017b) Early life exposure to per- and polyfluoroalkyl substances (PFASs): a critical review. *Emerging Contaminants* 3:55–68. <https://doi.org/10.1016/j.emcon.2017.05.001>

- Winqvist A, Steenland K (2014) Perfluorooctanoic acid exposure and thyroid disease in community and worker cohorts. *Epidemiology* 25:255–264. <https://doi.org/10.1097/EDE.0000000000000040>
- Wolf CJ, Rider CV, Lau C, Abbott BD (2014) Evaluating the additivity of perfluoroalkyl acids in binary combinations on peroxisome proliferator-activated receptor- α activation. *Toxicology* 316:43–54. <https://doi.org/10.1016/j.tox.2013.12.002>
- Wolf CJ, Takacs ML, Schmid JE, et al (2008) Activation of mouse and human peroxisome proliferator-activated receptor alpha by perfluoroalkyl acids of different functional groups and chain lengths. *Toxicological Sciences* 106:162–171. <https://doi.org/10.1093/toxsci/kfn166>
- Wong YM, Li R, Lee CKF, et al (2017) The measurement of bisphenol A and its analogues, perfluorinated compounds in twenty species of freshwater and marine fishes, a time-trend comparison and human health based assessment. *Marine Pollution Bulletin* 124:743–752. <https://doi.org/10.1016/j.marpolbul.2017.05.046>
- Wu T, Täubel M, Holopainen R, et al (2018) Infant and adult inhalation exposure to resuspended biological particulate matter. *Environmental Science & Technology* 52:237–247. <https://doi.org/10.1021/acs.est.7b04183>
- Wu X (May), Bennett DH, Calafat AM, et al (2015) Serum concentrations of perfluorinated compounds (PFC) among selected populations of children and Adults in California. *Environmental Research* 136:264–273. <https://doi.org/10.1016/j.envres.2014.09.026>
- Xiao F (2017) Emerging poly- and perfluoroalkyl substances in the aquatic environment: a review of current literature. *Water Research* 124:482–495. <https://doi.org/10.1016/j.watres.2017.07.024>
- Xiao F, Simcik MF, Halbach TR, Gulliver JS (2015) Perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) in soils and groundwater of a U.S. metropolitan area: Migration and implications for human exposure. *Water Research* 72:64–74. <https://doi.org/10.1016/j.watres.2014.09.052>
- Xie Z, Wang Z, Mi W, et al (2015) Neutral poly-/perfluoroalkyl substances in air and snow from the Arctic. *Scientific Reports* 5:8912. <https://doi.org/10.1038/srep08912>
- Xu Z, Fiedler S, Pfister G, et al (2013) Human exposure to fluorotelomer alcohols, perfluorooctane sulfonate and perfluorooctanoate via house dust in Bavaria, Germany. *Science of The Total Environment* 443:485–490. <https://doi.org/10.1016/j.scitotenv.2012.10.089>
- Yamada A, Bemrah N, Veyrand B, et al (2014a) Dietary exposure to perfluoroalkyl acids of specific French adult sub-populations: high seafood consumers, high freshwater fish consumers and pregnant women. *Science of The Total Environment* 491–492:170–175. <https://doi.org/10.1016/j.scitotenv.2014.01.089>

- Yamada A, Bemrah N, Veyrand B, et al (2014b) Perfluoroalkyl acid contamination and polyunsaturated fatty acid composition of French freshwater and marine fishes. *Journal of Agricultural and Food Chemistry* 62:7593–7603. <https://doi.org/10.1021/jf501113j>
- Yan H, Zhang CJ, Zhou Q, et al (2012) Short- and long-chain perfluorinated acids in sewage sludge from Shanghai, China. *Chemosphere* 88:1300–1305. <https://doi.org/10.1016/j.chemosphere.2012.03.105>
- Yang L, Wang Z, Shi Y, et al (2016) Human placental transfer of perfluoroalkyl acid precursors: levels and profiles in paired maternal and cord serum. *Chemosphere* 144:1631–1638. <https://doi.org/10.1016/j.chemosphere.2015.10.063>
- Yao Y, Zhao Y, Sun H, et al (2018) Per- and polyfluoroalkyl substances (PFASs) in indoor air and dust from homes and various microenvironments in China: implications for human exposure. *Environmental Science & Technology* 52:3156–3166. <https://doi.org/10.1021/acs.est.7b04971>
- Ye F, Zushi Y, Masunaga S (2015) Survey of perfluoroalkyl acids (PFAAs) and their precursors present in Japanese consumer products. *Chemosphere* 127:262–268. <https://doi.org/10.1016/j.chemosphere.2015.02.026>
- Ye X, Strynar MJ, Nakayama SF, et al (2008) Perfluorinated compounds in whole fish homogenates from the Ohio, Missouri, and Upper Mississippi Rivers, USA. *Environmental Pollution* 156:1227–1232. <https://doi.org/10.1016/j.envpol.2008.03.014>
- Yeh A, Marcinek DJ, Meador JP, Gallagher EP (2017) Effect of contaminants of emerging concern on liver mitochondrial function in Chinook salmon. *Aquatic Toxicology* 190:21–31. <https://doi.org/10.1016/j.aquatox.2017.06.011>
- Yeung LWY, Mabury SA (2016) Are humans exposed to increasing amounts of unidentified organofluorine? *Environmental Chemistry* 13:102. <https://doi.org/10.1071/EN15041>
- Yoo H, Washington JW, Jenkins TM, Ellington JJ (2011) Quantitative determination of perfluorochemicals and fluorotelomer alcohols in plants from biosolid-amended fields using LC/MS/MS and GC/MS. *Environmental Science & Technology* 45:7985–7990. <https://doi.org/10.1021/es102972m>
- Young CJ, Furdui VI, Franklin J, et al (2007) Perfluorinated acids in Arctic snow: new evidence for atmospheric formation. *Environmental Science & Technology* 41:3455–3461. <https://doi.org/10.1021/es0626234>
- Yu N, Guo H, Yang J, et al (2018) Non-target and suspect screening of per- and polyfluoroalkyl substances in airborne particulate matter in China. *Environmental Science & Technology* 52:8205–8214. <https://doi.org/10.1021/acs.est.8b02492>

- Zareitalabad P, Siemens J, Hamer M, Amelung W (2013) Perfluorooctanoic acid (PFOA) and perfluorooctanesulfonic acid (PFOS) in surface waters, sediments, soils and wastewater – a review on concentrations and distribution coefficients. *Chemosphere* 91:725–732. <https://doi.org/10.1016/j.chemosphere.2013.02.024>
- Zhang D, Zhang W, Liang Y (2019a) Distribution of eight perfluoroalkyl acids in plant-soil-water systems and their effect on the soil microbial community. *Science of The Total Environment* 697:134146. <https://doi.org/10.1016/j.scitotenv.2019.134146>
- Zhang H, Liu W, He X, et al (2015) Uptake of perfluoroalkyl acids in the leaves of coniferous and deciduous broad-leaved trees. *Environmental Toxicology and Chemistry* 34:1499–1504. <https://doi.org/10.1002/etc.2968>
- Zhang L, Sun H, Wang Q, et al (2019b) Uptake mechanisms of perfluoroalkyl acids with different carbon chain lengths (C2–C8) by wheat (*Triticum aestivum* L.). *Science of The Total Environment* 654:19–27. <https://doi.org/10.1016/j.scitotenv.2018.10.443>
- Zhang R, Ye J, Wei Q, et al (2018) Plasma concentration of 14 perfluoroalkyl acids (PFAAs) among children from seven cities in Guangdong, China. *Science of The Total Environment* 616–617:1469–1476. <https://doi.org/10.1016/j.scitotenv.2017.10.167>
- Zhang T, Sun H, Lin Y, et al (2013) Distribution of poly- and perfluoroalkyl substances in matched samples from pregnant women and carbon chain length related maternal transfer. *Environmental Science & Technology* 47:7974–7981. <https://doi.org/10.1021/es400937y>
- Zhang Y, Tan D, Geng Y, et al (2016) Perfluorinated compounds in greenhouse and open agricultural producing areas of three provinces of China: Levels, sources and risk assessment. *International Journal of Environmental Research and Public Health* 13:1224. <https://doi.org/10.3390/ijerph13121224>
- Zhao H, Guan Y, Qu B (2018) PFCA uptake and translocation in dominant wheat species (*Triticum aestivum* L.). *International Journal of Phytoremediation* 20:68–74. <https://doi.org/10.1080/15226514.2017.1337066>
- Zhao L, Zhang Y, Zhu L, et al (2017) Isomer-specific transplacental efficiencies of perfluoroalkyl substances in human whole blood. *Environmental Science & Technology Letters* 4:391–398. <https://doi.org/10.1021/acs.estlett.7b00334>
- Zhao S, Zhu L (2017) Uptake and metabolism of 10:2 fluorotelomer alcohol in soil-earthworm (*Eisenia fetida*) and soil-wheat (*Triticum aestivum* L.) systems. *Environmental Pollution* 220:124–131. <https://doi.org/10.1016/j.envpol.2016.09.030>
- Zhao Z, Xie Z, Möller A, et al (2012) Distribution and long-range transport of polyfluoroalkyl substances in the Arctic, Atlantic Ocean and Antarctic coast. *Environmental Pollution* 170:71–77. <https://doi.org/10.1016/j.envpol.2012.06.004>

- Zucker HA, Butler J, Lyon N, et al (2017) Letter to Brenda Fitzgerald, M.D., Director, Centers for Disease Control and Prevention, Administrator, Agency for Toxic Substances and Disease Registry, US Department of Health & Human Services, August 24, 2017. Available at: https://www.health.ny.gov/environmental/investigations/drinkingwaterresponse/docs/2017-08-24_cdc_study_letter.pdf. Accessed 15 Sep 2017
- Zushi Y, Hogarh JN, Masunaga S (2012) Progress and perspective of perfluorinated compound risk assessment and management in various countries and institutes. *Clean Technologies and Environmental Policy* 14:9–20. <https://doi.org/10.1007/s10098-011-0375-z>
- Zushi Y, Masunaga S (2009) Identifying the nonpoint source of perfluorinated compounds using a geographic information system based approach. *Environmental Toxicology and Chemistry* 28:691–700

APPENDIX 1 – ADDITIONAL PFAS CANDIDATE CHEMICALS

DATABASE LISTINGS

Chemical Name	CAS RN	Authoritative List	Hazard Traits
10:2 Fluorotelomer acetate	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
10:2 Fluorotelomer acrylate	17741-60-5	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
10:2 Fluorotelomer carboxylic acid	53826-13-4	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
10:2 Fluorotelomer unsaturated carboxylic acid	70887-94-4	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
2-(N-Ethyl-perfluorooctane sulfonamido) acetic acid (Et-PFOA-AcOH)	2991-50-6	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
2-(N-Methyl-perfluorooctane sulfonamido) acetic acid (Me-PFOA-AcOH)	2355-31-9	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity

Chemical Name	CAS RN	Authoritative List	Hazard Traits
2,3,3,3-Tetrafluoro-2-(1,1,2,2,3,3,3-heptafluoropropoxy) propanoic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
4:2 Fluorotelomer sulfonic acid	757124-72-4	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
5:3 Fluorotelomer carboxylic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
6:2 Fluorotelomer acetate	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
6:2 Fluorotelomer acrylate	17527-29-6	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
6:2 Fluorotelomer carboxylic acid	53826-12-3	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
6:2 Fluorotelomer phosphate diester	57677-95-9	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity

Chemical Name	CAS RN	Authoritative List	Hazard Traits
6:2 Fluorotelomer phosphate monoester	57678-01-0	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
6:2 Fluorotelomer sulfonic acid	27619-97-2	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
6:2 Fluorotelomer unsaturated carboxylic acid	70887-88-6	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
6:2/8:2 Fluorotelomer phosphate diester	943913-15-3	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
7:3 Fluorotelomer carboxylic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
7:3 Fluorotelomer carboxylic acid	No CAS RN	ATSDR Neurotoxicants (3/3/2011)	Immunotoxicity
8:2 Fluorotelomer acetate	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity

Chemical Name	CAS RN	Authoritative List	Hazard Traits
8:2 Fluorotelomer acrylate	27905-45-9	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
8:2 Fluorotelomer carboxylic acid	27854-31-5	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
8:2 Fluorotelomer phosphate diester	678-41-1	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
8:2 Fluorotelomer sulfonic acid	39108-34-4	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
8:2 Fluorotelomer unsaturated carboxylic acid	70887-84-2	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Ammonium 4,8-dioxo-3H-perfluorononanoate (ADONA)	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Ammonium nonadecafluorodecanoate	3108-42-7	EC PBTs (1/12/2017)	Bioaccumulation, Environmental Persistence
Ammonium pentadecafluorooctanoate (APFO)	3825-26-1	EC PBTs (1/12/2017)	Bioaccumulation, Environmental Persistence

Chemical Name	CAS RN	Authoritative List	Hazard Traits
Ammonium pentadecafluorooctanoate (APFO)	3825-26-1	EC Annex VI CMRs – Cat. 1B (7/19/2016)	Reproductive Toxicity
Bis(perfluorohexyl) phosphinic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Bis(perfluorooctyl) phosphinic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Sum of branched perfluorooctanoic isomers	No CAS RN	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Heptacosafuorotetradecanoic acid	376-06-7	EC PBTs (1/12/2017)	Bioaccumulation, Environmental Persistence
Nonadecafluorodecanoic acid, sodium salt	3830-45-3	EC PBTs (1/12/2017)	Bioaccumulation, Environmental Persistence
n-Perfluorooctane sulfonic acid (n-PFOS) and n-Perfluorooctanoic acid (n-PFOA)	No CAS RN	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Pentacosafuorotridecanoic acid	72629-94-8	EC PBTs (1/12/2017)	Bioaccumulation, Environmental Persistence
Perfluorobutane sulfonic acid (PFBS)	375-73-5	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Perfluorodecanoic acid (PFDeA)	335-76-2	EC PBTs (1/12/2017)	Bioaccumulation, Environmental Persistence
Perfluorodecanoic acid (PFDeA)	335-76-2	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Perfluorodecylphosphonic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity

Chemical Name	CAS RN	Authoritative List	Hazard Traits
Perfluorododecanoic acid (PFDoA)	307-55-1	EC PBTs (1/12/2017)	Bioaccumulation, Environmental Persistence
Perfluorododecanoic acid (PFDoA)	307-55-1	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Perfluoroethylcyclohexane sulfonic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluoroheptane sulfonic acid	375-92-8	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluoroheptanoic acid (PFHpA)	375-85-9	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Perfluorohexadecanoic acid	67905-19-5	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluorohexane sulfonic acid (PFHxS)	355-46-4	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Perfluorohexanoic acid (PFHxA)	307-24-4	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluorohexyl-perfluorooctylphosphinic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity

Chemical Name	CAS RN	Authoritative List	Hazard Traits
Perfluorohexylphosphonic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Sum of perfluoromethylheptane sulfonic acid isomers	No CAS RN	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Perfluorononane sulfonic acid	474511-07-4	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluorononanoic acid (PFNA)	375-95-1	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Perfluorononanoic acid (PFNA)	375-95-1	EC Annex VI CMRs – Cat. 1B (7/19/2016)	Reproductive Toxicity
Perfluorooctadecanoic acid	16517-11-6	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluorooctane sulfonamide (PFOSA)	754-91-6	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Perfluorooctane sulfonic acid (PFOS)	1763-23-1	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity
Perfluorooctane sulfonic acid (PFOS)	1763-23-1	EC Annex VI CMRs – Cat. 1B (7/19/2016)	Reproductive Toxicity
Perfluorooctane sulfonic acid (PFOS)	1763-23-1	OSPAR Priority Action Part A (1/1/2002)	Bioaccumulation, Environmental Persistence
Perfluorooctane sulfonic acid (PFOS)	1763-23-1	WA PBT (1/3/2006)	Bioaccumulation, Environmental Persistence
Perfluorooctanoic acid (PFOA)	335-67-1	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity

Chemical Name	CAS RN	Authoritative List	Hazard Traits
Perfluorooctanoic acid (PFOA)	335-67-1	EC Annex VI CMRs – Cat. 1B (7/19/2016)	Reproductive Toxicity
Perfluorooctanoic acid (PFOA)	335-67-1	EC PBTs (1/12/2017)	Bioaccumulation, Environmental Persistence
Perfluorooctanoic acid (PFOA)	335-67-1	IARC Carcinogens – 2B (5/19/2017)	Carcinogenicity
Perfluorooctylphosphonic acid	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluoropentane sulfonic acid	2706-91-4	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluoropentanoic acid	2706-90-3	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluorotetradecanoic acid	376-06-7	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluorotridecanoic acid	72629-94-8	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Perfluoroundecanoic acid (PFUA)	2058-94-8	CDC 4 th National Exposure Report (1/1/2017)	Developmental Toxicity, Hepatotoxicity, Digestive System Toxicity, Reproductive Toxicity

Chemical Name	CAS RN	Authoritative List	Hazard Traits
Potassium 1,1,2,2,-tetrafluoro-2-(oxy)ethanesulfonate tetrafluoroethene	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Sodium 2-(N-ethylperfluorooctane-1-sulfonamido)ethyl phosphate	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity
Sodium bis-[2-(N-ethylperfluorooctane-1-sulfonamido)ethyl] phosphate	No CAS RN	CECBP - Priority Chemicals (12/22/2015) Priority Chemicals that are identified under the California Environmental Contaminant Biomonitoring Program	Bioaccumulation, Carcinogenicity, Developmental Toxicity, Endocrine Toxicity, Hepatotoxicity, Digestive System Toxicity, Immunotoxicity, Reproductive Toxicity

APPENDIX 2 – PFAS HAZARD TRAITS

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
Carcinogenicity	Limited evidence in humans for the carcinogenicity of perfluorooctanoic acid (PFOA). A positive association was observed for cancers of the testis and kidney. Overall evaluation: Perfluorooctanoic acid (PFOA) is possibly carcinogenic to humans (Group 2B). Review by authoritative body.	Positive	IARC (2016)
	Suggestive evidence of carcinogenic potential for PFOA. Epidemiology studies demonstrate an association of serum PFOA with kidney and testicular tumors among highly exposed members of the general population. Two chronic bioassays of PFOA support a positive finding for its ability to be tumorigenic in one or more organs of rats, including the liver, testes, and pancreas. Review by authoritative body.	Positive	U.S. EPA (2016a)
	Suggestive evidence of carcinogenic potential for PFOS. In a chronic oral toxicity and carcinogenicity study of PFOS in rats, liver and thyroid tumors, and mammary fibroadenomas were identified. Review by authoritative body.	Positive	U.S. EPA (2016b)
	Positive association of kidney cancer with PFOA exposure as observed in high PFOA exposure cohorts in occupational and community settings.	Positive	Barry et al. (2013); C8 Science Panel (2012); Steenland and Woskie (2012); Vieira et al. (2013)
	Positive association of testicular cancer with PFOA exposure in high PFOA exposure cohort.	Positive	Barry et al. (2013); C8 Science Panel (2012)
	Increased risk of thyroid cancer in occupational cohort exposure to PFOA.	Positive	Barry et al. (2013)
	Possible increase in prostate cancer mortality in workers with 10 years occupational exposure to PFOA.	Positive	Gilliland and Mandel (1993)
	Higher PFOA serum levels associated with testicular, kidney, prostate and ovarian cancer, and non-Hodgkin lymphoma. (High PFOA exposure cohort)	Positive	Lundin et al. (2009); Vieira et al. (2013)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
—	Higher risk of prostate cancer in individuals with PFAS levels above median and a first degree relative with prostate cancer. Positive of genetic susceptibility. PFHxS showed a statistically significant interaction.	Positive	Hardell et al. (2014)
—	Weak positive association of breast cancer risk with PFOS exposure in Danish National Birth Cohort.	Positive	Bonfeld-Jørgensen et al. (2014)
—	Tumor triad in male rats (liver, Leydig cells, pancreatic) dosed with APFO.	Positive	Biegel et al. (2001)
—	Ammonium 2,3,3,3-tetrafluoro-2-(heptafluoropropoxy)-propanoate was tested for chronic toxicity and carcinogenicity in Sprague–Dawley rats. Level of evidence sufficient to show that the findings of liver, pancreas, testicular tumors are relevant for humans.	Positive	Beekman et al. (2016); Rae et al. (2015)
—	General Danish population study did not indicate increased incidence of prostate, bladder, pancreatic, liver cancers.	Negative	Eriksen et al. (2009)
—	No association between PFOA levels and breast cancer risk in mothers from Danish National Birth Cohort.	Negative	Bonfeld-Jørgensen et al. (2014)
—	No increased prostate cancer risk compared to controls (Gleason score and prostate-specific antigen (PSA) levels).	Negative	Hardell et al. (2014)
—	No association of occupational APFO exposure with liver, pancreatic or testicular cancers.	Negative	Lundin et al. (2009)
—	Study reports lack of concordance between community exposures and occupational exposures one or two magnitudes higher than those for the general population. The discrepant findings across the study populations were described as likely due to chance, confounding, and/or bias.	Negative	Chang et al. (2014); (3M-funded independent tox/epi review)
—	No evidence of genotoxicity from NaPFHx in bacterial mutagenicity assays (333-5000 µg/mL), nor chromosome aberrations in human peripheral lymphocytes (with or without metabolic activation) (5-3860 µg/mL).	Negative	Loveless et al. (2009)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	No evidence of carcinogenicity in either male or female Sprague-Dawley rats treated with PFHxA when administered orally daily for 2 years.	Negative	Klaunig et al. (2015)
	PFASs modestly produce reactive oxygen species and DNA damage in the HepG2 cell line. Increase in ROS production was not concentration-dependent and the compounds did not generate DNA damage that could be detected by the alkaline comet assay.	Negative	Eriksen et al. (2010)
Developmental Toxicity	PFOA and PFOS are “known to the State of California to cause reproductive toxicity”, based on formal identification by an authoritative body (U.S. EPA). Review by authoritative body.	Positive	OEHHA (2017)
	The EPA Office of Water selected a RfD of 0.00002 mg/kg-day for PFOA based on reduced ossification and accelerated puberty (in male mice) (Lau et al. 2006).	Positive	U.S. EPA (2016a)
	EPA derived a reference dose for PFOS of 0.00002 mg/kg-day based on decreased neonatal rat body weight from the 2-generation study (Luebker et al. 2005).	Positive	U.S. EPA (2016b)
	Treatment of pregnant CD-1 mice with NH4+ PFBA by oral gavage is associated with delayed eye opening in pups and delayed vaginal opening in 175 or 350 mg/kg-day dose groups.	Positive	Das et al. (2008)
	Decreased birth weight with increased maternal/cord blood PFOA levels in general population.	Positive	Apelberg et al. (2007); Fei et al. (2007); Johnson et al. (2014); Maisonet et al. (2012)
	No association between PFOA levels and birth weight in high-exposure C8 community population.	Negative	Darrow et al. (2013); Nolan et al. (2009); Savitz et al. (2012a); Savitz et al. (2012b); Stein et al. (2009)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	K+ PFHxS given to Sprague Dawley rats by oral gavage. No treatment-related effects (reproductive, developmental, neurological) in dams or offspring compared with controls.	Negative	Butenhoff et al. (2009)
	2,3,3,3-Tetrafluoro-2-(heptafluoropropoxy)propanoate (a PFOA replacement technology) in rats at 100 and 1000 mg/kg-day decreases fetal weights and produces early delivery of offspring. Offspring were alive, with no increase in resorptions. Reproductive effects were observed at dose levels also inducing maternal toxicity, which “do not normally warrant classification.”	Negative	Beekman et al. (2016)
Reproductive Toxicity	Decreased fertility in humans with increased concentrations of PFOS, PFOA and PFHxS in maternal plasma.	Positive	Fei et al. (2009); Velez et al. (2015)
	Probable link between PFOA exposure and pregnancy-induced hypertension in humans.	Positive	C8 Science Panel (2012)
	PFOS, PFDoA, PFNA, and PFOA are cytotoxic in JEG-3 human placental cells (EC50s range from 107 - 647 µM). PFOS, PFOA, and PFBS are potential CYP19 aromatase inhibitors (IC50s 57 - 80 µM). Notable inhibitory effect of PFBS with relatively low uptake in cells.	Positive	Gorrochategui et al. (2014)
	Treatment of pregnant CD-1 mice with PFBA by oral gavage was associated with increased incidence of fetal resorption and increased offspring liver weight at postnatal day (PND) 1 (but not by PND 10). Delayed eye opening in pups was observed in all dose groups (35, 175 and 350 mg/kg-day) and delayed puberty in highest two dose groups.	Positive	Das et al. (2008)
	Increased ovarian follicular fluid levels of PFHxS were associated with reduced baseline follicle count and post-fertilization success in women receiving assistive reproductive technology.	Positive	McCoy et al. (2017)
	Probable link between PFOA exposure and elevated cholesterol.	Positive	C8 Science Panel (2012)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
Cardiovascular Toxicity	No increase in relative risk of heart disease. Significant 'trend' for increased risk in heart disease after 10-year lag in PFOA exposure.	Positive	Sakr et al. (2009)
	Positive correlation of serum cholesterol with PFOA, PFOS and PFHxS exposure.	Positive	Fisher et al. (2013); Costa et al. (2009); Nelson et al. (2010); Sakr et al. (2007b); Skuladottir et al. (2015); Steenland et al. (2010); Wang et al. (2012); Winqvist and Steenland (2014)
	No indication of increased serum lipids/cholesterol in humans with elevated PFOS/PFOA serum levels.	Negative	Emmett et al. (2006) (exposed residents); Olsen et al. (2000); Olsen and Zobel (2007)
	No electrocardiogram (EKG) alterations in PFOA-exposed workers (5-9550 ng/mL serum PFOA).	Negative	Sakr et al. (2007a); Sakr et al. (2007b)
	No increased mortality ratio in exposed workers for heart disease.	Negative	Lau et al. (2007); Lundin et al. (2009); Steenland and Woskie (2012)
Endocrine Toxicity	Probable link between PFOA exposure and thyroid disease in human based on occupational and community cohort studies.	Positive	C8 Science Panel (2012); Lopez-Espinosa et al. (2012); Melzer et al. (2010); Winqvist and Steenland (2014b)
	Increased risk of gestational diabetes development with serum PFOA	Positive	Zhang et al. (2015)
	Increased mortality from diabetes in occupational exposure cohorts. Authors specify that mortality may not be a good surrogate for incidence.	Positive	Lau et al. (2007); Leonard et al. (2008); Lundin et al. (2009) (SMR = 197); Steenland and Woskie (2012) (SMR = 1.90)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	Positive association of serum estradiol and testosterone with serum PFOA in men. Authors indicated circadian variations (collection time) may confound data.	Positive	Sakr et al. (2007a); Sakr et al. (2007b)
	Decreased free and total T4 in PFOA-treated Cynomolgus monkeys.	Positive	Butenhoff et al. (2002)
	PFBS (IC50 = 68 µM) and PFHxS are aromatase inhibitors in placental cells. Inhibitory effect of PFBS important despite low uptake in cells.	Positive	Gorrochategui et al. (2014)
	Activation of estrogen receptor in reporter gene assay for 4:2, 6:2 and 8:2 FTOH. C4+ PFCAs all activate PPAR-alpha; no PFBA activation of PPAR-gamma.	Positive	Rosenmai et al. (2016)
	PPAR-alpha activities are induced at lower cellular concentrations for short-chain homologs relative to long-chains.	Positive	Rosenmai et al. (2016)
	PFHxSK inhibits 11 -HSD2 in human and rat kidney microsomes. PFHxSK is more potent in human than rat microsomes.	Positive	Zhao et al. (2010)
	Non-cytotoxic levels of PFHxDI (dodecafluoro-1,6-diiodohexane) significantly promote the proliferation of MCF-7 breast cancer cells. PFHxDI's preferential binding affinity for estrogen receptor α and β isoforms mediated this cellular response.	Positive	Song et al. (2018)
	PFHxS can disrupt the thyroid system of Wistar rats, as measured by T4 levels in dams and offspring. PFHxS potentiated the antiandrogenic effect of an endocrine disruptor mixture in male offspring.	Positive	Ramhøj et al. (2018)
	Competitive binding to transthyretin (for TH transport) in binding assay, reducing free thyroxine in blood. Competitive binding of PFHxS > PFOS/PFOA > PFHpA. Binding of PFASs 12-300 times lower than thyroxine.	Positive	Weiss et al. (2009)
	No effect from PFBA and FTOHs.	Negative	

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	PFBA/PFBS had lowest potency of activation of PPAR-alpha in COS-1 cells in multiple chain lengths tested.	Negative	Wolf et al. (2008)
	The potassium salts of PFBS and PFHxS has no effect on 3β or 17β-HSD activity in human or rat testes microsomes, even at high concentrations.	Negative	Zhao et al. (2010)
	No evidence of altered thyroid function with occupational exposure, serum thyroid stimulating hormone (TSH), T3, T4.	Negative	Mundt et al. (2007); Olsen et al. (1998); Olsen et al. (2003); Olsen and Zobel (2007); Sakr et al. (2007a); Sakr et al. (2007b)
	No association between serum PFOA and Type 2 diabetes incidence in general or worker populations.	Negative	MacNeil et al. (2009); Steenland et al. (2015)
	No association between serum PFOA and measures of metabolic syndrome in general or worker populations.	Negative	Lin et al. (2009)
Hemato toxicity	Reduced red blood cell (RBC) count, hemoglobin and hematocrit in male rats (90 day PFBS dosage).	Positive	Lieder et al. (2009)
	No correlation in hematological parameters (including hematocrit, hemoglobin, red blood cells, white blood cells, platelets) and serum PFOS in plant workers.	Negative	Olsen et al. (2003); Olsen et al. (1998); Olsen and Zobel (2007)
	No alterations in blood counts in workers exposure to PFOA.	Negative	Sakr et al. (2007a); Sakr et al. (2007b)
Hepatotoxicity and Digestive System Toxicity	Positive minor association of serum PFOA with serum liver enzyme levels (AST, ALT, GGT).	Positive	Costa et al. (2009); Gallo et al. (2012); Olsen et al. (2000); Olsen et al. (2003); Olsen and Zobel (2007); Sakr et al. (2007a); Sakr et al. (2007b)
	Probable link between PFOA exposure and ulcerative colitis.	Positive	C8 Science Panel (2012); Steenland et al. (2015)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	Small, increased liver weight in Cynomolgous monkeys (PFOA capsules).	Positive (low N, 1-2% increase)	Butenhoff et al. (2002)
	Hepatonecrosis, hepatic tumors in Cynomolgous monkeys. At termination of dosing, the mean PFOS serum level corresponding to the LOAEL is 171,000 ng/mL in females and 173,000 ng/mL in males (K-PFOS capsules).	Positive (low N)	Seatcat et al. (2002)
	In rats, PFHxS is more toxic to liver than PFOS and PFBS.	Positive	Danish Environmental Protection Agency (2015)
	In a 90-day study, PFHxA was administered to rats daily by oral gavage. The NOAEL is 50 mg/kg/day for males and >200 mg/kg/day for females based on increased liver weight, peroxisome beta oxidation activity and centrilobular hepatocyte hypertrophy.	Positive	Chengelis et al. (2009)
	Reversible increased liver weight at 30 and 150 mg/kg-day APFB, slight hepatocellular hypertrophy and decreased serum total cholesterol in male S-D rats at 150 mg/kg-day for 28 days APFB.	Positive	Butenhoff et al. (2012)
	Various repeated dose studies of varying duration (but 90 days or less) and dose in rodents of FRD-902, ammonium 2,3,3,3-tetrafluoro-2-(heptafluoropropoxy)propanoate revealed increased liver and kidney weights, hepatocellular hypertrophy and alterations in blood chemistries. Chronic feeding study revealed various liver pathologies.	Positive	Beekman et al. (2016)
	Increased incidence of focal cystic degeneration, and centrilobular hepatocellular hypertrophy and necrosis in male Sprague-Dawley rats given 50 mg/kg-day ammonium 2,3,3,3-tetrafluoro-2-(heptafluoropropoxy)-propanoate orally for 2 years. Similar results in female rats given 500 mg/kg-day. Clinical chemistry evaluations among males receiving 50 mg/kg-d showed increases in enzymes indicative of liver injury (alkaline phosphatase, alanine aminotransferase, aspartate aminotransferase, and sorbitol dehydrogenase).	Positive	Rae et al. (2015)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	K+ PFHxS to Sprague Dawley rats by oral gavage. Effects in parental males include reduced serum cholesterol (all doses), increased liver-to-body and liver-to brain weight ratio, centrilobular hypertrophy, hypertrophy AND hyperplasia of thyroid follicular cells, decreased hematocrit (3 and 10 mg/kg b.w. per day), decreased triglycerides and increased albumin, urea nitrogen, alkaline phosphatase, Ca ²⁺ and albumin/globulin ratio (10 mg/kg b.w. per day)	Positive	Butenhoff et al. (2009)
	4:2, 6:2 and 8:2 FTOHs were incubated in freshly isolated rat hepatocytes, causing cell and mitochondrial toxicity. Cytotoxicity is related to cytochrome P450-mediated bioactivation. LC50s 0.66, 3.7 and 1.4 mM, respectively.	Positive	Martin et al. (2009)
	Compared with PFOA and PFOS, long-chain PFAS alternatives 6:2 chlorinated polyfluorinated ether sulfonate, hexafluoropropylene trimer acid, HFPO tetramer acid, and 6:2 fluorotelomer sulfonic acid show greater cytotoxicity on human liver HL-7702 cells.	Positive	Sheng et al. (2017)
	6:2 chlorinated polyfluorinated ether sulfonate (6:2 Cl-PFESA) and hexafluoropropylene oxide (HFPO) show unique binding modes and higher binding affinity to human liver fatty acid binding protein (hl-FABP), and higher cytotoxicity in human liver cells, than PFOA and PFOS.	Positive	Sheng et al. (2018)
	PPAR-alpha induction in human hepatocellular carcinoma cells is similar for C4 and C5 ("short-chain") PFCAs as for C12 and C14 ("long-chain"), but is induced at lower cellular concentrations by the C4 and C5 compounds tested. Other PFCAs tested (C6-C10) induce PPAR-alpha greater than 2.5-fold.	Positive	Rosenmai et al. (2018)
	No significant relationship between serum PFOA and liver function in residents of community water district.	Negative	Emmett et al. (2006)
	No association of PFOA exposure with liver disease. The observed U-shaped dose-response for serum bilirubin may explain inverse responses in occupational cohorts.	Negative	C8 Science Panel (2012)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	No reported gastric ulcers or colon polyps in workers exposed to PFOS.	Negative	Grice et al. (2007)
Immunotoxicity	Several PFASs were associated with an increased number of respiratory tract infections in the first 10 years of life, suggesting immunosuppressive effects.	Positive	Impinen et al. (2018)
	Depressed antibody production post-vaccination, or increased odds of, with increasing PFOA, PFOS and PFHxS serum concentrations.	Positive	Grandjean et al. (2012); Grandjean et al. (2016); Granum et al. (2013); Looker et al. (2014)
	In human peripheral blood leukocytes, PFBS inhibits the release of TNF- α and IL-10, but IL-6 and IFN- γ are unaffected. In THP-1 cells, PFBS also inhibits NF- κ B activation (by inhibiting LPS-induced phosphorylation of P65, necessary for NF- κ B transcription), and prevents I- κ B degradation.	Positive	Corsini et al. (2012)
	Probable link between PFOA exposure and ulcerative colitis.	Positive	C8 Science Panel (2012); Steenland et al. (2015)
	Prenatal exposure to PFOA, PFDA, PFDoA and PFHxS increases the risk of childhood atopic dermatitis in female children during the first 24 months of life in a prospective birth cohort study.	Positive	Chen et al. (2018)
	Oral administration of 2,3,3,3-Tetrafluoro-2-(Heptafluoropropoxy)-Propanoate in mice for 28 days. Evaluated for T-cell dependent antibody response (TDAR) and splenic lymphocyte subpopulations 5 days post-antigen treatment. TDAR is suppressed in females at 100 mg/kg. T lymphocyte numbers are increased in males at 100 mg/kg.	Positive	Rushing et al. (2017)
Urinary System Toxicity	Increased mortality from renal disease (SMR 3.11) in workers at PFOA facility. PFOA exposure estimated based on job history and data from biomonitoring survey. Significant positive trend in nonmalignant kidney disease.	Positive	Steenland and Woskie (2012)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
Nephrotoxicity and other	No association of kidney function, (measured blood urea nitrogen and serum creatinine) with occupational PFOS/PFOA exposure.	Negative	Costa et al. (2009); Mundt et al. (2007); Olsen et al. (2003); Olsen et al. (1998)
Neurodevelopmental Toxicity	Positive association with Attention Deficit Hyperactivity Disorder (ADHD) in children in a highly PFOA-exposed community.	Positive	Hoffman et al. (2010)
	A single exposure to PFHxS (9.2 mg/kg) on postnatal day 10 affected the cholinergic system and altered adult spontaneous behavior and cognitive function in male and female mice.	Positive	Viberg et al. (2013)
	PFBS uniquely suppressed differentiation of Ach and Dopa phenotypes in vitro PC12 (rat neuronal) cells. Concentration-dependent decrease in expression of tyrosine hydroxylase and choline acetyltransferase. Unlikely for one shared mechanism of toxicity across perfluorinated chemicals.	Positive	Slotkin et al. (2008)
	No finding of adverse association between PFOA exposure and neuropsychological test performance in children.	Negative	Stein et al. (2013)
Neuro toxicity	PFASs such as PFOA and PFOS may induce neurobehavioral effects, particularly in developmentally exposed animals.	Positive	Mariussen (2012)
	Neurobehavioral endpoints were not affected in Sprague Dawley rats following exposure to PFHxA (0, 20, 100 or 500 mg/kg b.w. per day NaPFHx) for 90 days.	Negative	Loveless et al. (2009)
	90-day evaluation of NaPFHxA in Sprague Dawley rats following oral gavage, indicates a NOAEL of 20 and 100 mg/kg-day in males and females, respectively, based on mild to minimal degeneration of the nasal epithelium.	Positive	Loveless et al. (2009)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
Respiratory Toxicity	Three cases of young healthy adults, following occupational inhalation of a new waterproofing formulation containing an acrylate fluoropolymer, developed a rapidly progressive dyspnea; two had hypoxemia and flu-like reactions within 1-2 hours of exposure.	Positive	Lazor-Blanchet et al. (2004)
	A commercial nanofilm spray product containing a perfluorosilane induced a concentration-dependent decrease of the tidal volume, significant increases of protein level in bronchoalveolar lavage fluid, reduced body weight, atelectasis, emphysema, and lung hemorrhages in BALB/cJ mice compared to a formulation containing alkylsilane.	Positive	Nørgaard et al. (2010)
	Normal pulmonary function in fluoropolymer plant workers with serum PFOA.	Negative	Sakr et al. (2007a); Sakr et al. (2007b)
Other Toxicological Hazard Traits	Increased PPAR-alpha activation in human-transfected COS-1 monkey kidney cells by PFHxA, PFBA, PFHxS, and PFBS. PFDA inactive.	Positive	Wolf et al. (2008)
	Chlorinated polyfluorinated ether sulfonates (6:2 Cl-PFAES and 8:2 Cl-PFAES) bound to PPAR- α , - β , and - γ with affinity higher than PFOS and has agonistic activity toward PPAR signaling pathways with similar or greater potency.	Positive	Li et al. (2018)
	28-day 150 mg/kg-d APFB repeat-dose oral study in S-D rats. Delay in pupillary reflex response. Outer retinal degeneration and loss of photoreceptors. Decrease in inner retinal, nuclear and synaptic terminal layer thicknesses. No mortality.	Positive	3M (2006); Butenhoff et al. (2012)
	Ammonium 4,8-dioxa-3H-perfluorononanoate (ADONA) toxicity was evaluated in acute and repeat-dose studies of up to 90-days duration. It is a mild skin irritant and a moderate to severe eye irritant in rabbits.	Positive	Gordon (2011)
	Survival rate of female Sprague-Dawley rats significantly decreased at presumed maximally tolerated dose of PFHxA (200 mg/kg-d) in 2-year study.	Positive	Klaunig et al. (2015)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	Experimental and QSAR modeling results showed PFAS mixture cytotoxicity in amphibian fibroblasts was approximately additive, except with PFOS and PFOA, which were weakly synergistic.	Positive	Hoover et al. (2019)
	No generation of reactive oxygen species (ROS) nor DNA damage by PFBS and PFHxA in HepG2 cells.	Negative	Eriksen et al. (2010)
Phytotoxicity	Aquatic toxicity of PFHxA, PFHpA, PFOA and PFNA to several species of marine microalgae increases with increasing chain length: for every extra perfluoromethylene group in the alkyl chain, toxicity increases twofold. EC50 ranges from 0.28 mM to 12.84 mM.	Positive	Latala et al. (2009)
	Toxicity of PFBS, PFOS, PFHxA, PFOA, PFDoA and PFTeA on the membrane system of the freshwater algae <i>Scenedesmus obliquus</i> increased with chain length. PFOS, PFDoA and PFTeA inhibits algal growth in a concentration-dependent manner.	Positive	Liu et al. (2008)
	A broad range of acute and chronic toxicity responses in algae and aquatic plants have been observed, with stronger effects associated with increased perfluoroalkyl chain length.	Positive	Beach et al. (2006); Ding and Peijnenburg (2013); Environment Canada (2006); Environment Canada (2012); Giesy et al. (2010); UNEP/POPS/POPRC (2016a); UNEP/POPS/POPRC (2016b); UNEP/POPS/POPRC (2016c); UNEP/POPS/POPRC (2016d); Ding et al. (2012a); Ding et al. (2012b); Hoke et al. (2012); Latala et al. (2009); Liu et al. (2008)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	The aquatic toxicity of seven PFASs was investigated on the root elongation of lettuce (<i>L. sativa</i>) seeds and photosynthesis of green algae (<i>P. subcapitata</i>). The toxic effects on lettuce seeds and green algae were found to be similar in a relative sense and were shown to have a good relationship with the fluorinated carbon-chain length. The toxicity of these chemicals increases with increasing fluorinated carbon chain length. PFBA was more toxic than expected, likely due to acidification of the test solution. Because the unicellular <i>P. subcapitata</i> is sensitive to the pH value of the test solution, the actual EC50 value of PFBA is lower and deviates from the relationship between log EC50 value and chain length.	Positive	Ding et al. (2012a)
Wildlife Developmental Impairment	Tail deformation and uninflated swim bladders in zebrafish exposed to PFBA. Lowered heart rates and cardiac output. Malformations of the head in zebrafish exposed to PFBS. No structure-activity relationship for some endpoints. Exposure concentrations 50-3000 mg/L.	Positive	Hagenaars et al. (2011)
	Adverse impacts on embryonic development, hatchability, and time to hatch of Japanese medaka (<i>O. latipes</i>) eggs from exposure to PFOA and PFOS	Positive	Ji et al. (2008)
	Developmental toxicity observed in zebrafish embryos exposed to 6:2 FTCA. Observed effects include: decreased hatching and survival percentages, reduced heart rate, increased malformations (especially pericardial edema), decreased erythrocyte number, and disrupted erythroid differentiation during zebrafish embryonic development.	Positive	Shi et al. (2017a)
	Delayed development (time to metamorphosis) in northern leopard frog (<i>Rana pipiens</i>) tadpoles exposed to 10 ug/L PFHxS for 40 days.	Positive	Hoover et al. (2017)

Health Hazard Category	Summary of Findings	Adverse Impact Finding	Reference
	<p>F-53B [6:2 chlorinated polyfluorinated ether sulfonate] exposure (1.5 - 12 mg/L) induced developmental toxicity, including delayed hatching, increased occurrence of malformations, and reduced survival. Malformations, including pericardial and yolk sac edemas, abnormal spines, bent tails, and uninflated swim bladders increased with time course and dose. Continuous exposure resulted in high accumulation levels in zebrafish embryos, suggesting an inability for embryos to eliminate this compound and a high cumulative risk to fish. F-53B induced cardiac toxicity and reduced heart rate.</p>	Positive	Shi et al. (2017b)
Wildlife Reproductive Impairment	<p>Decreased fecundity of female Japanese medaka (<i>O. latipes</i>) after exposure to PFOS and PFOA.</p>	Positive	Ji et al. (2008)
	<p>Egg concentrations of PFASs are negatively correlated with hatching success in wild and laboratory birds.</p>	Positive	Custer et al. (2012); Custer et al. (2014); Nordén et al. (2016)
	<p>PFOS has adverse impacts on the reproduction of bumblebees (<i>Bombus terrestris</i>), including reduced ovarian size.</p>	Positive	Mommaerts et al. (2011)

APPENDIX 3 – REPORT PREPARATION*

Preparers:

Simona A. Balan (SCP), Ph.D., Senior Environmental Scientist (Specialist)

Armeen Etemad (SCP), Hazardous Substances Engineer

Vivek Mathrani (HERO), Ph.D., Staff Toxicologist (Specialist)

Reviewers:

Andre Algazi, Project Leader

Karl Palmer, Acting Deputy Director for Safer Consumer Products Program

* This document is based on information presented in the Product-Chemical Profile for Carpets and Rugs Containing Perfluoroalkyl or Polyfluoroalkyl Substances.

APPENDIX 4 – APPENDIX REFERENCES

- 3M (2006) Final Report: 28-Day Oral Study in Rats. ToxDocs Study Number 08-047, MTDID 8391. TSCA 8(E) Supplemental Submission: Docket No. 8EHQ-06-16638, 3M Strategic Toxicology Laboratory, St. Paul, MN, August 19, 2006
- Apelberg BJ, Witter FR, Herbstman JB, et al (2007) Cord serum concentrations of perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) in relation to weight and size at birth. *Environ Health Perspect* 115:1670–1676. <https://doi.org/10.1289/ehp.10334>
- Barry V, Winquist A, Steenland K (2013) Perfluorooctanoic acid (PFOA) exposures and incident cancers among adults living near a chemical plant. *Environ Health Perspect* 121:1313–1318. <https://doi.org/10.1289/ehp.1306615>
- Beach SA, Newsted JL, Coady K, Giesy JP (2006) Ecotoxicological evaluation of perfluorooctanesulfonate (PFOS). *Rev Environ Contam Toxicol* 186:133–174
- Beekman M, Zweers P, Muller A, et al (2016) Evaluation of substances used in the GenX technology by Chemours. Dordrecht. RIVM Letter Report 2006-0174. Natl Inst Public Health Environ Minist Health Welf Sport Neth
- Biegel LB, Hurtt ME, Frame SR, et al (2001) Mechanisms of extrahepatic tumor induction by peroxisome proliferators in male CD rats. *Toxicol Sci* 60:44–55. <https://doi.org/10.1093/toxsci/60.1.44>
- Bonefeld-Jørgensen EC, Long M, Fredslund SO, et al (2014) Breast cancer risk after exposure to perfluorinated compounds in Danish women: a case–control study nested in the Danish National Birth Cohort. *Cancer Causes Control* 25:1439–1448. <https://doi.org/10.1007/s10552-014-0446-7>
- Butenhoff J (2002) Toxicity of ammonium perfluorooctanoate in male cynomolgus monkeys after oral dosing for 6 months. *Toxicol Sci* 69:244–257. <https://doi.org/10.1093/toxsci/69.1.244>
- Butenhoff JL, Bjork JA, Chang SC, et al (2012) Toxicological evaluation of ammonium perfluorobutyrate in rats: twenty-eight-day and ninety-day oral gavage studies. *Reprod Toxicol* 33:513–530. <https://doi.org/10.1016/j.reprotox.2011.08.004>
- Butenhoff JL, Chang SC, Ehresman DJ, York RG (2009) Evaluation of potential reproductive and developmental toxicity of potassium perfluorohexanesulfonate in Sprague Dawley rats. *Reprod Toxicol* 27:331–341. <https://doi.org/10.1016/j.reprotox.2009.01.004>

- C8 Science Panel (2012) Probable link evaluation of cancer. April 15, 2012. Available at: http://www.c8sciencepanel.org/pdfs/Probable_Link_C8_Cancer_16April2012_v2.pdf. Accessed 9 Oct 2017
- Cai Y, Chen H, Yuan R, et al (2019) Toxicity of perfluorinated compounds to soil microbial activity: Effect of carbon chain length, functional group and soil properties. *Sci Total Environ* 690:1162–1169. <https://doi.org/10.1016/j.scitotenv.2019.06.440>
- Chang ET, Adami HO, Boffetta P, et al (2014) A critical review of perfluorooctanoate and perfluorooctanesulfonate exposure and cancer risk in humans. *Crit Rev Toxicol* 44:1–81. <https://doi.org/10.3109/10408444.2014.905767>
- Chen Q, Huang R, Hua L, et al (2018) Prenatal exposure to perfluoroalkyl and polyfluoroalkyl substances and childhood atopic dermatitis: a prospective birth cohort study. *Environ Health* 17:. <https://doi.org/10.1186/s12940-018-0352-7>
- Chengelis CP, Kirkpatrick JB, Radovsky A, Shinohara M (2009) A 90-day repeated dose oral (gavage) toxicity study of perfluorohexanoic acid (PFHxA) in rats (with functional observational battery and motor activity determinations). *Reprod Toxicol* 27:342–351. <https://doi.org/10.1016/j.reprotox.2009.01.006>
- Corsini E, Sangiovanni E, Avogadro A, et al (2012) In vitro characterization of the immunotoxic potential of several perfluorinated compounds (PFCs). *Toxicol Appl Pharmacol* 258:248–255. <https://doi.org/10.1016/j.taap.2011.11.004>
- Costa G, Sartori S, Consonni D (2009) Thirty years of medical surveillance in perfluorooctanoic acid production workers. *J Occup Environ Med* 51:364–372. <https://doi.org/10.1097/JOM.0b013e3181965d80>
- Costantini D, Blévin P, Herzke D, et al (2019) Higher plasma oxidative damage and lower plasma antioxidant defences in an Arctic seabird exposed to longer perfluoroalkyl acids. *Environ Res* 168:278–285. <https://doi.org/10.1016/j.envres.2018.10.003>
- Custer CM, Custer TW, Dummer PM, et al (2014) Exposure and effects of perfluoroalkyl substances in tree swallows nesting in Minnesota and Wisconsin, USA. *Arch Environ Contam Toxicol* 66:120–138. <https://doi.org/10.1007/s00244-013-9934-0>
- Custer CM, Custer TW, Schoenfuss HL, et al (2012) Exposure and effects of perfluoroalkyl compounds on tree swallows nesting at Lake Johanna in east central Minnesota, USA. *Reprod Toxicol* 33:556–562. <https://doi.org/10.1016/j.reprotox.2011.01.005>

- Danish Environmental Protection Agency (2015) Short-chain polyfluoroalkyl substances (PFAS). A literature review of information on human health effects and environmental fate and effect aspects of short-chain PFAS. Project No. 1707, Copenhagen, Denmark
- Darrow LA, Stein CR, Steenland K (2013) Serum perfluorooctanoic acid and perfluorooctane sulfonate concentrations in relation to birth outcomes in the Mid-Ohio Valley, 2005-2010. *Environ Health Perspect* 121:1207–1213. <https://doi.org/10.1289/ehp.1206372>
- Das KP, Grey BE, Zehr RD, et al (2008) Effects of perfluorobutyrate exposure during pregnancy in the mouse. *Toxicol Sci* 105:173–181. <https://doi.org/10.1093/toxsci/kfn099>
- Ding G, Peijnenburg WJGM (2013) Physicochemical properties and aquatic toxicity of poly- and perfluorinated compounds. *Crit Rev Environ Sci Technol* 43:598–678. <https://doi.org/10.1080/10643389.2011.627016>
- Ding G, Wouterse M, Baerselman R, Peijnenburg WJGM (2012a) Toxicity of polyfluorinated and perfluorinated compounds to lettuce (*Lactuca sativa*) and green algae (*Pseudokirchneriella subcapitata*). *Arch Environ Contam Toxicol* 62:49–55. <https://doi.org/10.1007/s00244-011-9684-9>
- Ding GH, Frömel T, van den Brandhof EJ, et al (2012b) Acute toxicity of poly- and perfluorinated compounds to two cladocerans, *Daphnia magna* and *Chydorus sphaericus*. *Environ Toxicol Chem* 31:605–610. <https://doi.org/10.1002/etc.1713>
- Emmett EA, Zhang H, Shofer FS, et al (2006) Community exposure to perfluorooctanoate: relationships between serum levels and certain health parameters. *J Occup Environ Med* 48:771–779. <https://doi.org/10.1097/01.jom.0000233380.13087.37>
- Environment Canada (2006) Ecological screening assessment report on perfluorooctane sulfonate, its salts and its precursors that contain the C8F17SO2 or C8F17SO3, or C8F17SO2N moiety. June, 2006. Available at: https://www.ec.gc.ca/lcpe-cepa/documents/substances/spfo-pfos/ecological_sar_pfos_eng.pdf. Accessed 10 Oct 2017
- Environment Canada (2012) Ecological screening assessment report: Long-chain (C9–C20) perfluorocarboxylic acids, their salts and their precursors. August, 2012. Available at: https://www.ec.gc.ca/ese-ees/CA29B043-86E4-4308-BBC0-5CE163FEAC40/PFCAs_EN.pdf. Accessed 10 Oct 2017
- Eriksen KT, Raaschou-Nielsen O, Sørensen M, et al (2010) Genotoxic potential of the perfluorinated chemicals PFOA, PFOS, PFBS, PFNA and PFHxA in human HepG2 cells. *Mutat Res* 700:39–43. <https://doi.org/10.1016/j.mrgentox.2010.04.024>

- Eriksen KT, Sorensen M, McLaughlin JK, et al (2009) Perfluorooctanoate and perfluorooctanesulfonate plasma levels and risk of cancer in the general Danish population. *J Natl Cancer Inst* 101:605–609. <https://doi.org/10.1093/jnci/djp041>
- Fei C, McLaughlin JK, Lipworth L, Olsen J (2009) Maternal levels of perfluorinated chemicals and subfecundity. *Hum Reprod* 24:1200–1205. <https://doi.org/10.1093/humrep/den490>
- Fei C, McLaughlin JK, Tarone RE, Olsen J (2007) Perfluorinated chemicals and fetal growth: a study within the Danish National Birth Cohort. *Environ Health Perspect* 115:1677–1682. <https://doi.org/10.1289/ehp.10506>
- Fisher M, Arbuckle TE, Wade M, Haines DA (2013) Do perfluoroalkyl substances affect metabolic function and plasma lipids?—Analysis of the 2007–2009, Canadian Health Measures Survey (CHMS) Cycle 1. *Environ Res* 121:95–103. <https://doi.org/10.1016/j.envres.2012.11.006>
- Gallo V, Leonardi G, Genser B, et al (2012) Serum perfluorooctanoate (PFOA) and perfluorooctane sulfonate (PFOS) concentrations and liver function biomarkers in a population with elevated PFOA exposure. *Environ Health Perspect* 120:655–660. <https://doi.org/10.1289/ehp.1104436>
- Giesy JP, Naile JE, Khim JS, et al (2010) Aquatic toxicology of perfluorinated chemicals. *Rev Environ Contam Toxicol* 202:1–52. https://doi.org/10.1007/978-1-4419-1157-5_1
- Gilliland FD, Mandel JS (1993) Mortality among employees of a perfluorooctanoic acid production plant. *J Occup Environ Med* 35:950–954
- Gordon SC (2011) Toxicological evaluation of ammonium 4,8-dioxa-3H-perfluorononanoate, a new emulsifier to replace ammonium perfluorooctanoate in fluoropolymer manufacturing. *Regul Toxicol Pharmacol* 59:64–80. <https://doi.org/10.1016/j.yrtph.2010.09.008>
- Gorrochategui E, Pérez-Albaladejo E, Casas J, et al (2014) Perfluorinated chemicals: differential toxicity, inhibition of aromatase activity and alteration of cellular lipids in human placental cells. *Toxicol Appl Pharmacol* 277:124–130. <https://doi.org/10.1016/j.taap.2014.03.012>
- Grandjean P, Andersen EW, Budtz-Jørgensen E, et al (2012) Serum vaccine antibody concentrations in children exposed to perfluorinated compounds. *JAMA* 307:. <https://doi.org/10.1001/jama.2011.2034>
- Grandjean P, Heilmann C, Weihe P, et al (2016) Serum vaccine antibody concentrations in adolescents exposed to perfluorinated compounds. *Environ Health Perspect* 125:077018. <https://doi.org/10.1289/EHP275>

- Granum B, Haug LS, Namork E, et al (2013) Pre-natal exposure to perfluoroalkyl substances may be associated with altered vaccine antibody levels and immune-related health outcomes in early childhood. *J Immunotoxicol* 10:373–379. <https://doi.org/10.3109/1547691X.2012.755580>
- Grice MM, Alexander BH, Hoffbeck R, Kampa DM (2007) Self-reported medical conditions in perfluorooctanesulfonyl fluoride manufacturing workers. *J Occup Environ Med* 49:722–729. <https://doi.org/10.1097/JOM.0b013e3180582043>
- Hagenaars A, Vergauwen L, De Coen W, Knapen D (2011) Structure–activity relationship assessment of four perfluorinated chemicals using a prolonged zebrafish early life stage test. *Chemosphere* 82:764–772. <https://doi.org/10.1016/j.chemosphere.2010.10.076>
- Hardell E, Kärrman A, van Bavel B, et al (2014) Case-control study on perfluorinated alkyl acids (PFAAs) and the risk of prostate cancer. *Environ Int* 63:35–39. <https://doi.org/10.1016/j.envint.2013.10.005>
- Hoffman K, Webster TF, Weisskopf MG, et al (2010) Exposure to polyfluoroalkyl chemicals and attention deficit/hyperactivity disorder in U.S. children 12–15 years of age. *Environ Health Perspect* 118:1762–1767. <https://doi.org/10.1289/ehp.1001898>
- Hoke RA, Bouchelle LD, Ferrell BD, Buck RC (2012) Comparative acute freshwater hazard assessment and preliminary PNEC development for eight fluorinated acids. *Chemosphere* 87:725–733. <https://doi.org/10.1016/j.chemosphere.2011.12.066>
- Hoover G, Kar S, Guffey S, et al (2019) In vitro and in silico modeling of perfluoroalkyl substances mixture toxicity in an amphibian fibroblast cell line. *Chemosphere* 233:25–33. <https://doi.org/10.1016/j.chemosphere.2019.05.065>
- Hoover GM, Chislock MF, Tornabene BJ, et al (2017) Uptake and depuration of four per/polyfluoroalkyl substances (PFASS) in northern leopard frog *Rana pipiens* tadpoles. *Environ Sci Technol Lett* 4:399–403. <https://doi.org/10.1021/acs.estlett.7b00339>
- IARC (2016) International Agency for Research on Cancer (IARC): Perfluorooctanoic acid. IARC Monographs on the evaluation of carcinogenic risks to humans. Volume 110. Lyon, France
- Impinen A, Nygaard UC, Lødrup Carlsen KC, et al (2018) Prenatal exposure to perfluoroalkyl substances (PFASs) associated with respiratory tract infections but not allergy- and asthma-related health outcomes in childhood. *Environ Res* 160:518–523. <https://doi.org/10.1016/j.envres.2017.10.012>

- Ji K, Kim Y, Oh S, et al (2008) Toxicity of perfluorooctane sulfonic acid and perfluorooctanoic acid on freshwater macroinvertebrates (*daphnia magna* and *moina macrocopa*) and fish (*oryzias latipes*). *Environ Toxicol Chem* 27:2159. <https://doi.org/10.1897/07-523.1>
- Johnson PI, Sutton P, Atchley DS, et al (2014) The Navigation Guide—Evidence-based medicine meets environmental health: systematic review of human evidence for PFOA effects on fetal growth. *Environ Health Perspect* 122:1028–1039. <https://doi.org/10.1289/ehp.1307893>
- Klaunig JE, Shinohara M, Iwai H, et al (2015) Evaluation of the chronic toxicity and carcinogenicity of perfluorohexanoic acid (PFHxA) in Sprague-Dawley rats. *Toxicol Pathol* 43:209–220. <https://doi.org/10.1177/0192623314530532>
- Latala A, Nędzi M, Stepnowski P (2009) Acute toxicity assessment of perfluorinated carboxylic acids towards the Baltic microalgae. *Environ Toxicol Pharmacol* 28:167–171. <https://doi.org/10.1016/j.etap.2009.03.010>
- Lau C, Anitole K, Hodes C, et al (2007) Perfluoroalkyl acids: a review of monitoring and toxicological findings. *Toxicol Sci* 99:366–394. <https://doi.org/10.1093/toxsci/kfm128>
- Lazor-Blanchet C, Rusca S, Vernez D, et al (2004) Acute pulmonary toxicity following occupational exposure to a floor stain protector in the building industry in Switzerland. *Int Arch Occup Environ Health* 77:244–248. <https://doi.org/10.1007/s00420-004-0505-6>
- Leonard RC, Kreckmann KH, Sakr CJ, Symons JM (2008) Retrospective cohort mortality study of workers in a polymer production plant including a reference population of regional workers. *Ann Epidemiol* 18:15–22. <https://doi.org/10.1016/j.annepidem.2007.06.011>
- Li Y, Fletcher T, Mucs D, et al (2018) Half-lives of PFOS, PFHxS and PFOA after end of exposure to contaminated drinking water. *Occup Environ Med* 75:46–51. <https://doi.org/10.1136/oemed-2017-104651>
- Lieder PH, Chang SH, York RG, Butenhoff JL (2009) Toxicological evaluation of potassium perfluorobutanesulfonate in a 90-day oral gavage study with Sprague-Dawley rats. *Toxicology* 255:45–52. <https://doi.org/10.1016/j.tox.2008.10.002>
- Lin CY, Chen PC, Lin YC, Lin LY (2009) Association among serum perfluoroalkyl chemicals, glucose homeostasis, and metabolic syndrome in adolescents and adults. *Diabetes Care* 32:702–707. <https://doi.org/10.2337/dc08-1816>
- Liu W, Chen S, Quan X, Jin YH (2008) Toxic effect of serial perfluorosulfonic and perfluorocarboxylic acids on the membrane system of a freshwater alga measured by flow cytometry. *Environ Toxicol Chem* 27:1597. <https://doi.org/10.1897/07-459.1>

- Looker C, Luster MI, Calafat AM, et al (2014) Influenza vaccine response in adults exposed to perfluorooctanoate and perfluorooctanesulfonate. *Toxicol Sci* 138:76–88. <https://doi.org/10.1093/toxsci/kft269>
- Lopez-Espinosa MJ, Mondal D, Armstrong B, et al (2012) Thyroid function and perfluoroalkyl acids in children living near a chemical plant. *Environ Health Perspect* 120:1036–1041. <https://doi.org/10.1289/ehp.1104370>
- Loveless SE, Slezak B, Serex T, et al (2009) Toxicological evaluation of sodium perfluorohexanoate. *Toxicology* 264:32–44. <https://doi.org/10.1016/j.tox.2009.07.011>
- Lundin JI, Alexander BH, Olsen GW, Church TR (2009) Ammonium perfluorooctanoate production and occupational mortality. *Epidemiology* 20:921–928. <https://doi.org/10.1097/EDE.0b013e3181b5f395>
- MacNeil J, Steenland NK, Shankar A, Ducatman A (2009) A cross-sectional analysis of type II diabetes in a community with exposure to perfluorooctanoic acid (PFOA). *Environ Res* 109:997–1003. <https://doi.org/10.1016/j.envres.2009.08.002>
- Maisonet M, Terrell ML, McGeehin MA, et al (2012) Maternal concentrations of polyfluoroalkyl compounds during pregnancy and fetal and postnatal growth in British girls. *Environ Health Perspect* 120:1432–1437. <https://doi.org/10.1289/ehp.1003096>
- Mariussen E (2012) Neurotoxic effects of perfluoroalkylated compounds: mechanisms of action and environmental relevance. *Arch Toxicol* 86:1349–1367. <https://doi.org/10.1007/s00204-012-0822-6>
- Martin JW, Chan K, Mabury SA, O'Brien PJ (2009) Bioactivation of fluorotelomer alcohols in isolated rat hepatocytes. *Chem Biol Interact* 177:196–203. <https://doi.org/10.1016/j.cbi.2008.11.001>
- McCoy JA, Bangma JT, Reiner JL, et al (2017) Associations between perfluorinated alkyl acids in blood and ovarian follicular fluid and ovarian function in women undergoing assisted reproductive treatment. *Sci Total Environ* 605–606:9–17. <https://doi.org/10.1016/j.scitotenv.2017.06.137>
- Melzer D, Rice N, Depledge MH, et al (2010) Association between serum perfluorooctanoic acid (PFOA) and thyroid disease in the U.S. National Health and Nutrition Examination Survey. *Environ Health Perspect* 118:686–692. <https://doi.org/10.1289/ehp.0901584>
- Mommaerts V, Hagenars A, Meyer J, et al (2011) Impact of a perfluorinated organic compound PFOS on the terrestrial pollinator *Bombus terrestris* (Insecta, Hymenoptera). *Ecotoxicology* 20:447–456. <https://doi.org/10.1007/s10646-011-0596-2>

- Mundt DJ, Mundt KA, Luippold RS, et al (2007) Clinical epidemiological study of employees exposed to surfactant blend containing perfluorononanoic acid. *Occup Environ Med* 64:589–594. <https://doi.org/10.1136/oem.2007.032664>
- Nelson JW, Hatch EE, Webster TF (2010) Exposure to polyfluoroalkyl chemicals and cholesterol, body weight, and insulin resistance in the general U.S. population. *Environ Health Perspect* 118:197–202. <https://doi.org/10.1289/ehp.0901165>
- Nolan LA, Nolan JM, Shofer FS, et al (2009) The relationship between birth weight, gestational age and perfluorooctanoic acid (PFOA)-contaminated public drinking water. *Reprod Toxicol* 27:231–238. <https://doi.org/10.1016/j.reprotox.2008.11.001>
- Nordén M, Berger U, Engwall M (2016) Developmental toxicity of PFOS and PFOA in great cormorant (*Phalacrocorax carbo sinensis*), herring gull (*Larus argentatus*) and chicken (*Gallus gallus domesticus*). *Environ Sci Pollut Res* 23:10855–10862. <https://doi.org/10.1007/s11356-016-6285-1>
- Nørgaard AW, Larsen ST, Hammer M, et al (2010) Lung damage in mice after inhalation of nanofilm spray products: the role of perfluorination and free hydroxyl groups. *Toxicol Sci* 116:216–224. <https://doi.org/10.1093/toxsci/kfq094>
- OEHHA (2017) Office of Environmental Health Hazard Assessment (OEHHA): Chemicals Listed Effective November 10, 2017 as Known to the State of California to Cause Reproductive Toxicity: Perfluorooctanoic Acid (PFOA) and Perfluorooctane Sulfonate (PFOS). Available at: <https://oehha.ca.gov/proposition-65/crn/chemicals-listed-effective-november-10-2017-known-state-california-cause>. Accessed 9 Jan 2018
- Olsen GW, Burris JM, Burlew MM, Mandel JH (2000) Plasma cholecystokinin and hepatic enzymes, cholesterol and lipoproteins in ammonium perfluorooctanoate production workers. *Drug Chem Toxicol* 23:603–620. <https://doi.org/10.1081/DCT-100101973>
- Olsen GW, Burris JM, Burlew MM, Mandel JH (2003) Epidemiologic assessment of worker serum perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA) concentrations and medical surveillance examinations. *J Occup Environ Med* 45:260–270. <https://doi.org/10.1097/01.jom.0000052958.59271.10>
- Olsen GW, Gilliland FD, Burlew MM, et al (1998) An epidemiologic investigation of reproductive hormones in men with occupational exposure to perfluorooctanoic acid. *J Occup Environ Med* 40:614–622. <https://doi.org/10.1097/00043764-199807000-00006>

- Olsen GW, Zobel LR (2007) Assessment of lipid, hepatic, and thyroid parameters with serum perfluorooctanoate (PFOA) concentrations in fluorochemical production workers. *Int Arch Occup Environ Health* 81:231–246. <https://doi.org/10.1007/s00420-007-0213-0>
- Rae CJM, Craig L, Slone TW, et al (2015) Evaluation of chronic toxicity and carcinogenicity of ammonium 2,3,3,3-tetrafluoro-2-(heptafluoropropoxy)-propanoate in Sprague–Dawley rats. *Toxicol Rep* 2:939–949. <https://doi.org/10.1016/j.toxrep.2015.06.001>
- Ramhøj L, Hass U, Boberg J, et al (2018) Perfluorohexane sulfonate (PFHxS) and a mixture of endocrine disruptors reduce thyroxine levels and cause anti-androgenic effects in rats. *Toxicol Sci* 163:579–591. <https://doi.org/10.1093/toxsci/kfy055>
- Rosenmai AK, Ahrens L, le Godec T, et al (2018) Relationship between peroxisome proliferator-activated receptor alpha activity and cellular concentration of 14 perfluoroalkyl substances in HepG2 cells. *J Appl Toxicol* 38:219–226. <https://doi.org/10.1002/jat.3515>
- Rosenmai AK, Taxvig C, Svingen T, et al (2016) Fluorinated alkyl substances and technical mixtures used in food paper-packaging exhibit endocrine-related activity in vitro. *Andrology* 4:662–672. <https://doi.org/10.1111/andr.12190>
- Rushing BR, Hu Q, Franklin JN, et al (2017) Evaluation of the immunomodulatory effects of 2,3,3,3-tetrafluoro-2-(heptafluoropropoxy)-propanoate in C57BL/6 mice. *Toxicol Sci*. <https://doi.org/10.1093/toxsci/kfw251>
- Sakr CJ, Kreckmann KH, Green JW, et al (2007a) Cross-sectional study of lipids and liver enzymes related to a serum biomarker of exposure (ammonium perfluorooctanoate or APFO) as part of a general health survey in a cohort of occupationally exposed workers. *J Occup Environ Med* 49:1086–1096. <https://doi.org/10.1097/JOM.0b013e318156eca3>
- Sakr CJ, Leonard RC, Kreckmann KH, et al (2007b) Longitudinal study of serum lipids and liver enzymes in workers with occupational exposure to ammonium perfluorooctanoate. *J Occup Environ Med* 49:872–879. <https://doi.org/10.1097/JOM.0b013e318124a93f>
- Sakr CJ, Symons JM, Kreckmann KH, Leonard RC (2009) Ischaemic heart disease mortality study among workers with occupational exposure to ammonium perfluorooctanoate. *Occup Environ Med* 66:699–703. <https://doi.org/10.1136/oem.2008.041582>
- Savitz DA, Stein CR, Bartell SM, et al (2012a) Perfluorooctanoic acid exposure and pregnancy outcome in a highly exposed community. *Epidemiology* 23:386–392. <https://doi.org/10.1097/EDE.0b013e31824cb93b>

- Savitz DA, Stein CR, Elston B, et al (2012b) Relationship of perfluorooctanoic acid exposure to pregnancy outcome based on birth records in the mid-Ohio Valley. *Environ Health Perspect* 120:1201–1207. <https://doi.org/10.1289/ehp.1104752>
- Seacat AM (2002) Subchronic toxicity studies on perfluorooctanesulfonate potassium salt in cynomolgus monkeys. *Toxicol Sci* 68:249–264. <https://doi.org/10.1093/toxsci/68.1.249>
- Sheng N, Cui R, Wang J, et al (2018) Cytotoxicity of novel fluorinated alternatives to long-chain perfluoroalkyl substances to human liver cell line and their binding capacity to human liver fatty acid binding protein. *Arch Toxicol* 92:359–369. <https://doi.org/10.1007/s00204-017-2055-1>
- Sheng N, Zhou X, Zheng F, et al (2017) Comparative hepatotoxicity of 6:2 fluorotelomer carboxylic acid and 6:2 fluorotelomer sulfonic acid, two fluorinated alternatives to long-chain perfluoroalkyl acids, on adult male mice. *Arch Toxicol* 91:2909–2919. <https://doi.org/10.1007/s00204-016-1917-2>
- Shi G, Cui Q, Pan Y, et al (2017a) 6:2 fluorotelomer carboxylic acid (6:2 FTCA) exposure induces developmental toxicity and inhibits the formation of erythrocytes during zebrafish embryogenesis. *Aquat Toxicol* 190:53–61. <https://doi.org/10.1016/j.aquatox.2017.06.023>
- Shi G, Cui Q, Pan Y, et al (2017b) 6:2 Chlorinated polyfluorinated ether sulfonate, a PFOS alternative, induces embryotoxicity and disrupts cardiac development in zebrafish embryos. *Aquat Toxicol* 185:67–75. <https://doi.org/10.1016/j.aquatox.2017.02.002>
- Skuladottir M, Ramel A, Rytter D, et al (2015) Examining confounding by diet in the association between perfluoroalkyl acids and serum cholesterol in pregnancy. *Environ Res* 143:33–38. <https://doi.org/10.1016/j.envres.2015.09.001>
- Slotkin TA, MacKillop EA, Melnick RL, et al (2008) Developmental neurotoxicity of perfluorinated chemicals modeled in vitro. *Environ Health Perspect* 116:716–722. <https://doi.org/10.1289/ehp.11253>
- Song W, Liu QS, Sun Z, et al (2018) Polyfluorinated iodine alkanes regulated distinct breast cancer cell progression through binding with estrogen receptor alpha or beta isoforms. *Environ Pollut* 239:300–307. <https://doi.org/10.1016/j.envpol.2018.04.037>
- Steenland K, Fletcher T, Savitz DA (2010) Epidemiologic evidence on the health effects of perfluorooctanoic acid (PFOA). *Environ Health Perspect* 118:1100–1108. <https://doi.org/10.1289/ehp.0901827>
- Steenland K, Woskie S (2012) Cohort mortality study of workers exposed to perfluorooctanoic acid. *Am J Epidemiol* 176:909–917. <https://doi.org/10.1093/aje/kws171>

- Steenland K, Zhao L, Winquist A (2015) A cohort incidence study of workers exposed to perfluorooctanoic acid (PFOA). *Occup Environ Med* 72:373–380. <https://doi.org/10.1136/oemed-2014-102364>
- Stein CR, Savitz DA, Bellinger DC (2013) Perfluorooctanoate and neuropsychological outcomes in children. *Epidemiology* 24:590–599. <https://doi.org/10.1097/EDE.0b013e3182944432>
- Stein CR, Savitz DA, Dougan M (2009) Serum levels of perfluorooctanoic acid and perfluorooctane sulfonate and pregnancy outcome. *Am J Epidemiol* 170:837–846. <https://doi.org/10.1093/aje/kwp212>
- UNEP/POPS/POPRC (2016a) United Nations Environment Programme, Stockholm Convention on Persistent Organic Pollutants, Persistent Organic Pollutants Review Committee (UNEP/POPS/POPRC): Draft risk profile: pentadecafluorooctanoic acid (CAS No: 335-67-1, PFOA, perfluorooctanoic acid), its salts and PFOA-related compounds (UNEP/POPS/POPRC.12/3, 14 June 2016). Twelfth meeting, 19–23 September 2016, Rome, Italy
- UNEP/POPS/POPRC (2016b) United Nations Environment Programme, Stockholm Convention on Persistent Organic Pollutants, Persistent Organic Pollutants Review Committee (UNEP/POPS/POPRC): Additional information document (UNEP/POPS/POPRC.12/INF/5, 18 July 2016). Twelfth meeting, 19–23 September 2016, Rome, Italy
- UNEP/POPS/POPRC (2016c) United Nations Environment Programme, Stockholm Convention on Persistent Organic Pollutants, Persistent Organic Pollutants Review Committee (UNEP/POPS/POPRC): Comments and responses relating to the draft profile (UNEP/POPS/POPRC.12/INF/6*, 20 July 2016). Twelfth meeting, 19–23 September 2016, Rome, Italy
- UNEP/POPS/POPRC (2016d) United Nations Environment Programme, Stockholm Convention on Persistent Organic Pollutants, Persistent Organic Pollutants Review Committee (UNEP/POPS/POPRC): Additional comments and responses relating to the third draft (UNEP/POPS/POPRC.12/INF/6/Add1, 14 September 2016). Twelfth meeting, 19–23 September 2016, Rome, Italy
- U.S. EPA (2016a) U.S. Environmental Protection Agency (U.S. EPA): Drinking water health advisories for PFOA and PFOS. Available at: <https://www.epa.gov/ground-water-and-drinking-water/drinking-water-health-advisories-pfoa-and-pfos>. Accessed 14 Oct 2017
- U.S. EPA (2016b) U.S. Environmental Protection Agency (U.S. EPA): Health effects support document for perfluorooctanoic acid (PFOA). Office of Water (4304T), Health and Ecological Criteria Division, Washington, DC, EPA Document Number 822-R-16-003, May 2016. Available at:

https://www.epa.gov/sites/production/files/2016-05/documents/pfoa_hesd_final_508.pdf.
Accessed 28 Nov 2017

- Velez MP, Arbuckle TE, Fraser WD (2015) Maternal exposure to perfluorinated chemicals and reduced fecundity: the MIREC study. *Hum Reprod* 30:701–709. <https://doi.org/10.1093/humrep/deu350>
- Viberg H, Lee I, Eriksson P (2013) Adult dose-dependent behavioral and cognitive disturbances after a single neonatal PFHxS dose. *Toxicology* 304:185–191.
<https://doi.org/10.1016/j.tox.2012.12.013>
- Vieira VM, Hoffman K, Shin HM, et al (2013) Perfluorooctanoic acid exposure and cancer outcomes in a contaminated community: a geographic analysis. *Environ Health Perspect* 121:318–323.
<https://doi.org/10.1289/ehp.1205829>
- Wang J, Zhang Y, Zhang W, et al (2012) Association of perfluorooctanoic acid with HDL cholesterol and circulating miR-26b and miR-199-3p in workers of a fluorochemical plant and nearby residents. *Environ Sci Technol* 46:9274–9281. <https://doi.org/10.1021/es300906q>
- Weiss JM, Andersson PL, Lamoree MH, et al (2009) Competitive binding of poly- and perfluorinated compounds to the thyroid hormone transport protein transthyretin. *Toxicol Sci* 109:206–216.
<https://doi.org/10.1093/toxsci/kfp055>
- Winquist A, Steenland K (2014a) Modeled PFOA exposure and coronary artery disease, hypertension, and high cholesterol in community and worker cohorts. *Environ Health Perspect* 122:1299–1305. <https://doi.org/10.1289/ehp.1307943>
- Winquist A, Steenland K (2014b) Perfluorooctanoic acid exposure and thyroid disease in community and worker cohorts. *Epidemiology* 25:255–264.
<https://doi.org/10.1097/EDE.0000000000000040>
- Wolf CJ, Takacs ML, Schmid JE, et al (2008) Activation of mouse and human peroxisome proliferator-activated receptor alpha by perfluoroalkyl acids of different functional groups and chain lengths. *Toxicol Sci* 106:162–171. <https://doi.org/10.1093/toxsci/kfn166>
- Zhang C, Sundaram R, Maisog J, et al (2015) A prospective study of prepregnancy serum concentrations of perfluorochemicals and the risk of gestational diabetes. *Fertil Steril* 103:184–189.
<https://doi.org/10.1016/j.fertnstert.2014.10.001>
- Zhao B, Lian Q, Chu Y, et al (2010) The inhibition of human and rat 11 β -hydroxysteroid dehydrogenase 2 by perfluoroalkylated substances. *J Steroid Biochem Mol Biol* 125:143–147.
<https://doi.org/10.1016/j.jsbmb.2010.12.017>

