



Work Plan Implementation:

Perfluoroalkyl and Polyfluoroalkyl Substances (PFASs) in Carpets, Rugs, Indoor Upholstered Furniture, and their Care and Treatment Products

NOVEMBER 15, 2016



Introduction

The Department of Toxic Substances Control's (DTSC) Safer Consumer Products (SCP) Regulations specify the process and criteria by which DTSC evaluates consumer products for possible designation as Priority Products. Every three years, DTSC issues a Priority Product Work Plan, which "... identifies and describes the product categories that [DTSC] will evaluate ..." over the next three years. DTSC's current Work Plan includes seven product categories and lists the five policy priorities that informed their selection.

Since issuing the Work Plan, DTSC has conducted a review of the product categories and has identified specific subcategories, chemicals, and chemical classes that align with our policy priorities. This document is one of three that summarizes the findings of the research on a specific topic. It describes our preliminary findings and concerns around the topic and identifies data gaps we seek to fill and specific questions we hope our stakeholders can answer. This document signals the beginning of an ongoing dialogue with interested stakeholders from all sectors (manufacturers, NGOs, governments, and academia). We believe the dialogue will ultimately inform our selection of future Priority Products.

Background

The class of perfluoroalkyl and polyfluoroalkyl substances* (PFASs) contains more than 3,000 man-made chemicals with at least one fully fluorinated carbon atom.¹⁻³ All PFASs are [Candidate Chemicals](#) for the DTSC's [Safer Consumer Products program \(SCP\)](#), because the entire class was added by Biomonitoring California to its list of Priority Chemicals⁴ for measuring in the bodily fluids of Californians. PFASs have a variety of applications, such as water-, oil-, soil- and stain-repellents. They are found in all of the SCP [2015-2017 Work Plan](#) product categories: beauty, personal care and hygiene products, building products, household and office furniture and furnishings, cleaning products, clothing, fishing and angling equipment; and office consumable products. Out of these six product categories, DTSC has identified carpets, rugs, indoor upholstered furniture, and their associated care and treatment products, as the largest potential sources of significant and widespread PFAS exposures.

DTSC's literature review of PFASs found evidence for (1) hazard characteristics (high persistence, bioaccumulation, potential toxicity) and (2) potential for widespread exposures for humans and other living organisms. DTSC is requesting additional information from stakeholders about the environmental and health hazards of PFASs, as well as the use of PFASs in carpets, rugs, indoor upholstered furniture, and their care and treatment products.

Hazard Traits of PFASs

The best-defined hazard trait of this chemical class is its high environmental persistence.⁵ PFASs that do break down in the environment and in living organisms convert to other PFASs that persist indefinitely and may be even more harmful.^{6,7} Unlike other persistent organic pollutants, PFASs accumulate in protein-rich organs rather than in fat.⁸

Most of the publicly-available research to date is limited to a few longer-chain[†] PFASs, such as perfluorooctanoate (PFOA) and perfluorooctane sulfonate (PFOS). Longer-chain PFASs are known to accumulate and persist in living organisms, including humans.⁹ They have been linked to acute and chronic toxicity in aquatic life, such as microorganisms, algae, plants, invertebrates, amphibians, fish, and birds.^{10,11} PFAS toxicity is less well characterized in terrestrial species, despite reports of their presence in a wide range of organisms.¹²⁻¹⁴ In humans, potential adverse health effects from chronic longer-chain PFAS exposure include

* "Perfluoro" means that all carbons in the molecule are fully fluorinated, while "polyfluoro" refers to molecules containing a mix of fully-, partially-, and non-fluorinated carbons.

† Longer-chain PFASs include perfluoroalkyl carboxylates and sulfonates with six or more fluorinated carbons, and their precursors.

increased serum cholesterol,^{15,16} thyroid disease,¹⁷⁻¹⁹ liver²⁰ and kidney²¹ damage, immune system disruption,²²⁻²⁴ pregnancy-induced hypertension,²⁵ and kidney and testicular cancers.²⁶

Under a voluntary agreement with the U.S. EPA, major PFASs manufacturers phased out the production and emission of longer-chain PFASs and precursors in 2015.²⁷ As a result, biomonitoring studies have found decreasing levels of longer-chain PFASs in human tissues.²⁸ Most replacements are also PFASs, including fluorinated ethers and short-chain PFASs.²⁹ Fluorinated ethers were thought to degrade in the environment, but recent studies have cast doubt on that assumption.^{30,31} Shorter-chain PFASs are marketed as less toxic compared to the long-chains, mainly because they bioaccumulate less and are more readily eliminated from living organisms. However, they are equally persistent and more mobile in the environment than the chemicals they are replacing, and also show potential for toxicity.³²

Potential Exposures to PFASs




Figure 1: Potential routes of exposure to PFASs from

PFASs are found globally in the aquatic^{33,34} and terrestrial environment,³⁵ indoor dust,³⁶ drinking water sources,³⁷ food,³⁸ wildlife,³⁹ and humans,^{40,41} including the milk and serum of breastfeeding women.^{42,43}

Exposure begins early in a person's life, since mothers transfer PFASs to their babies via the placenta and breastfeeding.⁴⁴⁻⁴⁸ Young children may experience higher exposure levels than adults due to greater dust ingestion rates⁴⁹ and hand-to-mouth transfer from PFASs-treated consumer products.⁵⁰ Industrial workers, carpet installers, carpet cleaners and workers in

furniture, furnishings, outdoor clothing, and carpet stores may also experience above average PFASs exposure levels.^{51,52} For the general population, PFASs exposure occurs mainly via contaminated food and drinking water,⁵³⁻⁵⁵ yet questions remain regarding how PFASs enter the environment and food and water supplies.

A large percentage of the PFASs produced worldwide are used to treat carpets and other home textiles to confer stain-, soil-, oil- or water-resistance.⁵⁶⁻⁵⁸ During normal use, treated carpets, rugs, indoor upholstered furniture and other consumer products release PFASs into indoor air and dust.^{59,60} An estimated 50% of the PFAS treatment may be lost from carpets due to walking and vacuuming over a typical nine-year product lifespan.⁶¹ People ingest or inhale the contaminated house dust,^{62,63} and inhale volatile PFASs.⁶⁴ They can also be exposed to large amounts of PFASs when using after-market carpet and upholstery cleaning and treatment products.^{65,66}




Figure 2: Exposure pathways to PFASs used in carpets, rugs, indoor upholstered furniture, and their care and treatment products.

DTSC seeks further information to help elucidate to what extent the manufacturing, use, and disposal of PFASs-treated carpets, rugs and indoor upholstered furniture contribute to the PFASs contamination of human food, drinking water, or aquatic and terrestrial ecosystems. Available information suggests that current disposal practices for carpet manufacturing and installation waste, unused and used carpeting, and other PFASs-treated consumer products that end up in landfills may release PFASs into the environment. Studies of landfills have

detected PFASs in leachates⁶⁷⁻⁷¹ and gaseous emissions,⁷²⁻⁷⁴ Additionally, wastewater treatment plants (WWTPs) collecting landfill leachates, surface runoff, and residential and commercial wastewater, cannot effectively remove PFASs.^{75,76} As a result, when wastewater effluent is discharged into surface waters, PFASs are released into the environment, contaminating aquatic ecosystems and drinking water sources.⁷⁷⁻⁷⁹ Sewage sludge also contains PFASs, thus the application of biosolids on soil can lead to the contamination of terrestrial ecosystems, drinking water, and food supplies.⁸⁰⁻⁸⁴

Next Steps

SCP is asking stakeholders to address the questions posted on our online information management system, [CalSAFER](#), by December 30, 2016. This will help the Department determine whether carpets, rugs, indoor upholstered furniture and their care and treatment products contribute to human and ecological exposures to PFASs and whether such exposures have the potential for significant and widespread adverse impacts. We will also host a [public workshop](#) on Tuesday, January 31, 2017, in the CalEPA Headquarters Building, Sacramento. Please monitor our [Priority Products Work Plan webpage](#) for updates on this topic.

References

- ¹ Buck RC, Franklin J, Berger U, Conder JM, Cousins IT, de Voogt P, Jensen AA, Kannan K, Mabury SA, van Leeuwen SPJ. Perfluoroalkyl and polyfluoroalkyl substances in the environment: Terminology, classification, and origins. *Integrated Environmental Assessment and Management* 2011;7(4):513-541.
- ² KEMI (Swedish Chemicals Agency) 2015. Occurrence and use of highly fluorinated substances and alternatives. Report from a government assessment. Report 7/15.
- ³ Biomonitoring California 2015. Potential Priority Chemicals: Perfluoroalkyl and Polyfluoroalkyl Substances (PFASs)
http://biomonitoring.ca.gov/sites/default/files/downloads/PotentialPriority_PFASs_111815.pdf
- ⁴ Biomonitoring California 2016. Perfluoroalkyl and Polyfluoroalkyl Substances (PFASs).
<http://biomonitoring.ca.gov/chemicals/perfluoroalkyl-and-polyfluoroalkyl-substances-pfass>.
- ⁵ Organisation for Economic Cooperation and Development (OECD). Synthesis paper on per- and polyfluorinated chemicals (PFCs). Environment Directorate, Paris 2013.
- ⁶ Krafft MP and Riess JG. Selected physicochemical aspects of poly- and perfluoroalkylated substances relevant to performance, environment and sustainability-Part One. *Chemosphere* 2015;129:4-19.
- ⁷ Lee H, D'eon J, Mabury SA. Biodegradation of Polyfluoroalkyl Phosphates as a Source of Perfluorinated Acids to the Environment. *Environmental Science and Technology* 2010;44:3305-10.

- ⁸ Pérez F, Nadal M, Navarro-Ortega A, Fàbrega F, Domingo JL, Barceló D, Farré M. Accumulation of perfluoroalkyl substances in human tissues. *Environment International* 2013;59:354-62.
- ⁹ Organisation for Economic Cooperation and Development (OECD). Synthesis paper on per- and polyfluorinated chemicals (PFCs). Environment Directorate, Paris 2013.
- ¹⁰ Giesy JP, Naile JE, Khim JS, Jones KC, Newsted JL. Aquatic toxicology of perfluorinated chemicals. *Reviews of Environmental Contamination and Toxicology* 2010;202:1-52.
- ¹¹ Ding G and Peijnenburg WJGM. Physicochemical properties and aquatic toxicity of poly- and perfluorinated compounds. *Critical Reviews in Environmental Science and Technology* 2013;43:598–678.
- ¹² Giesy JP and Kannan K. Global distribution of perfluorooctane sulfonate in wildlife. *Environmental Science and Toxicology* 2001;35:1339–1342.
- ¹³ Smithwick M, Muir DCG, Mabury SA, Solomon KR, Martin JW, Sonne C, Born EW, Letcher RJ, Dietz R. Perfluoroalkyl contaminants in liver tissue from East Greenland polar bears (*Ursus maritimus*). *Environmental Toxicology and Chemistry* 2005;24(4):981-6.
- ¹⁴ Falandysz J, Taniyasu S, Yamashita N, Rostkowski P, Zalewski K, Kannan K. Perfluorinated compounds in some terrestrial and aquatic wildlife species from Poland. *Journal of Environmental Science and Health Part A*. 2007;42:715-9.
- ¹⁵ Wang J, Zhang Y, Zhang W, Jin Y, Dai J. Association of perfluorooctanoic acid with HDL cholesterol and circulating miR-26b and miR-199-3p in workers of a fluorochemical plant and nearby residents. *Environmental Science & Technology* 2012;46(17):9274-81.
- ¹⁶ Fisher M, Arbuckle TE, Wade M, Haines DA. Do perfluoroalkyl substances affect metabolic function and plasma lipids? Analysis of the 2007-2009, Canadian Health Measures Survey (CHMS) Cycle 1. *Environmental Research* 2013;121:95-103.
- ¹⁷ Melzer D, Rice N, Depledge MH, Henley We, Galloway TS. Association between serum perfluorooctanoic acid (PFOA) and thyroid disease in the U.S. National Health and Nutrition Examination Survey. *Environmental Health Perspectives* 2010;118(5):686-692.
- ¹⁸ Lopez-Espinosa MJ, Debapriya M, Armstrong B, Bloom MS, Fletcher T. Thyroid function and perfluoroalkyl acids in children living near a chemical plant. *Environmental health perspectives* 2012;120(7):1036-41.
- ¹⁹ Winquist A and Steenland K. Perfluorooctanoic acid exposure and thyroid disease in community and worker cohorts. *Epidemiology* 2014;25(2):255-64.
- ²⁰ Gallo V, Leonardi G, Genser B, Lopez-Espinosa MJ, Frisbee SJ, Karlsson L, Ducatman AM, Fletcher T. Serum perfluorooctanoate (PFOA) and perfluorooctane sulfonate (PFOS) concentrations and liver function biomarkers in a population with elevated PFOA exposure. *Environmental health perspectives* 2012;120(5): 655-60.
- ²¹ Steenland K and Woskie S. Cohort mortality study of workers exposed to perfluorooctanoic acid. *American Journal of Epidemiology* 2012;176(10):909-17.
- ²² Grandjean P and Budtz-Jørgensen E. Immunotoxicity of perfluorinated alkylates: calculation of benchmark doses based on serum concentrations in children. *Environmental Health* 2013;12:35-42.
- ²³ Granum B, Haug LS, Namork E, Stølevik SB, Thomsen C, Aaberge IS, van Loveren H, Løvik M, Nygaard UC. Pre-natal exposure to perfluoroalkyl substances may be associated with altered

vaccine antibody levels and immune-related health outcomes in early childhood. *Journal of Immunotoxicology* 2013;10(4):373-9.

²⁴ Looker C, Luster MI, Calafat AM, Johnson VJ, Burleson GR, Burleson FG, Fletcher T. Influenza vaccine response in adults exposed to perfluorooctanoate and perfluorooctanesulfonate. *Toxicological Sciences* 2014;138(1):76-88.

²⁵ C8 Science Panel. Probable Link Evaluation of Pregnancy Induced Hypertension and Preeclampsia. December 5, 2011. Available at http://www.c8sciencepanel.org/pdfs/Probable_Link_C8_PIH_5Dec2011.pdf.

²⁶ C8 Science Panel. Probable Link Evaluation of Cancer. April 15, 2012. Available at http://www.c8sciencepanel.org/pdfs/Probable_Link_C8_Cancer_16April2012_v2.pdf.

²⁷ U.S. Environmental Protection Agency (EPA). 2010/2015 PFOA Stewardship Program. Available at <https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/and-polyfluoroalkyl-substances-pfass-under-tsca#tab-3>.

²⁸ Wu X, Bennett DH, Calafat AM, Kato K, Strynar M, Andersen E, Moran RE, Tancredi DJ, Tulse NS, Hertz-Picciotto IH. Serum concentrations of perfluorinated compounds (PFC) among selected populations of children and adults in California. *Environmental Research* 2015;136:264-73.

²⁹ Wang Z, Cousins IT, Scheringer M, Hugerbühler K. Fluorinated alternatives to long-chain perfluoroalkyl carboxylic acids (PFCAs), perfluoroalkane sulfonic acids (PFASs) and their potential precursors. *Environment International* 2013;60:242-8.

³⁰ Strynar M, Dagnino S, McMahan R, Liang S, Lindstrom A, Andersen E, McMillan L, Thurman M, Ferrer I, Ball C. Identification of novel perfluoroalkyl ether carboxylic acids (PFECAs) and sulfonic acids (PFESAs) in natural waters using accurate mass time-of-flight mass spectrometry (TOFMS). *Environmental Science & Technology* 2015;49(19):11622-30.

³¹ Ruan T, Lin Y, Wang T, Liu R, Jiang G. Identification of novel polyfluorinated ether sulfonates as PFOS alternatives in municipal sewage sludge in China. *Environmental Science & Technology* 2015;49:6519-27.

³² Wang Z, Cousins IT, Scheringer M, Hugerbuehler K. Hazard assessment of fluorinated alternatives to long-chain perfluoroalkyl acids (PFAAs) and their precursors: Status quo, ongoing challenges and possible solutions. *Environment International* 2015;75:172-9.

³³ Ahrens L and M Bundschuh. Fate and effects of poly- and perfluoroalkyl substances in the aquatic environment: A review. *Environmental Toxicology and Chemistry* 2014;33:1921-9.

³⁴ Benskin JP, Muir DCG, Scott BF, Spencer C, De Silva AO, Kylin H, Martin JW, Morris A, Lohmann R, Tomy G, Rosenberg B, Taniyasu S, Yamashita N. Perfluoroalkyl acids in the Atlantic and Canadian Arctic Oceans, *Environmental Science & Technology* 2012;46:5815-23.

³⁵ Rankin K, Mabury SA, Jenkins TM, Washington JW. A North American and global survey of perfluoroalkyl substances in surface soils: Distribution patterns and mode of occurrence. *Chemosphere* 2016;161:333-41.

³⁶ Eriksson U and Kärman. World-wide indoor exposure to polyfluoroalkyl phosphate esters (PAPs) and other PFASs in household dust. *Environmental Science & Technology* 2015;49(24):14503-11.

- ³⁷ Post GB, Cohn PD, Cooper KR. Perfluorooctanoic acid (PFOA), an emerging drinking water contaminant: A critical review of recent literature. *Environmental Research* 2012;116:93-117.
- ³⁸ Pérez F, Llorca M, Köck-Schulmeyer M, Škrbić B, Oliveira LS, da Boit Martinello K, Al-Dhabi NA, Antić I, Farré M, Barceló D. Assessment of perfluoroalkyl substances in food items at global scale. *Environmental Research* 2014;135:181-9.
- ³⁹ Houde M, De Silva AO, Muir DCG, Lechter RJ. Monitoring of perfluorinated compounds in aquatic biota: An updated review. *Environmental Science & Technology* 2011;45:7962-7973.
- ⁴⁰ Calafat AM, Kuklenyik Z, Reidy JA, Caudill SP, Tully JS, Needham LL. Serum concentrations of 11 polyfluoroalkyl compounds in the U.S. population: Data from the National Health and Nutrition Examination Survey (NHANES) 1999-2000. *Environmental Science & Technology* 2007;41(7):2237-2242.
- ⁴¹ Krafft MP and Riess JG. Per- and polyfluorinated substances (PFASs): Environmental challenges. *Current Opinion in Colloid & Interface Science* 2015;20:192-212.
- ⁴² Fujii Y, Yan J, Harada KH, Hitomi T, Yang H, Wang P, Koizumi A. Levels and profiles of long-chain perfluorinated carboxylic acids in human breast milk and infant formulas in East Asia. *Chemosphere* 2012;86:315-21.
- ⁴³ Cariou R, Veyrand B, Yamada A, Berrebi A, Zalkod D, Durand S, Pollono C, Marchand P, Leblanc JC, Antignac JP, Le Bizec B. Perfluoroalkyl acid (PFAA) levels and profiles in breast milk, maternal and cord serum of French women and their newborns. *Environment International* 2015;84:71-81.
- ⁴⁴ Yang L, Wang Z, Shi Y, Li J, Wang Y, Zhao Y, Wu Y, Cai Z. Human placental transfer of perfluoroalkyl acid precursors: Levels and profiles in paired maternal and cord serum. *Chemosphere* 2016;144:1631-8.
- ⁴⁵ Kim S, Choi K, Ji K, Seo J, Kho Y, Park J, Kim S, Park S, Hwang I, Jeon J, Yang H, Gisey JP. Trans-placental transfer of thirteen perfluorinated compounds and relations with fetal thyroid hormones. *Environmental Science & Technology* 2011;45:7465-7472.
- ⁴⁶ Lee YJ, Kim M, Bae J, Yang J. Concentrations of perfluoroalkyl compounds in maternal and umbilical cord sera and birth outcomes in Korea. *Chemosphere* 2013;90:1603-9.
- ⁴⁷ Fromme H, Mosch C, Morovitz M, Alba-Alejandre I, Boehmer S, Kiranoglu M, Fabre F, Hannibal I, Genzel-Boroviczeny O, Koletzko B, Volkel W. 2010. Pre- and Postnatal Exposure to Perfluorinated Compounds (PFCs) *Environmental Science & Technology* 2010;44:7123-7129.
- ⁴⁸ Mogensen UB, Grandjean P, Nielsen F, Weihe P, Budtz-Jørgensen. Breastfeeding as an exposure pathway for perfluorinated alkylates. *Environmental Science & Technology* 2015;49(17):10466-10473.
- ⁴⁹ Goosey E and Harrad S. Perfluoroalkyl compounds in dust from Asian, Australian, European, and North American homes and UK cars, classrooms, and offices. *Environment International* 2011;37:86-92.
- ⁵⁰ Trudel D, Horowitz L, Wormuth M, Scheringer M, Cousins IT, Hungerbühle K. Estimating Consumer Exposure to PFOS and PFOA. *Risk Analysis* 2008;28(2):251-69.
- ⁵¹ Agency for Toxic Substances & Disease Registry (ATSDR). Public Health Statement for Perfluoroalkyls. 2015. Available at <http://www.atsdr.cdc.gov/phs/phs.asp?id=1115&tid=237>.

- ⁵² Langer V, Dreyer A, Ebinghaus R. Polyfluorinated compounds in residential and nonresidential indoor air. *Environmental Science & Technology* 2010;44:8075-81.
- ⁵³ Egeghy PP and Lorber M. An assessment of the exposure of Americans to perfluorooctane sulfonate: A comparison of estimated intake with values inferred from NHANES data. *Journal of Exposure Science and Environmental Epidemiology* 2011;21:150–168.
- ⁵⁴ Noorlander CW, te Biesebeek JD, van Leeuwen SPJ, Zeilmaker MJ. Dietary intake of PFOS and PFOA in The Netherlands. RIVM/SIR Project V/320126 Perfluorinated compounds in food. Report number 320126001. May 2010.
- ⁵⁵ Domingo JL. Health risks of dietary exposure to perfluorinated compounds. *Environment International* 2012;40:187-95.
- ⁵⁶ Schultz MM, Barofsky DF, Field JA. Fluorinated alkyl surfactants. *Environmental Engineering Science* 2003;20:487–501.
- ⁵⁷ Guo Z, Liu X, Krebs KA, Roache NF. Perfluorocarboxylic acid content in 116 articles of commerce. U.S. Environmental Protection Agency, Washington, DC, EPA/600/R-09/033, 2009.
- ⁵⁸ Liu X, Guo Z, Krebs KA, Pope RH, Roache NF. Trends of perfluoroalkyl acid content in articles of commerce – Market monitoring from 2007 through 2011. U.S. Environmental Protection Agency, Washington, DC, EPA/600/R-09/033, 2009. EPA/600/R-12/585, 2012.
- ⁵⁹ Washburn ST, Bingman TS, Braithwaite SK, Buck RB, Buxton LW, Clewell HJ, Haroun LA, Kester JE, Rickard RW, Shipp AM. Exposure assessment and risk characterization for perfluorooctanoate in selected consumer articles. *Environmental Science and Technology* 2005;39: 3904-10.
- ⁶⁰ Gewurtz SB, Bhavsar SP, Crozier PW, Diamond ML, Helm PA, Marvin CH, Reiner EJ. Perfluoroalkyl contaminants in window film: Indoor/outdoor, urban/rural, and winter/summer contamination and assessment of carpet as a possible source. *Environmental Science & Technology* 2009;43:7317-23.
- ⁶¹ Battelle Memorial Institute. Part 1: Life-Cycle Waste Stream Estimates. Final Report for 3M Specialty Materials. April 21, 2000.
- ⁶² Tian Z, Kim SK, Shoeib M, Oh JE, Park JE. Human exposure to per- and polyfluoroalkyl substances (PFASs) via house dust in Korea: Implication to exposure pathway. *Science of the Total Environment* 2016;553:266-75.
- ⁶³ Strynar MJ and Lindstrom AB. Perfluorinated Compounds in House Dust from Ohio and North Carolina, USA. *Environmental Science and Technology* 2008;42:3751-56.
- ⁶⁴ Shoeib M, Harner T, Webster GM, Lee SC. Indoor Sources of Poly- and Perfluorinated Compounds (PFCS) in Vancouver, Canada: Implications for Human Exposure. *Environmental Science and Technology* 2011;45:7999-8005.
- ⁶⁵ Beeson S, Genuis SJ, Benskin JP, Martin JW. Exceptionally high serum concentrations of perfluorohexanesulfonate in a Canadian family are linked to home carpet treatment applications. *Environmental Science & Technology* 2012;46:12960-7.
- ⁶⁶ Guo Z, Liu X, Krebs KA, Roache NF. Perfluorocarboxylic acid content in 116 articles of commerce. U.S. Environmental Protection Agency, Washington, DC, EPA/600/R-09/033, 2009.
- ⁶⁷ Busch J, Ahrens L, Sturm R, Ebinghaus R. Polyfluoroalkyl compounds in landfill leachates. *Environmental Pollution* 2010;158:1467-71.

- ⁶⁸ Eggen T, Moeder M, Arukwe A. Municipal landfill leachates: A significant source for new and emerging pollutants. *Science of the Total Environment* 2010;408:5147-57.
- ⁶⁹ Huset CA, Barlaz MA, Barofsky DF, Field JA. Quantitative determination of fluorochemicals in municipal landfill leachates. *Chemosphere* 2011;82:1380-6.
- ⁷⁰ Benskin JP, Li B, Ikonomou MG, Grace JR, Li LY. Per- and polyfluoroalkyl substances in landfill leachate: Patterns, time trends, and sources. *Environmental Science & Technology* 2012;46:11532-40.
- ⁷¹ Shoaieoskouei S. Perfluorinated compounds in landfill leachate from discarded carpets. Master of Applied Sciences. The University of British Columbia. June 2012.
- ⁷² Weinberg I, Dreyer A, Ebinghaus R. Landfills as sources of polyfluorinated compounds, polybrominated diphenyl ethers and musk fragrances to ambient air. *Atmospheric Environment* 2011;45:935-41.
- ⁷³ Ahrens L, Shoeib M, Harner T, Lee SC, Guo R, Reiner EJ. Wastewater treatment plant and landfills as sources of polyfluoroalkyl compounds to the atmosphere. *Environmental Science & Technology* 2011;45:8098-105.
- ⁷⁴ Rankin K. Fluorotelomer-based acrylate polymers as an indirect source of perfluoroalkyl carboxylates. Thesis for the degree of Doctor of Philosophy, University of Toronto, 2015.
- ⁷⁵ Huset CA. Determination of fluorochemicals in waste-dominated aqueous systems. Doctor of Philosophy Thesis. Oregon State University. May 2007.
- ⁷⁶ Arvaniti OS and Stasinakis AS. Review on the occurrence, fate and removal of perfluorinated compounds during wastewater treatment. *Science of the Total Environment* 2015;524-525:81-92.
- ⁷⁷ Skutlarek D, Exner M, Färber H. Perfluorinated surfactants in surface and drinking waters. *Environmental Science and Pollution Research* 2006;13(5):299-307.
- ⁷⁸ Jin YH, Liu W, Sato I, Nakayama SF, Sasaki K, Saito N, Tsuda S. PFOS and PFOA in environmental and tap water in China. *Chemosphere* 2009;77:605-11.
- ⁷⁹ Chimeddulam D and Wu KY. River water contaminated with perfluorinated compounds potentially posing the greatest risk to young children. *Chemosphere* 2013;90:1617-24.
- ⁸⁰ Washington JW, Yoo H, Ellington JJ, Jenkins TM, Libelo L. Concentrations, distribution, and persistence of perfluoroalkylates in sludge-applied soils near Decatur, Alabama, USA. *Environmental Science & Technology* 2010;44:8390-6.
- ⁸¹ Yoo H, Washington JW, Ellington JJ, Jenkins TM, Neill MP. Concentrations, distribution, and persistence of fluorotelomer alcohols in sludge-applied soils near Decatur, Alabama, USA. *Environmental Science & Technology* 2010;44:8397-402.
- ⁸² Sepulvado JG, Blaine AC, Hundal LS, Higgins CP. Occurrence and fate of perfluorochemicals in soil following the land application of municipal biosolids. *Environmental Science & Technology* 2011;45:8106-12.
- ⁸³ Skutlarek D, Exner M, Färber H. Perfluorinated surfactants in surface and drinking waters. *Environmental Science and Pollution Research* 2006;13(5):299-307.
- ⁸⁴ Blaine AC, Rich CD, Lakhwinder SH, Lau C, Mills MA, Harris KM, Higgins CP. Uptake of perfluoroalkyl acids into edible crops via land applied biosolids: Field and greenhouse studies. *Environmental Science & Technology* 2013;47:14062-9.